

Protective effect of dietary oils containing omega-3 fatty acids against glucocorticoid-induced osteoporosis*

Amr Samir Elbahnasawy^{1,2†}, Emiliya Ramzievna Valeeva¹, Eman Mustafa El-Sayed² and Natalya Vladimirovna Stepanova¹

¹Department of Bioecology, Hygiene and public health, Institute of Fundamental Medicine and Biology, Kazan Federal University, Kazan, 420012, Russia

²Department of Nutrition and Food Sciences, National Research Centre, Dokki, Giza, 12622, Egypt

ABSTRACT

Purpose: Glucocorticoids (GCs) are implicated in secondary osteoporosis, and the resulting fractures cause significant morbidity. Polyunsaturated fatty acids (PUFAs) play a vital role in bone metabolism. However, few trials have studied the impact of omega-3 PUFA-containing oils against GC-induced osteoporosis. Therefore, the present study was undertaken to determine whether supplementation with omega-3 PUFA-containing dietary oils such as fish oil, flaxseed oil or soybean oil can impede the development of GC-induced osteoporosis. **Methods:** The fatty acids (FAs) content of oils was determined using gas chromatography. Male rats were subdivided into 5 groups (8 rats each): normal control (balanced diet), prednisolone control (10 mg/kg prednisolone daily), soybean oil (prednisolone 10 mg/kg + soybean oil 7% w/w), flaxseed oil (prednisolone 10 mg/kg + flaxseed oil 7% w/w), and fish oil (from cod liver; prednisolone 10 mg/kg + fish oil 7% w/w). **Results:** The study data exhibited a significant depletion in bone mineral density (BMD) and femur mass in the prednisolone control compared to the normal control, accompanied with a marked decrease in the levels of plasma calcium and 1,25-(OH)₂-vitamin D₃, and elevated levels of C-terminal telopeptide (CTX), tumor necrosis factor- α (TNF- α) and malondialdehyde (MDA). Supplementation with fish oil, soybean oil or flaxseed oil helped to improve plasma calcium levels, and suppress oxidative stress and inflammatory markers. Additionally, bone resorption was suppressed as reflected by the decreased CTX levels. However, fish oil was more effective than the other two oils with a significant improvement in BMD and normal histological results compared to the normal control. **Conclusion:** This study demonstrated that supplementation with dietary oils containing omega-3 PUFAs such as fish oil, soybean oil or flaxseed oil can play a role in the prevention of bone loss and in the regulation of bone metabolism, especially fish oil which demonstrated a greater level of protection against GC-induced osteoporosis.

KEY WORDS: glucocorticoid-induced osteoporosis, fish oil, soybean oil, flaxseed oil, inflammation

Introduction

Osteoporosis is recognized by decreased bone density and a deterioration in bone structure, causing a decrease in bone strength and an increase in bone fragility and fracture problems [1]. Approximately 200 million persons worldwide have osteoporosis-related health complications, with more than 9 million fractures being reported annually. About 20% of men and 33% of women over 50 years of age are at high risk of fractures due to osteoporosis

and with the rapid increase in this population worldwide, the incidence of hip fracture will continue to increase significantly in the near future [2]. The reported prevalence of osteoporosis in women was 16% in the United States of America, 15% in France and Germany, 9% in the United Kingdom and 38% in Japan, while in men, the prevalence was 1% in the United Kingdom, 4% in Japan, 3% in Canada, and 8% in France in 2014 [3].

Glucocorticoids (GCs) have been used in the treatment of many diseases for decades to suppress tissue inflammation

Received: June 3, 2019 / Revised: July 10, 2019 / Accepted: July 26, 2019

* This work was supported in cooperation with Kazan Federal University (Russian Federation), National Research Centre (Egypt), the cultural affairs and mission sector, Ministry of Higher Education (Egypt). The work was performed according to the Russian Government Program of Competitive Growth of Kazan Federal University.

† To whom correspondence should be addressed.

tel: +7-9375953532, e-mail: amelbahnasawi@stud.kpfu.ru

© 2019 The Korean Nutrition Society

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

and immunological reactions. Many patients with diseases of the lungs, joints, skin, muscle, eyes, nerves, etc., take prednisolone to control their disease activity and symptoms. A significant part of the population in the following countries (Belgium, Germany, Netherlands, UK, France, Italy, Spain, USA, Canada and Australia) has been reported to be receiving GC therapy [4]. GCs suppress bone formation, increase bone resorption cause negative calcium balance, induce a high risk of fractures and lead to GC-induced osteoporosis [5]. GCs are considered one of the main culprits of secondary osteoporosis which leads to dangerous fractures and causes significant morbidity. After starting of oral GCs, bone loss occurs rapidly, and fracture risk increases [6].

The omega-3 polyunsaturated fatty acids (ω -3 PUFAs) have been reported to have valuable effects in the regulation of bone metabolism and to be positively correlated with bone mineral density (BMD) [7]. The major ω -3 PUFAs include the essential fatty acid (FA), alpha-linolenic acid (ALA), and its longer chain derivatives, eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA), which are important in the control of osteoblastogenesis and inhibition of bone resorption [8]. A recent review of several micronutrients for improving health status among older people identified ω -3 PUFAs as selected micronutrients that might effectively increase the bone formation rate and skeleton functions [9].

Marine fish is a major source of omega-3 PUFA, Hutchins-Wiese et al. [10] found that fish oil can decrease bone resorption and inhibit osteoclastogenesis, thereby decreasing bone resorption. Some plant oils like flaxseed and soybean are rich sources of PUFAs. The essential FA, alpha-linolenic acid (ALA), is usually consumed from different food sources including nuts and seeds (walnuts and chia seeds) and plant-based oils (flaxseed oil, canola oil, and soybean oil) [7]. El-Saeed et al. [11] investigated the role of flaxseed oil in the homeostasis of bones and its effect on changes in the levels of the bone formation markers. They found that flaxseed oil has beneficial impacts on the prevention of osteoporosis. Hassan et al. [12] concluded that consumption of soybean oil-supplemented diet is considered a functional food for preventing and decreasing the problems of osteoporosis in rats.

The lack of effective treatments for the risk of osteoporosis in GC-treated individuals results in a special focus on prevention strategies, such as using diet to protect bone

loss as a side effect for patients being treated with prednisolone. Therefore, the main purpose of the present study was to evaluate the role of soybean, flaxseed, and fish oils in preventing bone loss and osteoporosis in GC-treated rats.

Methods

Fatty acid (FA) analyses of oils

Fish oil (from cod liver) was obtained from Tula pharmaceutical factory, Tula, Russian federation. Soybean and flaxseed oils were purchased from the Food Technology Research Institute, Agriculture Research Center, Giza, Egypt. Samples were stored at -20°C in dark glass bottles until used. The oil samples were esterified and methylated according to the AOCS Official Method Ce 2-66 (AOCS 2009) [13], Methyl esters of FA were determined by gas chromatography using a Perkin Elmer Auto System XL model supplied with flame ionization detector. The compounds were separated on a fused silica capillary column DB-Wax ($60\text{ m} \times 0.32\text{ mm}$ intern diameter, $0.20\text{ }\mu\text{m}$ film thickness). The temperature of the oven was initially kept at 50°C for 5 min, then programmed to increase from 50 to 220°C at a rate of $4^{\circ}\text{C}/\text{min}$. The carrier gas was helium at a flow rate of 1.1 ml/min. The temperatures of the injector and detector were 220 and 250°C , respectively. FAs were identified by comparing the retention times of pure standard methyl esters of FAs with the separate components of the samples, which were quantified by area normalization (%) [14]. A mixture composed of 37 FA methyl esters was used as a standard (Supelco, Sigma-Aldrich Co. LLC, Germany) with a purity between 99.1 and 99.9%.

Experimental animals

Twelve-week-old male Sprague Dawley rats ($135 \pm 15\text{ g}$) were bought from the Laboratory Animal House, National Research Center, Cairo, Egypt. Only male rats were used because of the hormonal fluctuation of female rats during the menstruation cycle, which may have influenced our experiment and results. The rats were housed individually in ventilated cages and maintained in a 12 h light/12 h darkness cycle at $23\text{-}25^{\circ}\text{C}$, with free access to food and deionized water. This study was approved by the ethical committee of the National Research Center, Egypt (approval number: 4/3/15).

Experimental design and diet

The diet was based on the purified balanced AIN-93GM diet, which was formulated for the maintenance of adult rats according to Reeves et al. [15]. After two weeks of acclimatization, the rats were divided into the following 5 groups (8 rats/group): I - (normal control), which was fed a balanced diet with sunflower oil; II - (prednisolone control) which was fed the balanced diet and administrated with prednisolone (10 mg/kg per daily) [16]; III - soybean oil (S), prednisolone (10 mg/kg per daily) + soybean oil (7% w/w in substitution with sunflower oil in the balanced diet); IV - flaxseed oil (X), prednisolone (10 mg/kg per daily) + flaxseed oil (7% w/w) [11]; and V - fish oil (F), prednisolone (10 mg/kg per daily) + fish oil (7% w/w). The experimental period was 3 weeks. Food intake was monitored daily and body weight was recorded weekly during the experimental period.

Blood assays

After the experimental period, the rats were fasted for 12 h and sacrificed by cervical dislocation. Blood samples were collected in heparinized tubes and centrifuged at 2,500 rpm under cooling. The separated plasma was frozen at -80°C for future biochemical analyses. The concentration of malondialdehyde (MDA), an index of lipid peroxidation, was measured according to the method of Draper and Hadley [17]. Tumor necrosis factor- α (TNF- α) was assayed using an enzyme-linked immunosorbent assay (ELISA) kit purchased from RayBio® Rat TNF- α (RayBiotech, Inc., Norcross, Georgia, USA). Calcium (Ca), phosphorus (P) and magnesium (Mg) values were assessed using colorimetric assay kits (BioSystems S.A., Costa Brava, Barcelona, Spain). 1,25-(OH)₂-Vitamin D₃ was quantitatively measured in plasma using an ELISA kit (IBL International GmbH, Hamburg, Germany). The C-terminal telopeptide (CTX) was determined using Plasma Cross Laps One-Step ELISA (Osteometer BioTech, Herlev, Denmark). Parathyroid Hormone (PTH) was determined using an ELISA kit (MyBiosource, Inc., San Diego, USA).

Bone examination

The right femur of each rat was anatomized, all tissues covering the bone were removed, and washed in saline. The femur mass was recorded by electronic balance, and the length was measured using an electronic caliper. The BMD of the femurs was determined using dual-energy

X-ray absorptiometry (DEXA) (Norland XR-600, Swissray International, Inc., Edison, NJ, USA). For histopathological examination, bones were put in 10% formalin for 48 hours. Decalcification was started by ethylene diamine tetra-acetic acid (EDTA) 10% in 7-7.4 pH for 4 weeks. The tissues were cleared with xylol. Femurs were cut transversely. The cleared fixed tissues were transferred into melted paraffin to form blocks of hard paraffin with tissues in the center. Each block of paraffin was cut into 6 μ m-thick sections using a rotatory microtome. Paraffin sections were then transferred to clean slides of glass smeared with glycerin-albumin. The slides were dried and stained by hematoxylin and eosin stain (H & E) for histopathological examination by light microscope. Images were analyzed by Optimas (Media Cybernetics, Inc., USA) 1998 version 6.21.19 [18].

Statistical analysis

The data were statistically analyzed using CoStat Version 6.451 statistical program (CoHort Software, USA). The differences in mean values among all groups were evaluated by one-way completely randomized analysis of variance (ANOVA), followed by Tukey's honestly significant difference (HSD) post hoc test to compare differences in values and determine the significance level ($p < 0.05$). The results are presented as mean \pm standard error (SE).

Results

Fatty acid (FA) profile of oils

As is well known, fish, flaxseed and soybean oils have different compositions of the various FAs. The specific FA profiles in the three study oils were determined and the results are shown in Table 1. Therefore, it was necessary to evaluate which oil is more effective in bone metabolism and in preventing osteoporosis.

Weight gain, food intake, BMD, mass and length of femur

As presented in Table 2, the body weight gain and the daily food intake values showed insignificant changes among the five groups. Femur mineral density, mass and length are also shown in Table 2. Administration of prednisolone in the positive control group significantly reduced femur mass and BMD compared to the negative normal group, and the reduction in femur mass was

Table 1. Fatty acid (FA) profiles of fish, flaxseed and soybean oils

FAs	Concentration (%) of total FAs		
	Fish oil	Flaxseed oil	Soybean oil
Saturated fatty acids (SFA)			
Myristic C14:0	5.19	ND	ND
Palmitic acid C16:0	8.17	8.80	12.19
Stearic acid C18:0	2.05	5.79	4.22
Arachidic acid C20:0	ND	0.26	0.37
Behenic acid, C22:0	ND	ND	0.13
Total SFA	15.41	14.85	16.91
Monounsaturated fatty acids (MUFAs)			
Palmitoleic acid, C16:1n7	6.39	ND	ND
Oleic acid, C18:1n9	12.21	21.40	24.35
Vaccenic acid, C18:1n7	4.14	ND	ND
Eicosenoic acid C20:1n9	8.35	ND	ND
Cetoleic acid, C22:1n11	5.19	ND	ND
Total MUFA	36.28	21.40	24.35
Polyunsaturated fatty acids (PUFAs)			
Linoleic acid, C18:2n6	1.5	22.47	53.58
α -Linolenic acid (ALA) C18:3n3	0.74	40.29	4.67
Stearidonic acid C18:4n3	3.6	ND	ND
Arachidonic acid, C20:4n6	0.25	ND	ND
Eicosapentaenoic acid (EPA) C20:5n3	19.12	ND	ND
Docosapentaenoic acid, (DPA) C22:5n3	1.9	ND	ND
Docosahexaenoic acid (DHA) C22:6n3	21.08	ND	ND
Total ω -6 PUFAs	1.75	22.47	53.58
Total ω -3 PUFAs	46.44	40.29	4.67
Total PUFA	48.19	62.76	58.25

ND: not detected

Table 2. Weight gain, food intake, bone mineral density (BMD), mass and length of femur in the experimental rats

Groups	Weight gain (g)	Daily food intake (g)	BMD (g/cm ²)	Femur mass (g/kg body weight)	Femur length (mm)
Normal control	41 ± 5 ^a	14.1 ± 0.7 ^a	0.1534 ± 0.0075 ^a	3.5 ± 0.19 ^a	31 ± 0.3 ^a
Prednisolone control	36 ± 4 ^a	14.7 ± 1.4 ^a	0.0950 ± 0.0078 ^b	2.6 ± 0.23 ^b	30 ± 0.5 ^a
S	37 ± 4 ^a	15.2 ± 1.2 ^a	0.1157 ± 0.0084 ^b	3.0 ± 0.22 ^{ab}	31 ± 0.5 ^a
X	40 ± 5 ^a	13.9 ± 1.3 ^a	0.1016 ± 0.0061 ^b	3.0 ± 0.18 ^{ab}	31 ± 0.4 ^a
F	47 ± 4 ^a	15.1 ± 0.8 ^a	0.1487 ± 0.0072 ^a	3.2 ± 0.16 ^{ab}	31 ± 0.4 ^a

The data are expressed as mean ± standard error (SE) (n = 8).

Different superscript letters in the same column are significantly different from each other at p < 0.05.

BMD, bone mineral density; S, soybean oil; X, flaxseed oil; F, fish oil

Table 3. Changes in some plasma minerals of the normal control, prednisolone control and treated groups

Groups	Calcium (mg/dl)	Phosphorus (mg/dl)	Magnesium (mg/dl)
Normal control	9.31 ± 0.10 ^a	4.43 ± 0.10 ^a	1.91 ± 0.07 ^a
Prednisolone control	8.01 ± 0.11 ^d	3.21 ± 0.10 ^d	1.89 ± 0.07 ^a
S	8.76 ± 0.11 ^b	3.68 ± 0.10 ^c	1.95 ± 0.09 ^a
X	8.43 ± 0.10 ^c	3.87 ± 0.09 ^{bc}	1.83 ± 0.08 ^a
F	9.03 ± 0.11 ^{ab}	4.08 ± 0.05 ^b	1.79 ± 0.08 ^a

The data are expressed as mean ± standard error (SE) (n = 8).

Different superscript letters in the same column are significantly different from each other at p < 0.05.

S, soybean oil; X, flaxseed oil; F, fish oil

improved insignificantly by the treatment with oils in all treated groups. Furthermore, the treatment with fish oil significantly increased BMD compared to the prednisolone control group. However, the femur length did not change significantly among the five groups.

Calcium, phosphorus and magnesium in blood

Plasma minerals were estimated (Table 3). The prednisolone control group demonstrated a significant decrease in the plasma levels of calcium and phosphorus compared to the normal group, and this decrease was significantly improved by the supplementation of the three oils in the treated groups. The fish oil group had the highest calcium level among all treatments and showed no significant change compared to the normal control, while the plasma magnesium levels in the five groups didn't show any significant differences.

Biomarkers related to bone metabolism, inflammation and oxidative stress

Table 4 shows the inflammatory marker (TNF- α), oxidative stress marker (MDA), and changes in 1,25-(OH) $_2$ -vitamin D $_3$, PTH and CTX. The prednisolone control group recorded the highest values of inflammatory and oxidative stress markers TNF- α and MDA, respectively, while the normal control recorded the lowest values. Treatment with oils significantly improved these values in the treated groups, especially fish oil, which showed a normal value of MDA compared to the normal control group. Supplementation with fish oil also demonstrated the highest value of vitamin D $_3$ with no significant differences from the normal control, while the supplementation with flaxseed oil showed the lowest value. Administration with prednisolone in the control group significantly increased PTH and CTX compared to the normal control, and the

PTH level was decreased after treatment with flaxseed oil and fish oil. Furthermore, the CTX levels in all treated groups were significantly improved compared to the prednisolone control group.

Histological findings of bones

The histological results of femoral examination showed that the normal control (Fig. 1A) and fish oil (Fig. 1E) groups had normal cortical bone thickness and normal osteocyte with no histopathological changes. In contrast, the prednisolone control group (Fig. 1B) clearly demonstrated




Fig. 1. Photomicrographs of bones of rats from (A) the normal control group showing no histopathological changes, normal cortical bone thickness and normal osteocyte; (B) the prednisolone control group showing several resorption cavities within the matrix, abnormal osteocyte, and thin cortical bone with presence of cracks and fissures; (C) the soybean oil group (S) showing few cracks and fissures in the cortical bone; (D) the flaxseed oil group (X) showing focal necrosis with few cracks and fissures in the cortical bone; and (E) the fish oil group (F) showing no histopathological changes, normal bone cortex and normal osteocyte (H & E X 100).

Table 4. Plasma levels of MDA, TNF- α , 1,25-(OH) $_2$ -vitamin D $_3$, PTH and CTX in plasma

Groups	MDA (μ mol/l)	TNF- α (pg/ml)	1,25-(OH) $_2$ -vitamin D $_3$ (pg/ml)	PTH (pg/ml)	CTX (pg/ml)
Normal control	1.45 \pm 0.08 ^c	27.2 \pm 0.8 ^c	59.7 \pm 0.8 ^a	51.5 \pm 0.8 ^c	346 \pm 6 ^d
Prednisolone control	2.33 \pm 0.08 ^a	35.9 \pm 1.2 ^a	49.9 \pm 1.3 ^b	55.6 \pm 0.9 ^{ab}	487 \pm 5 ^a
S	1.94 \pm 0.09 ^b	30.5 \pm 0.9 ^b	52.1 \pm 0.9 ^b	56.8 \pm 0.9 ^a	411 \pm 7 ^b
X	1.89 \pm 0.11 ^b	33.4 \pm 0.8 ^{ab}	48.5 \pm 1.5 ^b	53.1 \pm 0.8 ^{bc}	425 \pm 6 ^b
F	1.41 \pm 0.12 ^c	31.4 \pm 1.3 ^b	61.3 \pm 1.7 ^a	52.7 \pm 1.0 ^c	373 \pm 9 ^c

The data are expressed as mean \pm standard error (SE) (n = 8).

Different superscript letters in the same column are significantly different from each other at p < 0.05.

MDA, malondialdehyde; TNF- α , tumor necrosis factor-alpha; PTH, parathyroid hormone; CTX, C-terminal telopeptide; S, soybean oil; X, flaxseed oil; F, fish oil

a significantly thin cortical bone, accompanied by cracks and fissures, abnormal osteocytes, and several resorption cavities within the matrix. On the other hand, only few cracks and fissures were seen in the cortical bone of the soybean oil group (Fig. 1C). There was some focal necrosis with few cracks and fissures in the cortical bone of rats from the flaxseed oil group (Fig. 1D).

Discussion

GCs are absorbed through the cell membrane by special receptors and affect lipid, protein and carbohydrate metabolisms. The impact of GCs on bone could be linked to the imbalance between bone resorption and bone remodeling, vascular occlusion, and activation of apoptotic signaling pathway by some GCs [19]. Ton et al. [20] suggested that taking even low doses of prednisolone significantly decreases bone formation, may inhibit bone remodeling and could have different effects on bone strength. There are few published recommendations for preventing GC-induced osteoporosis. The prednisolone dosage in the present study was based on data from a previous study on rats [16]. Given that systemic metabolism and the biotransformation of drugs are stronger in rats than in humans, a ten-time transformation rate was adopted in biological experiments conducted on rats. Osseous tissues of rats are distinguished by a fast rate of metabolic turnover. Therefore, the three-week study duration was considered sufficient to evaluate the impact of prednisolone on bone loss in rats [21].

Several studies reported that PUFAs and their metabolites play significant roles in controlling bone metabolism. The link between PUFAs and bone metabolism could be related to some suggested mechanisms such as regulation of the activity and differentiation of the osteoclast and osteoblast, as well as changing the FA composition of bone cell membranes [22]. The omega-3 PUFAs have been reported to suppress osteoclast activity and improve osteoblast activity [23]. Dietary supplementation of long-chain PUFAs demonstrated a protective effect against aging-related bone loss [24].

In our study, we found that the main FA detected in flaxseed oil was ω -3 PUFAs α -linolenic acid (ALA), followed by the monounsaturated oleic acid, and these results are in line with some recent research [25]. Furthermore, the major dietary FA found in soybean oil was ω -6 PUFA (linoleic acid) at more than 50%, and

this result was similar to Dorni et al. [26] who found that ω -6 polyunsaturated linoleic acid was the highest (more than 50%) in soybean oil, followed by the monounsaturated oleic acid, and then the saturated palmitic acid, which together accounted for more than 90% of the total FAs. On the other hand, sunflower oil is also known to be rich in linoleic acid and oleic acid. Fish oil has been reported to be a good source of long-chain ω -3 PUFAs with significant quantities of EPA and DHA that are considered to be metabolically active compounds [27].

In the present work, the femur mass and BMD were reduced in rats with prednisolone-induced osteoporosis. Kaczmarczyk-Sedlak et al. [16] found that administration of prednisolone 10 mg/kg daily for three weeks significantly reduced caused a significant reduction in femur mass by (11.2%). The ratio of femur mass to body mass was lower in rats with prednisolone-induced osteoporosis than in those from the normal control group. This observed decrease in femur mass could be related to bone loss and bone resorption. The negative effect of GCs on the osseous tissue includes both a reduction in BMD and an increase in the number of bone fractures [28]. Pennisi et al. [29] administered prednisolone (5 mg/kg) for 4 weeks to rats and found a reduction in growth and low mineral density of the vertebrae, hip and femur. In our study, feeding rats with PUFAs with slightly different compositions from three different oils increased bone mass in all treated groups, which is in line with Lukas et al. [30] In the current study, only rats fed with fish oil had a significant improvement in BMD compared to the other treated groups. Sun et al. [31] reported that rats fed fish oil, which has high levels of EPA and DHA, had higher BMD and stronger bone than rats fed soybean oil, which is rich in ALA.

Our evaluation of plasma minerals demonstrated that administration with prednisolone significantly decreased plasma calcium and phosphorus compared to the normal group. Some mechanisms proposed in the literature lead to GC-induced bone loss by increasing renal calcium excretion and reducing the absorption of calcium through the intestinal tract [32]. GCs have many indirect negative effects on bones, including disturbance of calcium-phosphorus homeostasis (through blocking intestinal calcium and phosphorus absorption and increasing urinary calcium elimination), reducing growth hormone, and affecting PTH [33]. Therefore, non-pharmacological treatments such as suitable nutrition for preventing bone resorption are

necessary for all adults who are taking prednisolone. Our data showed significantly improved levels of plasma calcium and phosphorus after supplementation with the three oils. Beneficial interactions between calcium and fish oil were observed in the rat model of osteoporosis. These data support the proposed mechanism in which calcium absorption and excretion are regulated after supplementation of fish oil [31]. Omega-3 PUFAs affect bone health by improving calcium balance and bone turnover. The ω -3 PUFAs also influence bone structure by affecting osteoclastic and osteoblastic differentiation and activity [22]. Omega-3 improved plasma calcium levels, and thereby increased bone mass; hence, our data confirm the valuable effect of PUFAs on bone formation. Heaney et al. [34] found that supplementation of ω -3 PUFAs increased calcium absorption in humans. Long-chain PUFAs preserved Ca and P homeostasis and caused no disturbance in their nutritional balance. Soybean oil has been reported to promote bone health through increasing calcium and phosphorus retention [35].

In the current study, the group of rats fed with fish oil had the highest vitamin D₃ levels among the five groups. Fish oil is considered a good source of vitamin D₃ and is recognized as one of the major sources of vitamin D₃ in the human diet [36]. Vitamin D is hydroxylated to 25(OH)₂D₃ in the liver, followed by conversion to its most potent biological metabolite (1,25-(OH)₂D₃) by the enzyme 1 α -hydroxylase in renal proximal and distal convoluted tubules. An et al. [37] demonstrated that ω -3 PUFAs enhance 1,25-dihydroxyvitamin D level in dialysis patients. One hypothesis is that increased 1,25-(OH)₂D₃ concentrations were caused by the enzyme 1 α -hydroxylase activation by omega-3 PUFAs [38].

In addition to the enhanced bone formation, the treated groups showed a reduction in bone loss markers, as reflected by lowered CTX levels in plasma compared to the prednisolone control. Dong et al. [39] surveyed CTX levels in 164 postmenopausal women and found a significant decrease in CTX level in the intervention group after supplementation with ω -3 PUFA compared to the control group. High levels of PTH play a role in the development of bone loss by transferring bone calcium into the blood to restore the normal plasma calcium levels. Consequently, elevated levels of PTH are associated with increasing bone resorption rate, as seen in our results after prednisolone administration. Varela-López et al. [40] also suggested

that PTH is released when plasma calcium levels are low to restore calcium from bone into blood, which induces bone resorption process. On the other hand, fish oil significantly reduced the level of PTH. It has been reported that the protective effect of ω -3 PUFAs against bone loss was due to the modulation of systemic calcitrophic hormones, including PTH [41].

MDA is a secondary substance of lipid peroxidation that is strongly associated with oxidative stress [17]. According to our data, treatment with oils significantly decreased oxidative stress in all treated groups. Han et al. [42] found that flaxseed oil supplementation significantly reduced MDA levels in serum and liver, and suppressed hepatic oxidative stress. The immunological effects of soybean oil and its effect in improving pro-inflammatory status and oxidative stress have been reported [43]. Muga and Chao [44] found that treatment with fish oil reduced lipid peroxidation level to equal or below control levels and suggested that it may prevent or treat lipid peroxidation. Pipingas et al. [45] studied the effect of fish oil on health in 160 healthy individuals and suggested that consumption of daily fish oil prevents oxidative stress. Osteocytes can modulate the formation and function of both osteoblasts and osteoclasts. Apoptotic osteocytes release some pro-inflammatory cytokines such as TNF- α to stimulate osteoclastogenesis. Exposure to GCs can lead to osteocyte apoptosis and then bone loss. Our data showed that the prednisolone group recorded high values of inflammatory marker TNF- α . The GC-induced apoptotic osteocytes could increase the formation of TNF- α [5]. According to our research findings, treatment with oils rich in PUFAs decreased the inflammation. Recent investigation has reported that intake of PUFAs effectively decreased the inflammatory markers [27]. Muga and Chao [44] reported that treatment of fish oil regulated inflammation by decreasing the TNF- α concentration.

Our histological investigation of bones revealed that prednisolone affected bones at the cellular level and induced many abnormalities in the cortex and osteocytes, while treatment with oils, especially fish oil, reduced most of these side effects of prednisolone. According to Kasem et al. [46] histological examination of rat femur after injection of prednisolone to induce osteoporosis showed significantly decreased cortical thickness and osteocyte number of the osteoporosis group when compared with the normal control group. Derakhshanian et al. [47]

recorded a reduction in cortical and trabecular thickness, as well as a significant decrease in the number osteoblasts in GC-treated rats. At the cellular level, GCs can increase apoptosis, inhibit mitosis, and suppress type II collagen synthetic ability in osteoblasts. Moreover, GCs inhibit cartilage growth, suppress synthesis of proteoglycans and decrease osteocyte feasibility, thereby decreasing bone strength [5]. PUFAs suppress the action of osteoclasts and enhance the activity of osteoblasts, thereby inhibiting bone resorption and promoting bone formation. Supplementation of fish oil improved BMD and the mechanical and histological characteristics of bones [48].

Summary

The treatment of osteoporosis remains a considerable problem. Therefore, preventing bone loss is an important demand nowadays. The present data indicated that feeding dietary oils containing omega-3 PUFAs such as fish oil, soybean oil or flaxseed oil could reduce the potential effect of prednisolone in inducing the development of osteoporosis in rats. This may occur through some intracellular pathways, including the improvement of calcium absorption, regulation of bone metabolism, differentiation of the osteoblast and osteoclast, suppression of oxidative stress and modulation of inflammatory response. Thus, these oils and fish oil, in particular, could be used as a natural approach to help prevent the bone loss associated with GC therapy, and this potential offers an interesting prospect for future research.

ORCID

Amr Samir Elbahnasawy: <https://orcid.org/0000-0003-1012-478X>

Emiliya Ramzievna Valeeva: <https://orcid.org/0000-0002-8683-6241>

Eman Mustafa El-Sayed: <https://orcid.org/0000-0001-5825-8088>

Natalya Vladimirovna Stepanova: <https://orcid.org/0000-0002-2889-3109>

References

- Jeremiah MP, Unwin BK, Greenawald MH, Casiano VE. Diagnosis and management of osteoporosis. *Am Fam Physician* 2015; 92(4): 261-268.
- Akarimak U. Osteoporosis: a major problem - worldwide. *Arch Sports Med* 2018; 2(1): 106-108.
- Wade SW, Strader C, Fitzpatrick LA, Anthony MS, O'Malley CD. Estimating prevalence of osteoporosis: examples from industrialized countries. *Arch Osteoporos* 2014; 9(1): 182.
- Silverman S, Curtis J, Saag K, Flahive J, Adachi J, Anderson F, et al. International management of bone health in glucocorticoid-exposed individuals in the observational GLOW study. *Osteoporos Int* 2015; 26(1): 419-420.
- Wang T, Yu X, He C. Pro-inflammatory cytokines: cellular and molecular drug targets for glucocorticoid-induced-osteoporosis via osteocyte. *Curr Drug Targets* 2019; 20(1): 1-15.
- Compston J. Glucocorticoid-induced osteoporosis: an update. *Endocrine* 2018; 61(1): 7-16.
- Lavado-García J, Roncero-Martín R, Moran JM, Pedrera-Canal M, Aliaga I, Leal-Hernandez O, et al. Long-chain omega-3 polyunsaturated fatty acid dietary intake is positively associated with bone mineral density in normal and osteopenic Spanish women. *PLoS One* 2018; 13(1): e0190539.
- Kelly OJ, Gilman JC, Kim Y, Ilich JZ. Long-chain polyunsaturated fatty acids may mutually benefit both obesity and osteoporosis. *Nutr Res* 2013; 33(7): 521-533.
- Iolascon G, Gimigliano R, Bianco M, De Sire A, Moretti A, Giusti A, et al. Are dietary supplements and nutraceuticals effective for musculoskeletal health and cognitive function? A scoping review. *J Nutr Health Aging* 2017; 21(5): 527-538.
- Hutchins-Wiese HL, Picho K, Watkins BA, Li Y, Tannenbaum S, Claffey K, et al. High-dose eicosapentaenoic acid and docosahexaenoic acid supplementation reduces bone resorption in postmenopausal breast cancer survivors on aromatase inhibitors: a pilot study. *Nutr Cancer* 2014; 66(1): 68-76.
- El-Saeed GS, Elghoroury EA, Morsy S, Aly HM, Wafaey H. Phenotype of vitamin D receptor gene polymorphisms, impact of feeding flaxseed oil, and osteoporosis in ovariectomized diabetic rats. *Bull Natl Res Cent* 2018; 42(1): 11.
- Hassan HA, El Wakf AM, El Gharib NE. Role of phytoestrogenic oils in alleviating osteoporosis associated with ovariectomy in rats. *Cytotechnology* 2013; 65(4): 609-619.
- American Oil Chemists' Society (AOCS). Official methods and recommended practices of the AOCS. Champaign (IL): AOCS; 2009.
- Veronezi CM, Costa T, Jorge N. Basil (*Ocimum Basilicum* L.) as a natural antioxidant. *J Food Process Preserv* 2014; 38(1): 255-261.
- Reeves PG, Nielsen FH, Fahey GC Jr. AIN-93 purified diets for laboratory rodents: final report of the American Institute of Nutrition ad hoc writing committee on the reformulation of the AIN-76A rodent diet. *J Nutr* 1993; 123(11): 1939-1951.
- Kaczmarczyk-Sedlak I, Zych M, Rotko K, Sedlak L. Effects of thalidomide on the development of bone damage caused by prednisolone in rats. *Pharmacol Rep* 2012; 64(2): 386-395.
- Draper HH, Hadley M. Malondialdehyde determination as index of lipid peroxidation. *Methods Enzymol* 1990; 186: 421-431.
- Bancroft JD, Gamble M. Theory and practice of histological techniques. 5th edition. London: Churchill Livingstone; 2002.
- Le HT, Phi LT, Dao TT, Phan NK, Van Pham P, Vu NB.

- A mouse model of osteonecrotic femoral head induced by methylprednisolone and lipopolysaccharide. *Biomedical Research and Therapy* 2016; 3(3): 548-556.
20. Ton FN, Gunawardene SC, Lee H, Neer RM. Effects of low-dose prednisone on bone metabolism. *J Bone Miner Res* 2005; 20(3): 464-470.
 21. Kaczmarczyk-Sedlak I, Cegiela U, Nowińska B, Folwarczna J. Effects of catecholamines on the intramedullary pressure in the femur in rats with prednisolone-induced osteoporosis. *Pharmacol Rep* 2006; 58(4): 540-550.
 22. Lau BY, Cohen DJ, Ward WE, Ma DW. Investigating the role of polyunsaturated fatty acids in bone development using animal models. *Molecules* 2013; 18(11): 14203-14227.
 23. Watkins BA, Li Y, Lippman HE, Feng S. Modulatory effect of omega-3 polyunsaturated fatty acids on osteoblast function and bone metabolism. *Prostaglandins Leukot Essent Fatty Acids* 2003; 68(6): 387-398.
 24. Bullon P, Battino M, Varela-Lopez A, Perez-Lopez P, Granados-Principal S, Ramirez-Tortosa MC, et al. Diets based on virgin olive oil or fish oil but not on sunflower oil prevent age-related alveolar bone resorption by mitochondrial-related mechanisms. *PLoS One* 2013; 8(9): e74234.
 25. Antonacci LE, Bussetti M, Rodriguez MA, Cano AV, Gagliostro GA. Effect of diet supplementation with combinations of soybean and linseed oils on milk production and fatty acid profile in lactating dairy ewes. *Agric Sci* 2018; 9(2): 200-220.
 26. Dorni C, Sharma P, Saikia G, Longvah T. Fatty acid profile of edible oils and fats consumed in India. *Food Chem* 2018; 238: 9-15.
 27. Calder PC. Marine omega-3 fatty acids and inflammatory processes: Effects, mechanisms and clinical relevance. *Biochim Biophys Acta* 2015; 1851(4): 469-484.
 28. Shah SK1, Gecys GT. Prednisone-induced osteoporosis: an overlooked and undertreated adverse effect. *J Am Osteopath Assoc* 2006; 106(11): 653-657.
 29. Pennisi P, D'Alcamo MA, Leonetti C, Clementi A, Cutuli VM, Riccobene S, et al. Supplementation of L-arginine prevents glucocorticoid-induced reduction of bone growth and bone turnover abnormalities in a growing rat model. *J Bone Miner Metab* 2005; 23(2): 134-139.
 30. Lukas R, Gigliotti JC, Smith BJ, Altman S, Tou JC. Consumption of different sources of omega-3 polyunsaturated fatty acids by growing female rats affects long bone mass and microarchitecture. *Bone* 2011; 49(3): 455-462.
 31. Sun L, Tamaki H, Ishimaru T, Teruya T, Ohta Y, Katsuyama N, et al. Inhibition of osteoporosis due to restricted food intake by the fish oils DHA and EPA and perilla oil in the rat. *Biosci Biotechnol Biochem* 2004; 68(12): 2613-2615.
 32. Mazziotti G, Formenti AM, Adler RA, Bilezikian JP, Grossman A, Sbardella E, et al. Glucocorticoid-induced osteoporosis: pathophysiological role of GH/IGF-I and PTH/VITAMIN D axes, treatment options and guidelines. *Endocrine* 2016; 54(3): 603-611.
 33. Suarez-Bregua P, Guerreiro PM, Rotllant J. Stress, glucocorticoids and bone: a review from mammals and fish. *Front Endocrinol (Lausanne)* 2018; 9: 526.
 34. Heaney RP, Carey R, Harkness L. Roles of vitamin D, n-3 polyunsaturated fatty acid, and soy isoflavones in bone health. *J Am Diet Assoc* 2005; 105(11): 1700-1702.
 35. Maditz KH, Smith BJ, Miller M, Oldaker C, Tou JC. Feeding soy protein isolate and oils rich in omega-3 polyunsaturated fatty acids affected mineral balance, but not bone in a rat model of autosomal recessive polycystic kidney disease. *BMC Nephrol* 2015; 16(1): 13.
 36. Jaafarnejad F, Hosseini SF, Mazlom SR, Hami M. Comparison of the effect of fish oil and vitamin E on the duration of Cyclic mastalgia. *Evid Based Care J* 2013; 3(1): 69-76.
 37. An WS, Lee SM, Son YK, Kim SE, Kim KH, Han JY, et al. Omega-3 fatty acid supplementation increases 1,25-dihydroxyvitamin D and fetuin-A levels in dialysis patients. *Nutr Res* 2012; 32(7): 495-502.
 38. Lee SM, Son YK, Kim SE, An WS. The effects of omega-3 fatty acid on vitamin D activation in hemodialysis patients: a pilot study. *Mar Drugs* 2015; 13(2): 741-755.
 39. Dong H, Hutchins-Wiese H, Kleppinger A, Annis K, Liva E, Lammi-Keefe C, et al. Effects of omega-3 polyunsaturated fatty acid supplementation on bone turnover in older women. *Int J Vitam Nutr Res* 2014; 84(3-4): 124-132.
 40. Varela-López A, Ochoa JJ, Llamas-Elvira JM, López-Frías M, Planells E, Speranza L, et al. Loss of bone mineral density associated with age in male rats fed on sunflower oil is avoided by virgin olive oil intake or coenzyme Q supplementation. *Int J Mol Sci* 2017; 18(7): 1397.
 41. Chen AB, Minami K, Raposo JF, Matsuura N, Koizumi M, Yokota H, et al. Transient modulation of calcium and parathyroid hormone stimulates bone formation. *Endocrine* 2016; 54(1): 232-240.
 42. Han H, Qiu F, Zhao H, Tang H, Li X, Shi D. Dietary flaxseed oil prevents western-type diet-induced nonalcoholic fatty liver disease in apolipoprotein-E knockout mice. *Oxid Med Cell Longev* 2017; 2017: 3256241.
 43. Klein K, Gay S. Epigenetics in rheumatoid arthritis. *Curr Opin Rheumatol* 2015; 27(1): 76-82.
 44. Muga MA, Chao JC. Effects of fish oil and spirulina on oxidative stress and inflammation in hypercholesterolemic hamsters. *BMC Complement Altern Med* 2014; 14(1): 470.
 45. Pipingas A, Sinclair A, Croft KD, Januszewski AS, Jenkins AJ, Mori TA, et al. Fish oil and multivitamin supplementation reduces oxidative stress but not inflammation in healthy older adults: a randomised controlled trial. *J Funct Foods* 2015; 19: 949-957.
 46. Kasem MA, Khedr ES, Abdel-Aleem AM, Said AS. Histological effect of bisphosphonate, vitamin D and olive oil on glucocorticoid induced osteoporosis (Gio) in Albino Rat. *Egypt J Hosp Med* 2016; 65: 699-708.
 47. Derakhshanian H, Djalali M, Djazayeri A, Nourijelyani K, Ghadbeigi S, Pishva H, et al. Quercetin prevents experimental glucocorticoid-induced osteoporosis: a comparative study with alendronate. *Can J Physiol Pharmacol* 2013; 91(5): 380-385.
 48. Liu D, Veit HP, Wilson JH, Denbow DM. Long-term supplementation of various dietary lipids alters bone mineral content, mechanical properties and histological characteristics of Japanese quail. *Poult Sci* 2003; 82(5): 831-839.