

Increasing Prevalence of Type 2 Diabetes in a Rural Bangladeshi Population: A Population Based Study for 10 Years

Bishwajit Bhowmik¹, Faria Afsana², Lien My Diep^{1,3}, Sanjida Binte Munir⁴, Erica Wright⁵, Sharif Mahmood⁶, A. K. Azad Khan², Akhtar Hussain¹

¹Department of Community Medicine, Institute of Health and Society, University of Oslo Faculty of Medicine, Oslo, Norway,

²Department of Endocrinology, Bangladesh Institute of Research and Rehabilitation in Diabetes, Endocrine and Metabolic Disorders (BIRDEM), Dhaka, Bangladesh,

³Unit for Biostatistics and Epidemiology, Oslo University Hospital, Oslo, Norway,

⁴Executive Diabetes Care Centre, NHN, Diabetic Association of Bangladesh, Dhaka, Bangladesh,

⁵ACT Diabetes Service, Division of Medicine, Canberra Hospital, Canberra, Australia,

⁶Institutes of Statistical Research and Training, University of Dhaka, Dhaka, Bangladesh

Diabetes Metab J 2013;37:46-53. <http://dx.doi.org/10.4093/dmj.2013.37.1.46>

With regard to the paper by Bhowmik et al. (Diabetes Metab J 2013;37:46-53), the statement that the protocol was approved by the Norwegian Ethical Committee is incorrect. The authors deeply apologize for the imprecise statement. All the subjects were from Bangladesh and ethical clearance was approved only by the Ethical Committee from Bangladesh. All necessary information was collected from the participants following participant consent. The research was conducted following Declaration of Helsinki.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.