



Corrigendum

Korean J Anesthesiol 2020;73(3):268
<https://doi.org/10.4097/kja.19497.e1>
pISSN 2005-6419 • eISSN 2005-7563

Tips for troublesome sample-size calculation

Junyong In¹, Hyun Kang², Jong Hae Kim³, Tae Kyun Kim⁴, Eun Jin Ahn⁵, Dong Kyu Lee⁶, Sangseok Lee⁷, Jae Hong Park⁸

Department of Anesthesiology and Pain Medicine, ¹Dongguk University Ilsan Hospital, Goyang, ²Chung-Ang University College of Medicine, Seoul, ³Daegu Catholic University School of Medicine, Daegu, ⁴Yongsan Hospital, Pusan National University School of Medicine, Busan, ⁵Inje University Seoul Paik Hospital, Inje University College of Medicine, ⁶Guro Hospital, Korea University School of Medicine, ⁷Sanggye Paik Hospital, Inje University College of Medicine, Seoul, ⁸Inje University Haeundae Paik Hospital, Inje University College of Medicine, Busan, Korea

Korean Journal of Anesthesiology 2020; 73(2):114-120
<https://doi.org/10.4097/kja.19497>

The article by In J, et al. entitled, “Tips for troublesome sample-size calculation.” contained an error in the Equation 12.

Before correction:

$$\text{Adjusted sample size} = \frac{\text{Calculated sample size}}{1 - \text{dropout rate}} = \frac{39}{1 - 0.9} = 43.33$$

(Equation 12)

The correct information is found below:

$$\text{Adjusted sample size} = \frac{\text{Calculated sample size}}{1 - \text{dropout rate}} = \frac{39}{1 - 0.1} = 43.33$$

(Equation 12)

The value for dropout rate was incorrectly printed as 0.9. The value for dropout rate should be corrected to 0.1.

The authors apologize for any inconvenience this mistake may have caused.

© The Korean Society of Anesthesiologists, 2020

© This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.