



Corrigendum

Korean J Anesthesiol 2020;73(6):572
<https://doi.org/10.4097/kja.d.18.00242.e1>
pISSN 2005-6419 • eISSN 2005-7563

What is the proper way to apply the multiple comparison test?

Sangseok Lee¹, Dong Kyu Lee²

¹Department of Anesthesiology and Pain Medicine, Sanggye Paik Hospital, Inje University College of Medicine, ²Department of Anesthesiology and Pain Medicine, Guro Hospital, Korea University School of Medicine, Seoul, Korea

Korean Journal of Anesthesiology 2018;71(5):353-360.
<https://doi.org/10.4097/kja.d.18.00242>

The article Lee S and Lee DK, "What is the proper way to apply the multiple comparison test?" contained an error in the first appeared equation on page 353.

Before correction:

$$\mu_A \neq \mu_B \neq \mu_C \text{ or } \mu_A \neq \mu_B = \mu_C \text{ or } \mu_A = \mu_B \neq \mu_C \text{ or } \mu_A \neq \mu_C = \mu_B$$

The correct information is found below:

$$\mu_A \neq \mu_B \neq \mu_C \text{ or } \mu_A \neq \mu_B = \mu_C \text{ or } \mu_A = \mu_B \neq \mu_C \text{ or } \mu_A = \mu_C \neq \mu_B$$

The authors would like to apologize for any inconvenience caused.

© The Korean Society of Anesthesiologists, 2020

© This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.