



Power spectrum density analysis for the influence of complete denture on the brain function of edentulous patients - pilot study

Praveen Perumal, Gopi Naveen Chander*, Kuttae Viswanathan Anitha, Jetti Ramesh Reddy, Balasubramaniam Muthukumar

Department of Prosthodontics, SRM Dental College, Ramapuram, Chennai, India

PURPOSE. This pilot study was to find the influence of complete denture on the brain activity and cognitive function of edentulous patients measured through Electroencephalogram (EEG) signals. **MATERIALS AND METHODS.** The study recruited 20 patients aged from 50 to 60 years requiring complete dentures with inclusion and exclusion criteria. The brain function and cognitive function were analyzed with a mental state questionnaire and a 15-minute analysis of power spectral density of EEG alpha waves. The analysis included edentulous phase and post denture insertion adaptive phase, each done before and after chewing. The results obtained were statistically evaluated. **RESULTS.** Power Spectral Density (PSD) values increased from edentulous phase to post denture insertion adaption phase. The data were grouped as edentulous phase before chewing (EEG p1-0.0064), edentulous phase after chewing (EEG p2-0.0073), post denture insertion adaptive phase before chewing (EEG p3-0.0077), and post denture insertion adaptive phase after chewing (EEG p4-0.0096). The acquired values were statistically analyzed using paired t-test, which showed statistically significant results ($P < .05$). **CONCLUSION.** This pilot study showed functional improvement in brain function of edentulous patients with complete dentures rehabilitation. [*J Adv Prosthodont 2016;8:187-93*]

KEY WORDS: Power Spectral Density; Fast Fourier Transform; Brain activity; Geriatric; Electroencephalography; Complete denture

INTRODUCTION

World Health Organization (WHO) and the National Institute on Ageing (NIA) reported that tooth loss could be a risk for Alzheimer's disease.¹⁻⁴ The basal ganglia, limbic system, thalamus, and cerebral cortex control the mastication and are linked with the centers of mastication, deglutition, and respiration.^{5,6} Inhibition of masticatory movement

linked with the mastication center and deglutition center can cause a sudden deterioration of brain function.⁷⁻⁹ Masticatory efficacy could be improved by several ways in edentulous patients, from conventional complete denture to implant supported prosthesis.^{10,11} The effect of complete denture on restoring patients' mastication has been sufficiently evaluated by different methods, but the role of balanced complete dentures in the functional improvement of brain function has not been elucidated. This pilot study aims to know complete denture's influence on the brain and cognitive functions in edentulous patients.

MATERIALS AND METHODS

The protocol of the study was approved by the Institutional Ethical committee of SRM University, Chennai, India (SRMU/M &HS/SRMDC/2012/MDS-PG student /205). The patients of age group between 50 and 60 years were informed of the procedure and their consents were obtained (Fig. 1). The patients selected were edentulous for

Corresponding author:
Gopi Naveen Chander
Department of Prosthodontics, SRM Dental College, #496, 3rd Main Road, TNHB Colony, Velachery, Chennai -42, India
Tel. 919840749441; e-mail, drgopichander@gmail.com
Received September 12, 2015 / Last Revision February 1, 2016 / Accepted March 24, 2016

© 2016 The Korean Academy of Prosthodontics
This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.




Fig. 1. Study procedure flow chart.

a year and they were treated with complete denture for the first time. Patients participated in this study had class 1 ridges and adequate interarch spaces. A comprehensive examination excluded patients with symptoms of temporomandibular disorders, xerostomia, orofacial motor disorders, severe oral manifestations of systemic diseases, and psychological or psychiatric conditions that could influence their response to treatment.

The electroencephalogram (EEG) signals were used to check the brain activity.¹² Patients were instructed to take adequate sleep the night before and were restricted from consuming caffeine at least eight hours before the EEG analysis. The electrode placement for EEG was done in accordance to the international federation in electroencephalography and clinical neurophysiology as a 10-20 electrode placement system.^{13,14} EEG recordings were obtained in two settings: before chewing paraffin gum (Saliva-Check kit, GC Corporation, Japan) and after three minutes of chewing gum with one-minute resting intervals between each EEG recording. The recordings were made for the total of fifteen minutes. The signals were sampled at 256 samples per second with 16-bit resolution and with fifteen minutes of digitized signal duration. This data was saved to be analyzed and compared with the EEG records two months after denture placement (Fig. 2, Fig. 3 and Fig. 4).

Complete dentures were fabricated with balanced occlusion for all the patients with maximized comfort. After two months of denture usage, the EEG recordings made before and after chewing were similar to pre-treatment EEG results. Both the pre-treatment and post-treatment EEG data were analyzed for differences in power spectral density (PSD) with gum chewing.




Fig. 2. Flow chart for EEG measurement.




Fig. 3. Data acquisition of EEG.




Fig. 4. Flow Chart for EEG Analysis.

The algorithm was tested using a single channel (C_3-P_3). The alpha waves of the greatest amplitude were recorded. The data was pre-processed by computing average spectrum with 50% overlap in the epoch signal. EEG signal baseline wanders were viewed in the LAB VIEW platform (National Instrument, India). The recordings were corrected for artifacts and the signal amplitude was quantified to micro-volts.¹⁵ A digital low pass finite element filter (FIR) filtered the EEG signal by Hamming window technique. The order of the filter was 40 and the cut off frequency was 32 Hz. Flatness without a ripple in the pass band was desirable in the analysis of EEG signals. The filtered EEG segments were chosen for analysis.

Averaged spectrum and power measure were calculated. The power spectrum of the signal was computed using Fast Fourier Transformation (FFT). The equation for FFT is given as a formula¹⁶:

$$X(k) = \sum_{n=0}^{N-1} x(n)W_N^{kn} : k=0, \dots, N-1$$

$$W_N = e^{-j\frac{2\pi}{N}}$$

where one value of 'k' has N complex multiplications, since 'k' = 0, 1, ..., N-1. The multiplication of x(n) and w^{kn} was done for N times, since n = 0 to N-1.

Spectral analysis is the function of power over frequency. In medicine, a spectral analysis of various signals measured from a patient, such as electrocardiogram (ECG) or electroencephalogram (EEG) signal, provides useful information for diagnosis. A random signal usually has finite average power and is characterized by an average power spectral density as in equation:

$$PSD_f(w) = \lim_{T \rightarrow \infty} \frac{|F_{X_T}(w)|^2}{2T}$$

where $F(w)$ XT represents the FFT output and T is the total duration of the input signal (Fig. 5). The power spec-

tral density of alpha waves calculated from the data was obtained (Table 1). The Kolmogorov-Smirnov test was used to check the normal distribution of the values and the results were statistically analyzed using a paired t-test.




Fig. 5. Power spectral density value.

Table 1. Power spectral density (relative units)

| S. NO: | EEG P1 | EEG P2 | EEG P3 | EEG P4 |
|--------|--------|--------|--------|--------|
| 1 | 0.0025 | 0.0034 | 0.0038 | 0.0053 |
| 2 | 0.0039 | 0.0042 | 0.0051 | 0.0074 |
| 3 | 0.0095 | 0.0107 | 0.0131 | 0.0188 |
| 4 | 0.0039 | 0.0042 | 0.0057 | 0.0061 |
| 5 | 0.0133 | 0.0141 | 0.0133 | 0.0154 |
| 6 | 0.0062 | 0.0078 | 0.0099 | 0.0121 |
| 7 | 0.0021 | 0.0023 | 0.0039 | 0.0042 |
| 8 | 0.0058 | 0.0119 | 0.0085 | 0.0123 |
| 9 | 0.0012 | 0.0067 | 0.0042 | 0.0080 |
| 10 | 0.0016 | 0.0025 | 0.0026 | 0.0054 |
| 11 | 0.0121 | 0.0131 | 0.0167 | 0.0170 |
| 12 | 0.0049 | 0.0049 | 0.0032 | 0.0046 |
| 13 | 0.0026 | 0.0013 | 0.0018 | 0.0032 |
| 14 | 0.0133 | 0.0143 | 0.0133 | 0.0178 |
| 15 | 0.0039 | 0.0045 | 0.0042 | 0.0055 |
| 16 | 0.0082 | 0.0093 | 0.0095 | 0.0096 |
| 17 | 0.0056 | 0.0058 | 0.0067 | 0.0069 |
| 18 | 0.0124 | 0.0135 | 0.0114 | 0.0144 |
| 19 | 0.0067 | 0.0074 | 0.0087 | 0.0089 |
| 20 | 0.0075 | 0.0076 | 0.0078 | 0.0083 |

EEG p1: Edentulous phase before chewing, EEG p2: Edentulous phase after chewing, EEG p3: Post denture insertion adaptive phase before chewing, EEG p4: Post denture insertion adaptive phase after chewing.

RESULT

The mean values were grouped as edentulous phase before chewing (EEG p1-0.006), edentulous phase after chewing (EEG p2-0.007), post denture insertion adaptive phase before chewing (EEG p3-0.007), and post denture insertion adaptive phase after chewing (EEG p4-0.009). The values were statistically analyzed using a paired t-test, which showed a significant increase in alpha waves PSD values after chewing in both edentulous and post denture adaptive phase. The alpha waves PSD related to before-chewing values increased in post denture adaptive phase ($P < .05$). (Table 2, Fig. 6)




Fig. 6. Mean values of edentulous phase and post denture insertion adaptive phase before and after chewing.

DISCUSSION

The sensory information transmitted through the trigeminal nerve from the masticatory muscles, temporomandibular joint, and residual mucous membrane increased when subjects chewed. This transmission was unique and different from the control mechanisms involved in the movements of the arms and legs. Otsuka *et al.*¹⁶ proved the relationship between malocclusion and the limbic circuits. Narita *et al.*¹⁷ and Hirano *et al.*¹⁸ estimated that gum chewing was more effective in stimulating brain activity than other treatments. Yamazaki *et al.*¹⁹ demonstrated that, in both mature and old animals, the spatial memory declined as the number of hippocampal neurons decreased from impairment in hippocampal function due to early tooth loss. Hippocampal dysfunction caused impairment in episodic memory such as learning new information and retrieving information.²⁰ Like hippocampus, other brain areas, such as the prefrontal cortex (PFC), the striatum, and the cerebellum, play an important role in executive abilities like performing multiple tasks simultaneously, set-shifting, and inhibition. These areas are sensitive to ageing.^{21,22} Morokuma²³ and Klimesch²⁴ showed a positive relationship between alpha waves and cognitive performance as well as the speed of processing information. With this concept, brain activity was evaluated by studying power spectral density of alpha waves in edentulous patients before and after denture insertion. PSD values were calculated and a comparison was made for the EEG taken in two phases - edentulous phase and post denture insertion adaptive phase - before and after chewing paraffin gum.

The power spectral density of alpha waves increased in both edentulous phase and post denture adaptive phase after chewing. The masticatory stimulation travels from the masticatory muscles to the hypothalamus via trigeminal nerve. This mechanism is believed to involve a wide area of the brain and gum chewing is thought to be effective in

Table 2. Paired samples t-test to compare the mean value

| Pair | Variables | N | Mean | SD | t value | P value |
|--------|-----------|----|---------|--------|---------|---------|
| Pair 1 | EEG p1 | 20 | 0.00641 | 0.0039 | 3.566 | .002 |
| | EEG p3 | 20 | 0.0077 | 0.0042 | | |
| Pair 2 | EEG p2 | 20 | 0.0073 | 0.0039 | 4.974 | < .001 |
| | EEG p4 | 20 | 0.0096 | 0.0048 | | |
| Pair 3 | EEG p1 | 20 | 0.0064 | 0.0039 | 3.671 | .002 |
| | EEG p2 | 20 | 0.0073 | 0.0039 | | |
| Pair 4 | EEG p3 | 20 | 0.0077 | 0.0042 | 5.202 | < .001 |
| | EEG p4 | 20 | 0.0096 | 0.0048 | | |

EEG p1: Edentulous phase before chewing, EEG p2: Edentulous phase after chewing, EEG p3: Post denture insertion adaptive phase before chewing, EEG p4: Post denture insertion adaptive phase after chewing.

stimulating the brain activity.²⁵ The study result did not synchronize with the studies performed by Masumoto *et al.*,²⁶ which stated no significant differences in EEG frequencies between a control and post gum condition. Masumoto *et al.*²⁷ contradicted their earlier study with the findings indicating that chewing flavorless and odorless gum base led to increased alpha and theta activity.

Comparison of PSD values before chewing between edentulous phase and post denture insertion adaption phase showed that post denture insertion adaptive phase PSD value increased and influenced the brain function. The results concluded that dentures showed eminent improvement in brain function.

The limitation of this study is that the brain function can potentially be affected by other factors involving the subject, including his/her living environment, hospital visits, conversations with the attending physician, or the treatment administered. There is also a concern over the potential effect of circadian variability of EEG on the measurement sequence of brain function. These factors have to be standardized in future studies.

CONCLUSION

Within the limitations of the study, it can be concluded that the brain activity of edentulous patients improved with mastication and denture reconstruction.

ORCID

Gopi Naveen Chander <http://orcid.org/0000-0002-2040-4550>

REFERENCES

- Musha T, Matsuzaki H. Neuronal impairment and instability. *Int J Bioelectromagn* 2007;9:113-5.
- Onozuka M, Watanabe K, Nagasaki S, Jiang Y, Ozono S, Nishiyama K, Kawase T, Karasawa N, Nagatsu I. Impairment of spatial memory and changes in astroglial responsiveness following loss of molar teeth in aged SAMP8 mice. *Behav Brain Res* 2000;108:145-55.
- Shalat SL, Seltzer B, Pidcock C, Baker EL Jr. Risk factors for Alzheimer's disease: a case-control study. *Neurology* 1987;37:1630-3.
- Weiner MW, Veitch DP, Aisen PS, Beckett LA, Cairns NJ, Cedarbaum J, Green RC, Harvey D, Jack CR, Jagust W, Luthman J, Morris JC, Petersen RC, Saykin AJ, Shaw L, Shen L, Schwarz A, Toga AW, Trojanowski JQ; Alzheimer's Disease Neuroimaging Initiative. 2014 Update of the Alzheimer's Disease Neuroimaging Initiative: A review of papers published since its inception. *Alzheimers Dement* 2015;11:e1-120.
- Frota de Almeida MN, de Siqueira Mendes Fde C, Gurgel Felício AP, Falsoni M, Ferreira de Andrade ML, Bento-Torres J, da Costa Vasconcelos PF, Perry VH, Picanço-Diniz CW, Kronka Sosthenes MC. Spatial memory decline after masticatory deprivation and aging is associated with altered laminar distribution of CA1 astrocytes. *BMC Neurosci* 2012;13:23.
- Soboļeva U, Lauriņa L, Slaidiņa A. The masticatory system—an overview. *Stomatologija* 2005;7:77-80.
- Avivi-Arber L. Neuroplasticity and the edentulous patient—toward a paradigm shift in oral rehabilitation. *Int J Prosthodont* 2015;28:115.
- De Cicco V. Central syntropic effects elicited by trigeminal proprioceptive equilibrium in Alzheimer's disease: a case report. *J Med Case Rep* 2012;6:161.
- Teixeira FB, Pereira Fernandes Lde M, Noronha PA, dos Santos MA, Gomes-Leal W, Ferraz Maia Cdo S, Lima RR. Masticatory deficiency as a risk factor for cognitive dysfunction. *Int J Med Sci* 2014;11:209-14.
- Hirano Y, Obata T, Takahashi H, Tachibana A, Kuroiwa D, Takahashi T, Ikehira H, Onozuka M. Effects of chewing on cognitive processing speed. *Brain Cogn* 2013;81:376-81.
- Stellingsma K, Slagter AP, Stegenga B, Raghoobar GM, Meijer HJ. Masticatory function in patients with an extremely resorbed mandible restored with mandibular implant-retained overdentures: comparison of three types of treatment protocols. *J Oral Rehabil* 2005;32:403-10.
- Dauwels J, Vialatte F, Latchoumane C, Jeong J, Cichocki A. EEG synchrony analysis for early diagnosis of Alzheimer's disease: a study with several synchrony measures and EEG data sets. *Conf Proc IEEE Eng Med Biol Soc* 2009;2009:2224-7.
- Homan RW, Herman J, Purdy P. Cerebral location of international 10-20 system electrode placement. *Electroencephalogr Clin Neurophysiol* 1987;66:376-82.
- Atcherson SR, Gould HJ, Pousson MA, Prout TM. Variability of electrode positions using electrode caps. *Brain Topogr* 2007;20:105-11.
- Stewart CM, Newlands SD, Perachio AA. Spike detection, characterization, and discrimination using feature analysis software written in LabVIEW. *Comput Methods Programs Biomed* 2004;76:239-51.
- Otsuka T, Watanabe K, Hirano Y, Kubo K, Miyake S, Sato S, Sasaguri K. Effects of mandibular deviation on brain activation during clenching: an fMRI preliminary study. *Cranio* 2009;27:88-93.
- Narita N, Kamiya K, Yamamura K, Kawasaki S, Matsumoto T, Tanaka N. Chewing-related prefrontal cortex activation while wearing partial denture prosthesis: pilot study. *J Prosthodont Res* 2009;53:126-35.
- Hirano Y, Obata T, Takahashi H, Tachibana A, Kuroiwa D, Takahashi T, Ikehira H, Onozuka M. Effects of chewing on cognitive processing speed. *Brain Cogn* 2013;81:376-81.
- Yamazaki K, Wakabayashi N, Kobayashi T, Suzuki T. Effect of tooth loss on spatial memory and trkB-mRNA levels in rats. *Hippocampus* 2008;18:542-7.
- Burke DM, Mackay DG. Memory, language, and ageing. *Philos Trans R Soc Lond B Biol Sci* 1997;352:1845-56.
- Perry RJ, Hodges JR. Attention and executive deficits in Alzheimer's disease. A critical review. *Brain* 1999;122:383-404.
- Salat DH, Buckner RL, Snyder AZ, Greve DN, Desikan RS, Busa E, Morris JC, Dale AM, Fischl B. Thinning of the cere-

- bral cortex in aging. *Cereb Cortex* 2004;14:721-30.
23. Morokuma M. Influence of the functional improvement of complete dentures on brain activity. *Nihon Hotetsu Shika Gakkai Zasshi* 2008;52:194-9.
 24. Klimesch W. EEG alpha and theta oscillations reflect cognitive and memory performance: a review and analysis. *Brain Res Brain Res Rev* 1999;29:169-95.
 25. Lund JP. Mastication and its control by the brain stem. *Crit Rev Oral Biol Med* 1991;2:33-64.
 26. Masumoto Y, Morinushi T, Kawasaki H, Takigawa M. Spectral analysis of changes in electroencephalographic activity after the chewing of gum. *Psychiatry Clin Neurosci* 1998;52:587-92.
 27. Masumoto Y, Morinushi T, Kawasaki H, Ogura T, Takigawa M. Effects of three principal constituents in chewing gum on electroencephalographic activity. *Psychiatry Clin Neurosci* 1999;53:17-23.