



The Use of a Pipeline Embolization Device for Treatment of a Ruptured Dissecting Middle Cerebral Artery M3/M4 Aneurysm: Challenges and Technical Considerations

Robert P. Berwanger, BS¹, Madeline C. Hoover, BS¹, John A. Scott, MD², Andrew J. DeNardo, MD², Krishna Amuluru, MD², Troy D. Payner, MD², Charles G. Kulwin, MD², Daniel H. Sahlein, MD²

¹Department of Neurosurgery, Indiana University School of Medicine, Indianapolis, IN, USA

²Division of Interventional Neuroradiology, Goodman Campbell Brain and Spine at Ascension St. Vincent Hospital, Carmel, IN, USA

Prompt, effective treatment is necessary following aneurysmal subarachnoid hemorrhage to prevent recurrent rupture, which is thought to double mortality. Atypical ruptured aneurysms, such as blister or dissecting pseudoaneurysms, or those that are unusually distal in the middle cerebral artery (MCA) are challenging to treat with either open or endovascular options, though the pipeline embolization device (PED) has shown promise in multiple case series. We present a case of a ruptured dissecting pseudoaneurysm in the distal MCA (distal M3/proximal M4) prefrontal division in a healthy young patient (<60 years) successfully treated with a PED. The PED was chosen both as the only vessel sparing option in the young patient as well as for its potential as a vessel sacrifice tool if the pseudoaneurysm was felt to be incompletely treated, which in this case was not necessary—though would have leveraged the thrombogenicity of the device as a therapeutic advantage.

Key Words: Pseudoaneurysm; Subarachnoid hemorrhage; Embolization, therapeutic; Middle cerebral artery

INTRODUCTION

Spontaneous subarachnoid hemorrhage (SAH) is a potentially devastating neurological event with an associated mortality rate of 25% within the first 24 hours.¹ The overwhelming majority of aneurysms are located on the circle of Willis or proximal anterior or posterior distributions and are saccular in etiology.² Less common types of aneurysms, such as those in the distal middle cerebral or anterior cerebral locations or those with an unusual morphology, are often more complex to treat.³ Currently,

there is no standard of treatment available for distal or dissecting aneurysms.

Increasingly, flow diversion devices, such as the pipeline embolization device (PED), have been utilized in the treatment of dissecting aneurysms.⁴ The PED was originally developed for large unruptured aneurysms in the anterior circulation,^{5,6} but has become a promising treatment option for ruptured and/or wide neck and fusiform aneurysms.⁷

We present a case in which a younger patient with a likely dissecting aneurysm in the distal left middle cerebral artery (MCA) was successfully treated with flow

Correspondence to:

Daniel H. Sahlein, MD

Department of Interventional Neuroradiology, Goodman Campbell Brain and Spine at Ascension St. Vincent Hospital, 13345 Illinois Street, Carmel, IN 46032, USA

Tel: +1-3173961160

Fax: +1-3174156175

E-mail: dsahlein@goodmancampbell.com

Received: February 1, 2022

Revised: February 23, 2022

Accepted: February 28, 2022

Copyright © 2022 Korean Society of Interventional Neuroradiology

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

pISSN 2093-9043

eISSN 2233-6273

diversion. This is the most distal use of the PED to our knowledge, and the excellent technical and clinical outcomes are instructive.

CASE REPORT

Presentation

A patient (<60 years old) with a past medical history of anxiety, depression, fibromuscular dysplasia, and breast cancer treated with lumpectomy and radiation in 2014–2015 was admitted to an outside hospital in early December 2019 with headache, neck pain, and vomiting. At this time, the

patient was alert and oriented with intact speech. A computed tomography (CT) scan (Fig. 1A) demonstrated SAH and parenchymal hemorrhage focused on the left superior and middle frontal gyrus in the prefrontal MCA distribution (Fig. 1). In addition, there was subdural hemorrhage. A catheter angiogram was performed and was consistent with dissecting pseudoaneurysm within the prefrontal division of the left MCA at the level of M3/M4, and this artery traversed the center of the collection of parenchymal and SAH (Fig. 1B). Five days after symptom onset, the patient was transferred to our hospital for endovascular intervention. On exam, the patient was alert and oriented and reported having a headache, but was negative for neurological deficits.




Fig. 1. (A) Noncontrast head axial CT demonstrates parenchymal (large black arrow), subarachnoid (small black arrow), and subdural (grey arrow) hemorrhage in the left hemisphere prefrontal mid convexity. There is surrounding edema (white arrow). (B) Catheter angiography of left internal carotid in the capillary phase shows the dissecting segment (white arrow) coursing through the center of a region of oligemia (black arrows) confirming that this ruptured M3/M4 pseudoaneurysm is the cause of hemorrhage. (C) 3D angiography demonstrates the ruptured M3/M4 dissecting aneurysm. (D) Magnified postprocedural angiography demonstrates appropriate positioning of the stent (white arrows) across the diseased and dissecting segment. (E) Six-month follow-up 3D angiography demonstrates patency of the stent (white arrows) and normal antegrade flow into the dominant (grey arrow) and jailed (black arrow) divisions of the prefrontal segment of the left middle cerebral artery. The stent (white arrows) has expanded to the nominal diameter of 2.5 mm during the 6-month interval follow-up. (F) Six-month follow-up MRI shows encephalomalacia/gliosis in the area of prior parenchymal hemorrhage (black arrows) with no additional signal abnormality either superior to the dominant collection or posterior to the collection (white arrow). CT, computed tomography; MRI, magnetic resonance imaging.

Catheter angiography showed a left MCA dissecting pseudoaneurysm in the M3 segment over the course of an 8 mm length of the artery with a maximum dimension of 2 mm and a minimum arterial diameter of slightly less than 1 mm (Fig. 1C). After a discussion of various treatment options, largely revolving around forms of arterial sacrifice, the decision was made to attempt to spare the artery using a PED. The plan was that if the implant looked ineffective in any way—for example, if the dissecting pseudoaneurysm continued to opacify briskly (or enlarge) or if the luminal diameter appeared too constrained following PED implant to continue to act as an effective conduit for antegrade flow, then the patient would be taken off their antiplatelet regimen and the highly thrombogenic PED would be used as an arterial sacrifice device.

Procedure

The patient was given 90 mg of ticagrelor at 4 AM and 90 mg of ticagrelor at 6 AM with 81 mg of aspirin for an 8 AM procedure start time as per our practice protocol for ruptured flow diversion cases. Maintenance thereafter was with 60 mg of ticagrelor twice daily and 81 mg of aspirin. The case was performed on a Siemens Artis Q biplane unit (Siemens, Munich, Germany). Femoral arterial access was utilized, and a 90 cm Neuron Max (Penumbra, Alameda, CA, USA) was utilized as a guide sheath. An 0.044-inch Phenom Plus (Medtronic, Irvine, CA, USA) was utilized as an intermediate catheter. All arteriography was performed through the Phenom Plus. Given the degree of vasospasm noted during routine catheterization, 5 mg of verapamil was given as premedication into the internal carotid artery, though this is not part of our practice routine.

Distal arterial access was achieved using a Phenom 27 microcatheter (Medtronic) advanced over a 0.014-inch Traxcess guidewire (MicroVention, Tustin, CA, USA). As per our practice routine, intravenous heparin (5,000 U intravenous bolus) was administered after catheter access across the segmental pseudoaneurysm.

Following a latency of 4 minutes, a 2.5×10 mm PED was deployed across the lesion with the distal aspect within the proximal M4 segment and proximally terminating within the M3 segment, spanning the dissecting segment with 3–4 mm landing zones on either zone. Left internal carotid arteriography demonstrated mild vasospasm within the prefrontal M3 division and stent patency with normal antegrade flow, and there was no longer visualization of the pseudoan-

eurysm (image not shown).

Another left internal carotid arteriographic run was performed following a latency of 310 minutes that showed stent patency and normal antegrade flow as well as decreased vasospasm (Fig. 1D). The decision was made to continue the patient's dual antiplatelets and bring the patient back later in the day to see if the stent remained widely patent and if the pseudoaneurysm remained stable.

Post Procedure

The patient tolerated the procedure well and post-operatively remained neurologically stable, following commands. The patient was maintained on nimodipine, levetiracetam, and a statin as per SAH protocol. Catheter angiography later the same day demonstrated stent patency and excellent antegrade flow with normal arterial transit (image not shown). There was no visualization of the pseudoaneurysm, and there was minimal vasospasm distal to the PED. The decision was made to continue the antiplatelet regimen to preserve the artery. A CT done 3 days postoperatively showed a stable left frontal opercular intraparenchymal hemorrhage and significant improvement of the SAH. Transcranial dopplers were negative for vasospasm. Magnetic resonance angiography 5 days post-procedure demonstrated a normal flow-related signal within the stent suggesting persistent normal antegrade flow. There were stable left frontal parenchymal and subdural hematomas (image not shown).

The patient experienced transient episodes of encephalopathy twice during the course of recovery without focality characterized by mild word-finding difficulty and intermittent headaches as well as mild paresthesia in the left foot, which was unchanged from the pre-procedure baseline. On the fifteenth postoperative day, the patient was sent home with outpatient physical and occupational therapy.

Follow Up

The patient came in for follow-up catheter cerebral angiography 6 months after the procedure (Fig. 1E). Left internal carotid arteriography demonstrated good patency of the PED, as well as antegrade flow of the M4 branch that had been jailed by the stent. Interestingly, in the interim, the pipeline stent which had previously been constrained within less than 1 millimeter diameter of the artery, had expanded to its nominal diameter of 2.5 mm. The entire lumen of the 2.5 mm device opacified with contrast resulting in a segment of the ectatic reconstructed artery that otherwise appeared

normal without residual luminal irregularity or pseudoaneurysm. Follow-up magnetic resonance imaging (MRI) demonstrated encephalomalacia limited to the original region of the parenchymal hemorrhage, not involving the territory of the dominant division of the prefrontal branch that supplied the territory posterior to the region of post-hemorrhage encephalomalacia (Fig. 1F).

DISCUSSION

This case is clinically significant for a multitude of reasons. First, this study provides further evidence for the efficacy of endovascular flow diversion with the PED in treating ruptured aneurysms. The use of a PED for the treatment of ruptured aneurysms with architecture that makes conventional saccular approaches (surgical clip reconstruction, endovascular coiling, Woven EndoBridge embolization [MicroVention, Aliso Viejo, CA, USA]) technically challenging or high risk is becoming increasingly prevalent because of the inherent advantages of the endoluminal approach in these cases and the empirically excellent clinical outcomes.^{7,8} This patient's aneurysm was unique in that it was likely a ruptured dissecting aneurysm as well as very distal. While there have been reports of PED use to treat dissecting aneurysms, to our knowledge there has not been a case of the PED for treatment of an aneurysm in such a distal location. The alternative and more conventional approach would have been arterial sacrifice, likely with coils; though, completely occluding pial arteries with coils can be challenging, and attempting curative coil occlusion in a dissecting aneurysm carries an inherent risk of perforation. Our approach was also designed to spare as much cortex as possible, and while the territory spared is not traditionally considered eloquent, there are benefits to minimizing cortical gliosis in a younger patient, such as lowering the likelihood of developing a chronic seizure disorder.⁹ Follow-up MRI in this case clearly demonstrated intact cortex and subcortical white matter dorsal to the area of parenchymal hemorrhage in the territory supplied by the dominant branch of the prefrontal division of the MCA.

While there is an inherent risk in using flow diversion in very-small caliber vessels, a recent case series of 67 patients demonstrated the use of flow diverters in small arteries in a caliber of less than 2 mm, resulting in only 1 case of flow delay associated with flow diameter reduction.¹⁰ In this case, The artery remained patent and expanded with the device

as evidenced by the ectasia within the stented segment when compared to the native artery immediately proximal and distal to the PED implant, which suggests that the relatively thin-walled distal pial arterial circulation is amenable to flow diversion and is tolerant of oversized devices over time.

Another application from this case is the use of the PED's thrombogenicity as a potential therapeutic strategy. As mentioned, if the segment containing the PED implant was not effective in treatment of the pseudoaneurysm, the strategy was to use the PED as a closure device, taking advantage of the thrombogenicity of a braided metal implant. While this strategy was unnecessary in this patient's case, it could provide an alternative course of action when treating distal arterial aneurysms.

This case showed the effectiveness of flow diversion in treating ruptured dissecting aneurysms as well as for treating aneurysms complicated by distal location. The device's thrombogenicity should be considered as a therapeutic strategy in the appropriate clinical circumstances.

Fund

None.

Ethics Statement

Institutional Review Board approval for case reports is not required at our institution; however, written informed consent was obtained for the procedure. Informed consent for publication was not obtained so we anonymized the patient.

Conflicts of Interest

DS serves as a speaker and receives grant support from MicroVention, a proctor and speaker with Medtronic and a consultant for Phenox. KA is an educational consultant for Medtronic. All other authors have no conflicts of interest.

Author Contributions

Concept and design: RB, MH, JS, AD, KA, TP, CK, and DS. Analysis and interpretation: RB, JS, AD, KA, CK, and DS. Data collection: RB, MH, and DS. Writing the article: RB, MH, JS, AD, KA, TP, CK, and DS. Critical revision of the article: RB, MH, JS, AD, KA, TP, CK, and DS. Final approval of the article: RB, MH, JS, AD, KA, TP, CK, and DS. Overall responsibility: DS.

ORCID

Robert P. Berwanger: <https://orcid.org/0000-0001-6797-5281>
Madeline C. Hoover: <https://orcid.org/0000-0001-5660-985X>

John A. Scott: <https://orcid.org/0000-0001-8739-3780>

Andrew J. DeNardo: <https://orcid.org/0000-0001-7585-8624>

Krishna Amuluru: <https://orcid.org/0000-0002-8859-8574>

Troy D. Payner: <https://orcid.org/0000-0002-8799-679X>

Charles G. Kulwin: <https://orcid.org/0000-0001-9990-2028>

Daniel H. Sahlein: <https://orcid.org/0000-0002-6591-9427>

REFERENCES

1. Long B, Koyfman A, Runyon MS. Subarachnoid hemorrhage: updates in diagnosis and management. *Emerg Med Clin North Am* 2017;35:803-824
2. Keedy A. An overview of intracranial aneurysms. *Mcgill J Med* 2006;9:141-146
3. Joo SP, Kim TS, Choi JW, Lee JK, Kim YS, Moon KS, et al. Characteristics and management of ruptured distal middle cerebral artery aneurysms. *Acta Neurochir (Wien)* 2007;149:661-667
4. Cunegatto-Braga M, Hogan B, Aguilar-Salinas P, Beier AD, Hanel RA. Pipeline embolization device flow diversion for a dissecting ruptured posterior cerebral artery aneurysm in a pediatric patient. *World Neurosurg* 2018;117:255-260
5. Becske T, Kallmes DF, Saatci I, McDougall CG, Szikora I, Lanzino G, et al. Pipeline for uncoilable or failed aneurysms: results from a multicenter clinical trial. *Radiology* 2013;267:858-868
6. Hanel RA, Kallmes DF, Lopes DK, Nelson PK, Siddiqui A, Jabbour P, et al. Prospective study on embolization of intracranial aneurysms with the pipeline device: the PREMIER study 1 year results. *J Neurointerv Surg* 2020;12:62-66
7. Chalouhi N, Zanaty M, Whiting A, Tjoumakaris S, Hasan D, Aji-boye N, et al. Treatment of ruptured intracranial aneurysms with the pipeline embolization device. *Neurosurgery* 2015;76:165-172; discussion 172
8. Lozupone E, Piano M, Valvassori L, Quilici L, Pero G, Visconti E, et al. Flow diverter devices in ruptured intracranial aneurysms: a single-center experience. *J Neurosurg* 2018;128:1037-1043
9. Johnson AM, Sugo E, Barreto D, Hiew CC, Lawson JA, Connolly AM, et al. The severity of gliosis in hippocampal sclerosis correlates with pre-operative seizure burden and outcome after temporal lobectomy. *Mol Neurobiol* 2016;53:5446-5456
10. Bender MT, Zarrin DA, Campos JK, Lin LM, Huang J, Caplan JM, et al. Tiny pipes: 67 cases of flow diversion for aneurysms in distal vessels measuring less than 2.0 mm. *World Neurosurg* 2019;127:e193-e201