



Diels-Alder Type Adducts from Hairy Root Cultures of *Morus macrourea*

Nizar Happyana^{1,*}, Euis H. Hakim¹, Yana M. Syah¹, Oliver Kayser², Lia D. Juliawaty¹, Didin Mujahidin¹,
Tri M. Ermayanti³, and Sjamsul A. Achmad¹

¹Chemistry Study Program, Faculty of Mathematics and Natural Sciences, Bandung Institute of Technology, Jl. Ganesha 10, 40132 Bandung, Indonesia

²Department of Technical Biochemistry, Technical University of Dortmund, Emil-Figge-Str. 66, 44227 Dortmund, Germany

³Research Center of Biotechnology, Indonesian Institute of Sciences (LIPI), Jl. Raya Bogor KM. 46 Cibinong 16911, Indonesia

Abstract – Three Diels-Alder type adducts, guangsangon E (**1**), chalcomoracin (**2**) and sorocein I (**3**) were isolated from hairy root cultures of *Morus macrourea*. The structures of the isolated compounds (**1** - **3**) were determined by spectroscopic method (NMR and MS), and spectral comparison to literature. Cytotoxic activities of the isolated compounds (**1** - **3**) were investigated against P-388 murine leukemia cell line. Guangsangon E (**1**) showed the most potent cytotoxicity against P-388 murine leukemia cell line with IC₅₀ value of 2.75 ± 0.32 µg/mL. To the best of our knowledge, guangsangon E (**1**) and sorocein I (**3**) were reported for the first time from the tissue cultures of *M. macrourea*.

Keywords – Diels-Alder type adduct, *Morus macrourea*, tissue cultures, cytotoxicity

Introduction

Morus macrourea, locally known as ‘Andalas’, is a rare plant found and native to the Malesian region.¹ It is a tree growing up to 35 m and found at altitudes of 900 - 1600 m.¹ In the region, the plant has been valued for the high quality of the wood due to its resistance to insects. Unfortunately, destruction of Indonesian forest threatens the existence and future study of this plant, especially the investigation of secondary metabolites. Plant tissue culture is a method that can be applied to solve the obstacle in investigating compounds in this plant.

Previous phytochemical studies on *M. macrourea* revealed the isolation of Diels-Alder type adducts.²⁻⁷ Two new benzofuran derivatives, macourins A and B, were also isolated from this plant.⁸ Our previous works on the chemical constituents of this plant have resulted in the isolation of other phenolic compounds, including andalasin A, andalasin B, quercimeritrin, moracin B, moracin P, oxyresveratrol, and mulberroside C.^{9,10} Furthermore, our work on hairy root cultures of *M. macrourea* afforded chalcomoracin and mulbereofuran P, Diels-Alder type adducts.¹¹

Hairy root culture is one of the interesting methods in the plant tissue cultures since the culture able to grow fast and produce secondary metabolites as well. In our previous work, hairy root cultures of *M. macrourea* were successfully produced via transformation of the plant leaves using *Agrobacterium rhizogenes*.¹¹ Furthermore, the effect of *Saccharomyces cerevisiae* extract on production of secondary metabolites of *M. macrourea* hairy root cultures had been also investigated.¹¹ In continuation of our studies on these elicited hairy root cultures, here we report the isolation and identification of Diels-Alder type adducts from the methanol extract of the cultures. The isolated compounds (**1** - **3**) were tested for cytotoxicity against P-388 murine leukemia cell line.

Experimental

General experimental procedures – IR spectra were recorded on a Perkin Elmer spectrum One FT-IR spectrometer. UV spectra were recorded on a Varian Corry 100 Conc spectrometer. NMR spectra were recorded on JEOL JNM A 5000 at 400 MHz (¹H) and 100 MHz (¹³C). MS spectrum of compound **1** was recorded on an ESI-TOF Waters Premier LCT XE mass spectrometer. Meanwhile, MS spectra of compound **2** and compound **3** were recorded using an API 3000 mass spectrometer (Applied Biosystems/MDS Sciex). Vacuum liquid chromatography

*Author for correspondence

Nizar Happyana, Chemistry Study Program, Faculty of Mathematics and Natural Sciences, Bandung Institute of Technology, Jl. Ganesha 10, 40132 Bandung, Indonesia
Tel: +62-22-250-2103; E-mail: nizar@chem.itb.ac.id




Fig. 1. Growth of hairy root cultures of *M. macroura*.

graphy (VLC) and radial chromatography were carried out using Merck Silica gel 60 GF₂₅₄. Column chromatography was performed using Sephadex LH-20. Flash column chromatography was carried out by Merck Silica gel 60 (70-230 mesh). TLC was performed using Merck precoated silica gel F₂₅₄ plates.

***M. macroura* hairy root cultures** – Hairy root transformation of *M. macroura* by *A. rhizogenes* had been assisted by Research Center of Biotechnology, Indonesian Institute of Sciences (LIPI). Hairy root cultures of *M. macroura* were grown in 50 mL of Murashige Skoog (MS) media with 0.5 ppm of indol-3-butyric acid (IBA) at room temperature and shaken at 100 rpm. Subculture (propagation) of the hairy root cultures was carried out at week 4 until the number of the cultures are enough for the elicitation. Fig. 1 showed the growth of hairy root cultures of *M. macroura* from week 0 till week 4. Elicitation was conducted by adding 500 μ L of *S. cerevisiae* (2.5% w/v) into the hairy root cultures at the week 8 and followed by shaking at 100 rpm for 7 days and then harvested.

***S. cerevisiae* preparation** – *S. cerevisiae* was obtained from Microbiology Laboratory, School of Life Sciences and Technology, Bandung Institute of Technology, Indonesia. *S. cerevisiae* was grown in 50 mL of glucose-yeast extract media (GYE) at room temperature and shaken at 100 rpm. The cultures of *S. cerevisiae* were subcultivated every 2 days by transferring 10 mL of the culture into 40 mL of the fresh liquid medium. The cultures were harvested at 16 hours old. The supernatant was removed, and the cells were dried and grinded into powders.

Extraction and isolation – Dried powdered of hairy

roots of *M. macroura* (35.5 g) were macerated with MeOH yielding 7.9 g of dried extract. The MeOH extract was fractionated with vacuum liquid chromatography (hexane-EtOAc, gradient concentrations) into seven fractions (A-H). Fraction C was subjected to silica gel radial chromatography (EtOAc-chloroform, 2:8) yielding 7 fractions. Fraction C7 was successively purified by Sephadex LH-20 column chromatography and flash silica gel column chromatography (hexane-acetone, 6:4), afforded compound **1** (8 mg) and compound **2** (20 mg). Fraction E was successively purified by silica gel radial chromatography (chloroform-methanol, 9:1) and flash silica gel column chromatography (hexane-acetone, 6:4) yielding compound **3** (5 mg).

Guangsangon E (1): Brown amorphous gum; UV (MeOH) λ_{\max} (log ϵ) = 203 (8.45), 225 (7.34), 282 (5.61), 323 (4.93) nm; IR ν_{\max} (KBr) = 3445, 2927, 1619, 1497, 1461 cm^{-1} ; ^1H NMR (500 MHz, acetone- d_6): δ (ppm) = 1.56 (3H, s, H-24"), 1.67 (3H, s, H-25"), 1.89 (3H, s, H-7"), 2.07 (1H, m, H-6a"), 2.46 (1H, m, H-6b"), 3.19 (2H, d, J = 7.0 Hz, H-21"), 3.67 (1H, m, H-5"), 4.38 (1H, br s, H-3"), 4.68 (1H, dd, J = 6.5, 7.0 Hz, H-4"), 5.12 (1H, t, J = 7.0 Hz, H-22"), 5.55 (1H, s, H-2"), 6.14 (1H, dd, J = 1.5, 8.5 Hz, H-19"), 6.26 (1H, d, J = 1.5 Hz, H-17"), 6.33 (1H, d, J = 2.0 Hz, H-4'), 6.42 (1H, d, J = 8.5 Hz, H-13"), 6.81 (1H, s, H-7), 6.83 (1H, d, J = 2.0 Hz, H-2'), 6.83 (1H, d, J = 2.0 Hz, H-6'), 6.88 (1H, d, J = 8.5 Hz, H-20"), 7.04 (1H, s, H-3), 7.45 (1H, s, H-4), 8.03 (1H, d, J = 8.5 Hz, H-14"), 13.03 (1H, s, HO-10"); ^{13}C NMR (125 MHz, acetone- d_6): δ (ppm) = 17.9 (CH₃, C-25"), 22.3 (CH₂, C-21"), 23.8 (CH₃, C-7"), 25.9 (CH₃, C-24"), 33.5 (CH, C-5"), 37.4 (CH₂, C-6"), 37.9 (CH, C-3"), 48.5 (CH, C-4"), 97.2 (CH, C-7), 102.6 (CH, C-3), 103.4 (CH, C-17"), 103.8 (CH, C-4'), 103.8 (CH, C-2'), 103.9 (CH, C-6'), 107.3 (CH, C-13"), 107.6 (CH, C-19"), 115.4 (C-11"), 115.8 (C-9"), 121.9 (C-3 α), 123.2 (C-15"), 123.6 (CH, C-4), 123.8 (CH, C-22"), 124.5 (CH, C-2"), 125.6 (C-5), 127.1 (C-8"), 129.8 (CH, C-20"), 129.8 (C-1'), 130.6 (CH, C-14"), 131.2 (C-23"), 135.2 (C-1"), 155.1 (C-7 α), 155.1 (C-16"), 155.3 (C-2), 156.4 (C-6), 157.3 (C-18"), 159.8 (C-3'), 159.8 (C-5'), 161.9 (C-12"), 163.9 (C-10"). HRESI-MS: m/z 649.2426 (calcd. 649.2438 for C₃₉H₃₇O₉). The data were accordance with the literature.³

Chalcomoracin (2): Yellow amorphous gum; UV (MeOH) λ_{\max} (log ϵ) = 203 (7.45), 319 (7.49), 333 (3.84) nm; IR ν_{\max} (KBr) = 3349, 2913, 1669, 1619, 1489, 1440 cm^{-1} ; ^1H NMR (400 MHz, acetone- d_6): δ (ppm) = 1.56 (3H, s, H-25"), 1.70 (3H, s, H-24"), 1.95 (3H, s, H-7"), 2.21 (1H, m, H-6a"), 2.52 (1H, m, H-6b"), 3.25 (2H, d, J = 6.9 Hz, H-21"), 3.75 (1H, t, J = 4.5 Hz, H-5"), 4.11 (1H, br s, H-

3"), 4.63 (1H, t, $J = 4.5$ Hz, H-4"), 5.14 (1H, t, $J = 6.9$ Hz, H-22"), 5.77 (1H, br s, H-2"), 6.30 (1H, dd, $J = 8.0, 2.1$ Hz, H-19"), 6.43 (1H, d, $J = 9.3$ Hz, H-13"), 6.50 (1H, d, $J = 2.1$ Hz, H-17"), 6.76 (1H, br s, H-2'), 6.76 (1H, br s, H-6'), 6.76 (1H, m, H-5), 6.92 (1H, br s, H-3), 6.92 (1H, br s, H-7), 6.98 (1H, d, $J = 8.0$ Hz, H-20"), 7.34 (1H, d, $J = 8.1$ Hz, H-4), 8.44 (1H, d, $J = 9.3$ Hz, H-14"); ^{13}C NMR (100 MHz, acetone- d_6): δ (ppm) = 17.8 (CH₃, C-25"), 22.2 (CH₂, C-21"), 23.8 (CH₃, C-7"), 25.8 (CH₃, C-24"), 32.2 (CH₂, C-6"), 33.1 (CH, C-3"), 36.5 (CH, C-5"), 47.8 (CH, C-4"), 98.3 (CH, C-7), 101.8 (CH, C-3), 103.4 (CH, C-17"), 104.8 (CH, C-2'), 104.8 (CH, C-6'), 107.4 (CH, C-19"), 108.1 (CH, C-13"), 113.0 (CH, C-5), 113.4 (C-9"), 115.8 (C-4'), 116.6 (C-11"), 121.8 (CH, C-4), 121.8 (C-15"), 122.6 (C-3a), 123.1 (CH, C-22"), 124.4 (CH, C-2"), 128.7 (CH, C-20"), 130.9 (C-1'), 131.5 (C-23"), 132.1 (CH, C-14"), 133.8 (C-1"), 155.4 (C-7a), 156.3 (C-2), 156.5 (C-3'), 156.5 (C-5'), 156.6 (C-16"), 157.8 (C-6), 157.8 (C-18"), 163.3 (C-12"), 164.4 (C-10"), 209.7 (C-8"); ESI-MS: m/z 648.5 (C₃₉H₃₆O₉). The data were in agreement with the literature.^{12,13}

Sorocin I (3): Purple amorphous gum; UV (MeOH) λ_{max} (log ϵ) = 205 (4.45), 276 (4.25), 332 (4.05) nm; IR ν_{max} (KBr) = 3439, 2920, 1614, 1508, 1452 cm⁻¹; ^1H NMR (400 MHz, acetone- d_6): δ (ppm) = 1.58 (3H, s, H-25"), 1.71 (3H, s, H-24"), 1.77 (3H, s, H-7"), 2.00 (1H, overlapped, H-6a"), 2.72 (1H, dd, $J = 17.0, 4.8$ Hz, H-6b"), 2.96-3.00 (1H, m, H-4"), 2.96-3.00 (1H, m, H-5"), 3.32 (1H, d, $J = 6.9$ Hz, H-21"), 3.63 (1H, br t, $J = 5.4$ Hz, H-3"), 5.20 (1H, t, $J = 6.9$ Hz, H-22"), 6.34 (1H, d, $J = 8.4$ Hz, H-13"), 6.35 (1H, dd, $J = 8.4, 2.6$ Hz, H-5), 6.36 (1H, d, $J = 2.6$ Hz, H-17"), 6.41 (1H, d, $J = 2.6$ Hz, H-3), 6.44 (1H, br d, $J = 5.4$ Hz, H-2"), 6.51 (1H, dd, $J = 8.4, 2.6$ Hz, H-19"), 6.63 (1H, br s, H-2'), 6.65 (1H, br s, H-6'), 6.86 (1H, d, $J = 16.5$ Hz, H- β), 7.12 (1H, d, $J = 8.4$ Hz, H-14"), 7.15 (1H, d, $J = 8.4$ Hz, H-20"), 7.30 (1H, d, $J = 16.5$ Hz, H- α), 7.38 (1H, $J = d, 8.4$ Hz, H-6); ^{13}C NMR (100 MHz, acetone- d_6): δ (ppm) = 17.9 (CH₃, C-24"), 23.0 (CH₂, C-21"), 23.8 (CH₃, C-7"), 25.9 (CH₃, C-25"), 28.6 (CH, C-5"), 35.4 (CH, C-3"), 35.9 (CH, C-6"), 37.2 (CH, C-4"), 103.5 (CH, C-3), 103.6 (C, C-8"), 103.9 (CH, C-17"), 106.9 (CH, C-6'), 107.1 (CH, C-13"), 107.5 (CH, C-2'), 108.4 (CH, C-5), 110.2 (CH, C-19"), 111.6 (C, C-4'), 116.9 (C, C-11"), 117.1 (C, C-1), 117.1 (C, C-9"), 117.2 (C, C-15"), 112.8 (CH, C-2"), 123.8 (CH, C-22"), 124.6 (CH, C- α), 125.3 (CH, C- β), 126.1 (CH, C-14"), 127.9 (CH, C-20"), 128.3 (CH, C-6), 131.2 (C, C-23"), 133.5 (C, C-1"), 139.4 (C, C-1'), 152.6 (C, C-16"), 153.3 (C, C-3'), 155.5 (C, C-5'), 156.9 (C, C-10"), 157.5 (C, C-12"), 157.6 (C, C-4), 157.8 (C, C-18"), 159.1 (C, C-2); ESI-

MS: m/z 632.5 (C₃₉H₃₆O₈). The data were in agreement with the literature.¹⁴

Cytotoxicity assay – Cytotoxicity assay against P388 murine leukemia cell was carried out based on published reports.^{15,16} The leukemia cells at a density of 3×10^4 cell/mL, were plated in 96-well cultures dishes and incubated at 37 °C in humidified CO₂ incubator (5%) for 24 h. Guangsangon E (**1**), chalcomoracin (**2**) and sorocin I (**3**) were first dissolved in DMSO at the required concentrations and prepared subsequently in six desirable concentrations (100 – 0.1 $\mu\text{g/mL}$) using PBS (phosphoric buffer solution, pH = 7.30-7.65). The prepared samples were added into the sample wells. Meanwhile, the control wells received only DMSO and PBS. Afterward the cell cultures were incubated again for 48 h in 5% CO₂ incubator at 37 °C. After the incubation, MTT solution [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide] was added and incubated for 4 h. In the next step, MTT-stop solution containing sodium dodecyl sulphate (SDS) were added, and then the last incubation for 24 h was conducted. Cell survival was evaluated with a multi-well scanning spectrophotometer (ELX 800 UV, universal Microplate Reader, Bio-Tek Instruments, Inc.) at 540 nm. All data are presented as the mean values of triplicates.

Result and Discussion

Compound **1** was isolated from fraction C of the hairy root extract of *M. macroura* as a brown amorphous gum. According to the spectroscopic analysis (NMR and MS) and comparing with the literature,³ compound **1** was assigned to guangsangon E. The molecular structure of guangsangon E (**1**) is depicted in Fig. 2. Guangsangon E (**1**) is a Diels-Alder type adduct derived from a substituted chalcone and a substituted 2-phenylbenzofuran. In this work, guangsangon E (**1**) was tested against P-388 murine leukemia cell line. Compared to the other isolated compounds (**2** and **3**), guangsangon E (**1**) exhibited the most potent cytotoxicity against P-388 murine leukemia cell line with IC₅₀ value of 2.75 ± 0.32 $\mu\text{g/mL}$. To the best of our knowledge, this is the first report concerning cytotoxic activity of guangsangon E (**1**). Compound **1** was firstly isolated from stem bark of *M. macroura* and possessed anti-oxidant and anti-inflammation activities.³ Recently, guangsangon E (**1**) has been reported actively inhibiting tyrosinase.⁶

Compound **2** was isolated and purified from fraction C as a yellow amorphous gum. NMR and MS spectra analysis and data comparison to references^{12,13} concluded that compound **2** is chalcomoracin, a Diels-Alder type




Fig. 2. The molecular structures of **1**–**3**, isolated from hairy root cultures of *M. macroura*.

adduct derived from a dehydroprenylchalcone and a substituted 2-arylbenzofuran. The structure of chalcomoracin (**2**) is depicted in Fig. 2. Chalcomoracin (**2**) is a phytoalexin, collected for the first time from diseased mulberry leaves (*M. alba*).¹² Chalcomoracin (**2**) has been reported having antibiotic activity via inhibition of enoyl-acyl carrier reductase.¹⁷

According to the cytotoxic assay, chalcomoracin (**2**) showed moderate cytotoxicity against P-388 murine leukemia cell line with IC_{50} value of 5.46 ± 0.55 μ g/mL. Chalcomoracin (**2**) was reported possessing moderate cytotoxic activities against five human cancer cell lines, including A549, Bel 7402, BGC 823, HCT-8, and A2780.¹⁸ Recently, molecular mechanism of chalcomoracin (**2**) as anticancer agent had been reported.¹⁹ Chalcomoracin (**2**) inhibited the cancer cell growth by promoting paraptosis and triggering oxidative stress via a mitophagy.¹⁹ Therefore, the result of cytotoxic assay of chalcomoracin (**2**) against P-388 murine leukemia cell line was in accordance with

the previous reports^{18,19} and confirmed chalcomoracin (**2**) as potent anticancer agent.

Compound **3** was isolated as a purple amorphous gum. The structure of compound **3** was determined based on NMR and MS data analysis and spectral comparison to reported work.¹⁴ Compound **3** was assigned as sorocein I, a ketalized Diels-Alder type adduct formed from a dehydroprenylchalcone and a dehydroprenylstilbene. Sorocein I (**1**) was isolated for the first time from *Sorocea ilicifolia*.¹⁴ Molecular structure of sorocein I (**3**) was depicted in Fig. 2. Cytotoxic assay revealed that sorocein I (**3**) was not active against P-388 murine leukemia cell line.

In this report, three Diels-Alder type adducts, guangsangon E (**1**), chalcomoracin (**2**) and sorocein I (**3**) were isolated from hairy root cultures of *M. macroura*. Guangsangon E (**1**) and sorocein I (**3**) were recorded for the first time in this tissue culture. Furthermore, cytotoxic activities of guangsangon E (**1**) and sorocein I (**3**) were reported for the first time as well.

Acknowledgments

This study was partially funded by Directorate General of Higher Education under research contract No. 127/SP2H/PTNBH/DRPM/2018. We are grateful to Dr. Jalifah Latif, UKM, Malaysia, for the NMR measurements.

References

- (1) Heyne, K. Research and Development Agency; Ministry of Forestry: Indonesia, **1987**, p 659.
- (2) Dai, S. J.; Ma, Z. B.; Wu, Y.; Chen, R. Y.; Yu, D. Q. *Phytochemistry* **2004**, *65*, 3135-3141.
- (3) Dai, S. J.; Mi, Z. M.; Ma, Z. B.; Li, S.; Chen, R. Y.; Yu, D. Q. *Planta Med.* **2004**, *70*, 758-763.
- (4) Dai, S. J.; Wu, Y.; Wang, Y. H.; He, W. Y.; Chen, R. Y.; Yu, D. Q. *Chem. Pharm. Bull.* **2004**, *52*, 1190-1193.
- (5) Dai, S. J.; Yu, D. Q. *Nat. Prod. Res.* **2006**, *20*, 676-679.
- (6) Wang, Y. F.; Xu, L.; Gao, W.; Niu, L.; Huang, C.; Yang, P.; Hu, X. *Planta Med.* **2018**, *84*, 336-343.
- (7) Wang, Y. F.; Yu, M. H.; Xu, L.; Niu, L.; Huang, C.; Xu, H.; Yang, P.; Hu, X. *Phytochem. Lett.* **2018**, *26*, 149-153.
- (8) Sun, S. G.; Chen, R. Y.; Yu, D. Q. *J. Asian Nat. Prod. Res.* **2001**, *3*, 253-259.
- (9) Syah, Y. M.; Achmad, S. A.; Ghisalberti, E. L.; Hakim, E. H.; Iman, M. Z.; Makmur, L.; Mujahiddin, D. *Fitoterapia* **2000**, *71*, 630-635.
- (10) Syah, Y. M.; Achmad, S. A.; Ghisalberti, E. L.; Hakim, E. H.; Makmur, L.; Soekamto, N. H. *J. Chem. Res.* **2004**, *2004*, 339-340.
- (11) Happyana, N.; Mujahiddin, D.; Juliawaty, L. D.; Makmur, L.; Achmad, S. A.; Syah, Y. M.; Hakim, E. H. Proceeding of The International Seminar on Chemistry; Universitas Padjadjaran: Indonesia, **2008**, pp 257-261.
- (12) Kang, J.; Chen, R. Y.; Yu, D. Q. *Planta Med.* **2006**, *72*, 52-59.
- (13) Takasugi, M.; Nagao, S.; Masamune, T.; Shirata, A.; Takahashi, K. *Chem. Lett.* **1980**, *9*, 1573-1576.
- (14) Ferrari, F.; Delle Monache, F. *Fitoterapia* **2001**, *72*, 301-303.
- (15) Sahidin.; Hakim, E. H.; Juliawaty, L. D.; Syah, Y. M.; bin Din, L.; Ghisalberti, E. L.; Latip, J.; Said, I. M.; Achmad, S. A. *Z. Naturforsch. C.* **2005**, *60*, 723-727.
- (16) Alley, M. C.; Scudiero, D. A.; Monks, A.; Hursey, M. L.; Czerwinski, M. J.; Fine, D. L.; Abbott, B. J.; Mayo, J. G.; Shoemaker, R. H.; Boyd, M. R. *Cancer Res.* **1988**, *48*, 589-601.
- (17) Kim, Y. J.; Sohn, M. J.; Kim, W. G. *Biol. Pharm. Bull.* **2012**, *35*, 791-795.
- (18) Zhang, Q. J.; Tang, Y. B.; Chen, R. Y.; Yu, D. Q. *Chem. Biodivers.* **2007**, *4*, 1533-1540.
- (19) Han, H.; Chou, C. C.; Li, R.; Liu, J.; Zhang, L.; Zhu, W.; Hu, J.; Yang, B.; Tian, J. *Sci. Rep.* **2018**, *8*, 9566.

Received February 23, 2019

Revised April 11, 2019

Accepted April 12, 2019