

Abdominal Bloating: Pathophysiology and Treatment

A Young Seo,¹ Nayoung Kim^{1,2*} and Dong Hyun Oh¹

¹Department of Internal Medicine, Seoul National University, Bundang Hospital, Seongnam, Gyeonggi-do, Korea; and ²Department of Internal Medicine and Liver Research Institute, Seoul National University College of Medicine, Seoul, Korea

Abdominal bloating is a very common and troublesome symptom of all ages, but it has not been fully understood to date. Bloating is usually associated with functional gastrointestinal disorders or organic diseases, but it may also appear alone. The pathophysiology of bloating remains ambiguous, although some evidences support the potential mechanisms, including gut hypersensitivity, impaired gas handling, altered gut microbiota, and abnormal abdominal-phrenic reflexes. Owing to the insufficient understanding of these mechanisms, the available therapeutic options are limited. However, medical treatment with some prokinetics, rifaximin, lubiprostone and linaclotide could be considered in the treatment of bloating. In addition, dietary intervention is important in relieving symptom in patients with bloating.

(J Neurogastroenterol Motil 2013;19:433-453)

Key Words

Bloating; Pathophysiology; Rifaximin; Therapy

Introduction

Bloating is one of the most common gastrointestinal (GI) symptoms, which is a frequent complaint in the patients of all ages. This symptom is very common in patients with irritable bowel syndrome (IBS) and other functional gastrointestinal disorders (FGIDs) as well as in patients with organic disorders. Many clinicians encounter the patients' complaints such as "too much gas in abdomen," "heavy and uncomfortable feeling in abdomen" and "full belly." The severity of bloating is varied from

mild discomfort to severe, and it is one of the bothersome symptoms of the patients, affecting their quality of life. Despite being one of the frequent and bothersome complaints, bloating remains incompletely understood of all the symptoms. Therefore clinicians need to be more considerate when evaluating patients with abdominal bloating.

The possible causes of bloating are various and complicated, thus intestinal gas production and transit, gut microflora and hypersensitivity of the patient's gut might be the factors for the symptom generation. As the underlying mechanism of bloating remains elusive to date, there are few evidences for diagnostic and

Received: September 3, 2013 Revised: September 10, 2013 Accepted: September 16, 2013

© This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Correspondence: Nayoung Kim, MD

Department of Internal Medicine, Seoul National University Bundang Hospital, 82 Gumi-ro, 173 Beon-gil, Bundang-gu, Seongnam, Gyeonggi-do 463-707, Korea

Tel: +82-31-787-7008, Fax: +82-31-787-4051, E-mail: nayoungkim49@empas.com

Financial support: This work was supported by the National Research Foundation of Korea (NRF) grant (No. 2011-0030001) for the Global Core Research Center (GCRC) funded by the Korea government (MSIP).

Conflicts of interest: None.

Author contributions: A Young Seo analyzed the references and wrote a manuscript. Dong Hyun Oh collected references and adapted the figures. Nayoung Kim provided ideas, designed and supervised the manuscript.

therapeutic options available. In addition, patients who complain of bloating tend to have IBS or functional dyspepsia (FD), thus most of the therapeutic approaches of abdominal bloating are based on the treatments of IBS and FD.¹⁻⁴ However bloating could occur alone without associated diseases and there has been not enough data of randomized controlled trials for treatment of bloating alone. The treatments for bloating have not been standardized and there is no evidence-based algorithm. Although there have been several comprehensive reviews for pathophysiology or treatment of abdominal bloating in FGID (Table 1),⁵⁻¹⁰ both clinical trials and systematic reviews regarding functional bloating (FB) alone are scarce so far. Only one available pilot study was from Spain, which indicated that sugar intolerance was frequently observed in patients with FB and associated with bloating symptom. Additionally, a malabsorbed sugar-free diet gave rise to clinical improvement in high percentage of patients.¹¹ Thus, in the clinical setting, rational treatment for bloating is hard and the result of treatment is frequently unsatisfactory. Most treatment options for bloating are similar to treatments for IBS, but as previously mentioned, FB and IBS are in different disease entities.

Aims and Methods

Our aim was to explain the clinical importance, pathophysiologic mechanisms, and management of abdominal bloating, thus to provide a better understanding of this specific problem. We reviewed the literature of mechanisms and treatment interventions for abdominal bloating based on a PubMed search on the following terms; “abdominal bloating,” “intestinal gas and IBS,” “distension and IBS” and “FGID.” We also quoted important knowledge from a standard textbook, the chapter “intestinal gas.”¹²

Definition

Bloating is defined as subjective discomfort by patient’s sensation of intestinal gas; otherwise, abdominal distension is a visible increase in abdominal girth. In the past, bloating had been considered to be related to abdominal distension directly, but recent studies have suggested that it is not always accompanied by abdominal distension.¹³ There have been many studies to evaluate the relationship between bloating and abdominal distension. One study has shown that actual abdominal distension only occurred in about half of the patients suffering from bloating.¹⁴ In addition, some patients with both visceral hypersensitivity and FGID

complained of bloating in the absence of visible distension.^{15,16} Briefly, abdominal bloating is the subjective symptom and distension is the objective sign, so bloating and distension should be considered as separate disorders with different mechanisms. Although bloating has been considered as a supportive symptom for IBS or FD according to Rome classification, FB is also included as an independent entity in Rome criteria.¹⁷⁻¹⁹ The diagnosis of FB is made in patients who do not meet the diagnostic criteria of IBS or other FGIDs, but have recurrent symptoms of bloating. According to Rome III, the diagnostic criteria include recurrent feeling of bloating or visible distension at least 3 days a month in the last 3 months with symptom onset at least 6 months prior to diagnosis. Also it should exclude FD, IBS or other FGIDs. The name has been changed from functional abdominal bloating in Rome I and II criteria to functional bloating in Rome III criteria.¹⁷⁻¹⁹

Epidemiology

‘Bloating’ has been first described by Alvarez of the Mayo Clinic in 1949, in a woman patient with psychological problem.²⁰ In USA, 15-30% of general population has been reported to experience bloating.^{16,21,22} Also in Asia, similar result has been shown (15-23%), suggesting that the prevalence of bloating is not interracially different.²³ Though the data for FB alone are relatively little, women typically have higher rates of bloating than men according to the reports of IBS.^{16,21,22} This relevance between female gender and bloating has long been suggested and the hormonal effect in connection with menstrual cycle is regarded as one of the possible explanation.^{24,25} Besides, there are some reports of obese people experiencing more GI symptoms such as abdominal pain or bloating.^{26,27}

Bloating is the second most common reported symptom in patients with IBS following abdominal pain.²⁸ In a study from USA which assessed bloating in 542 IBS patients, 76% of the patients reported that they experienced bloating.²⁹ Other study revealed that more than 90% of patients with IBS suffered from bloating.³⁰ In addition, on comparing constipation dominant IBS (IBS-C) with diarrhea dominant IBS (IBS-D), the prevalence of bloating was higher in IBS-C.³¹

A survey from the USA suggested that more than 65% of patients with bloating rated their symptom as moderate to severe, and 54% of patients complained of decreased daily activity due to bloating. Furthermore, 43% of patients took medication for bloating or needed medication.²²

Table 1. Summary of the Comprehensive Reviews for Abdominal Bloating

Author (yr)	Study aim (method)	Suggested mechanisms of bloating	Treatment strategy
Zar et al ⁵ (2002)	Pathophysiology and treatment for bloating in functional bowel disorders	<ol style="list-style-type: none"> 1. Abnormal gas trapping 2. Fluid retention 3. Food intolerance 4. Altered gut microflora ^c altered abdominal muscle tone, altered mucosal sensitivity, colonic motor abnormalities	<ol style="list-style-type: none"> 1. Limited efficacy; dietary intervention, activated charcoal, simethicone, hypnotherapy 2. No efficacy; antibiotics, probiotics, prokinetics
Houghton et al ⁶ (2005)	Pathophysiology and treatment for bloating in FGIDs	<ol style="list-style-type: none"> 1. Gas accumulation 2. Visceral hypersensitivity 3. Fluid retention ^a motor dysfunction, abnormal gut flora, weak abdominal musculature	<ol style="list-style-type: none"> 1. Some benefit; tegaserod, non-absorbable antibiotics, hypnotherapy, surfactant 2. No efficacy; probiotics, activated charcoal, exercise, weight loss, 5-HT₃ antagonist
Azpiroz et al ⁷ (2005)	Clinical importance, pathophysiology and management of abdominal bloating (literature review from January 1989 to September 2004, based on a PubMed search)	<ol style="list-style-type: none"> 1. Altered abdominal wall activity 2. Abnormal perception 3. Intraluminal contents 4. Impaired gut and gas handling ^b Intraluminal gas volume	<ol style="list-style-type: none"> 1. Some benefit; hypnosis, antidepressant 2. Undetermined; prokinetics, spasmolytics, peppermint oil, gas-reducing substances, exclusion diet, exercise
Agrawal et al ⁸ (2008)	Epidemiology and pathophysiology of abdominal bloating in FGIDs (literature review up to 2006, based on a Medline search)	<ol style="list-style-type: none"> 1. Abnormal gas handling 2. Visceral hypersensitivity 3. Abnormal anterior wall muscular activity ^a carbohydrate intolerance, altered gut flora, small bowel bacterial overgrowth ^b gas excess, altered intestinal transit	<ol style="list-style-type: none"> 1. Effective; prokinetics (tegaserod, neostigmine), antibiotics (neomycin, rifaximin) 2. Worth trying; dietary intervention 3. Little evidence; simethicone, charcoal, antispasmodics, probiotics
Schmulson et al ⁹ (2011)	Treatment for abdominal bloating and distension (literature review up to February 2010 in Medline)	Not available	<ol style="list-style-type: none"> 1. Some efficacy; 5-HT₄ agonist (cisapride, tegaserod), lubiprostone, rifaximin, some probiotics (<i>B. infantis</i> 35624 and <i>B. animalis</i>), antifoaming agents 2. New suggestion; low FODMAPs diet 3. Undetermined; antispasmodics, SSRIs, dopamine antagonist 4. No efficacy; bulking agents, laxatives
Lacy et al ¹⁰ (2011)	Pathophysiology, evaluation, and treatment of bloating and distension	<ol style="list-style-type: none"> 1. Altered gut flora 2. Impaired gas transit 3. Impaired evacuation 4. Abnormal abdominal-diaphragmatic reflexes 5. Abnormal perception ^a psychosocial aspects	<ol style="list-style-type: none"> 1. Some efficacy; exercise, rifaximin, cisapride, neostigmine, tegaserod, lubiprostone, inaclootide 2. New suggestion; diet (low FODMAPs) 3. Undetermined; simethicone, probiotics, TCAs, dopamine antagonist 4. No efficacy; antispasmodics

^apossible mechanisms, ^blittle or no evidence, ^cundetermined.FGIDs, functional gastrointestinal disorders; 5-HT₄, 5-hydroxytryptamine; FODMAPs, fermentable oligo-, di- and mono-saccharides and polyols; SSRIs, selective serotonin reuptake inhibitors; TCAs, Tricyclic antidepressants.

Pathophysiology

Abnormal Gut Microbiota

The GI tract microbiota play an important role in host immune system, and there are more than 500 different species of GI microbiota in adult, which mostly are obligate anaerobes.¹⁰ Only a fraction of these organisms can be cultured; therefore, the understanding of the functions of various microbes in the GI tract is still limited. However, researches over the past decades have shown that altered colonic flora were found in stool samples of patients with IBS.³²⁻³⁴ Parkes et al³⁵ suggested that the GI microbiota can be divided into 2 ecosystems; the luminal bacteria and the mucosa-associated bacteria (Fig. 1). Luminal microbiota form the majority of the GI tract flora, and they play a key role in bloating and flatulence in IBS through carbohydrate fermentation and gas production.³⁵

One comprehensive study of the luminal microbiota in IBS examined the fecal samples of IBS subgroups (diarrhea predominant, constipation predominant and mixed subtype) and the controls using 16S ribosomal RNA (rRNA) sequencing. It has been shown that fecal microbiota are significantly altered in IBS. That is, some patients with IBS seem to have different patterns of colonization with coliforms, such as *lactobacillus* and *bifidobacterium* compared to the controls.³⁶ Similar study from Korea using 16S rRNA gene signatures also has demonstrated significant differences in diversity and dominance between IBS and non-IBS fecal samples.³⁷ In addition, these microbial changes altered protein

and carbohydrate metabolism in the gut.³⁷ A study from Japan also showed higher counts of *Veillonella* ($P = 0.046$) and *Lactobacillus* ($P = 0.031$) in IBS patients than in controls. Besides, they expressed significantly higher levels of acetic acid ($P = 0.049$), propionic acid ($P = 0.025$) and total organic acids ($P = 0.014$) than controls, which is related to symptoms such as abdominal pain, bloating and changes in bowel habits.³⁸ Another study demonstrated that the patients with IBS produced more H_2 but the total gas excretion was similar in both IBS patients and controls.³⁹ This may be associated with alteration in colonic fermentation by hydrogen-consuming bacteria, which may be an important factor in the pathogenesis of IBS.

Collins et al⁴⁰ have proposed that disruption of the balance between the host and intestinal microbiota produces changes in the mucosal immune system from microscopic to overt inflammation and this also results in changes in gut sensory-motor function and immune activity. Besides, these altered microflora may produce differences in fermented gas type and volume, which may be the causes of symptom in patients with bloating.^{36,40,41}

There have been some reports to verify the relationship between the types of gas produced by colonic microflora and bloating. The low producers of methane reported significantly increased bloating and cramping after ingestion of sorbitol and fiber, and the high producers of methane revealed lower prevalence of severe lactulose intolerance than low producers. Hence, the role of methanogenic flora may be important in the pathogenesis of bloating.^{42,43}

Small Intestinal Bacterial Overgrowth

The patients with IBS who specifically complain of bloating have been reported to have increased gas production from bacterial fermentation caused by small intestinal bacterial overgrowth (SIBO). Pimentel and colleagues had established the concept that SIBO might be a major pathogenesis of IBS.⁴⁴ Moreover, several studies found significant improvement of symptoms such as abdominal pain or bloating, when they were treated with antibiotics.⁴⁴⁻⁴⁶ These findings, however, have not been supported by other studies;⁴⁷⁻⁴⁹ they found mildly increased counts of small intestinal bacteria by culture to be more common in IBS, but the breath H_2 concentration was not significantly different between IBS patients and controls.⁴⁸ Also, there was no correlation between bacterial alteration and symptom pattern, and even lactulose breath test was considered as an unreliable method to detect an association between bacterial overgrowth and IBS.^{47,48}




Figure 1. Luminal and mucosal colonic microbiota and their roles in gut homeostasis. Modified from Parkes et al.³⁵

In another study, breath hydrogen concentration was similar in IBS group and control group, and did not correlate with pain ratings in IBS patients, owing to the lack of objective diagnostic measures and inconsistent data.⁵¹

It is unclear whether changes in small bowel bacterial flora could contribute to bloating in IBS patients, thus further studies are required to confirm these observations.

Intestinal Gas Accumulation

In the fasting state, the healthy GI tract contains only about 100 mL of gas distributed almost equally among 6 compartments - stomach, small intestine, ascending colon, transverse colon, descending colon and distal (pelvic) colon. Postprandial volume of gas increases by about 65%, primarily in the pelvic colon.¹² The excessive volume of intestinal gas has been proposed as the likely cause of bloating and distension, and many researchers have attempted to determine this view. A few studies using plain abdominal radiography demonstrated that intestinal gas volume was greater in patients with IBS than in controls (54% vs. 118%), however, the correlation between intra-abdominal gas contents and bloating was poor.^{52,53} The vast majority of studies do not support that excessive gas induces bloating or abdominal pain. Lasser et al⁵⁴ conducted a study using argon washout technique, which demonstrated no differences in the accumulation of intestinal gas between patients with bloating and healthy subjects. More recent studies using CT scans combined with modern imaging analysis software have also shown that excess gas was not associated with abdominal bloating in most patients.³ Thus, these observations suggest that increased volume of gas may not be the main mechanism of bloating, but rather impaired gas transit or distribution are more often the sources of problem.

Altered Gut Motility and Impaired Gas Handling

Various abnormal motility patterns have been described in IBS patients, but none of those parameters can be used as diagnostic markers.⁵⁵ Some authors have suggested that slow transit of food representing alteration in gut motility is related to bloating in IBS-C patients.⁵⁶ Also in a traditional experiment, normal volunteers being made constipated with loperamide, an agent known to slow transit, experienced bloating.⁵⁷ Recently, IBS-C patients with delayed orocecal and colonic transits have shown abdominal distension rather than bloating.¹⁵ Although delayed gastric emptying and slow intestinal transit in IBS-C patients were reported in many Asian studies, there are still controversies to de-

fine these motor disturbances as unique features in Asian IBS patients. Besides, the association between altered gut motility and IBS symptoms is pretty obscure.⁵⁸ A recent study has also suggested that altered colon transit is of no or minor importance for IBS symptoms such as bloating or pain.⁵⁹

However, there are some different points with respect to the intestinal gas handling or transit. In a study by Serra et al,⁶⁰ they have shown that infused gas into the jejunum resulted in distension and abdominal bloating in most of the IBS patients (18 of 20), while only 20% (4 of 20) of control subjects developed symptoms like that. Another study using gas challenge technique has demonstrated that small intestinal gas transit (especially, jejunum) was more prolonged in patients with bloating than in controls, whereas colonic transit was normal.⁶¹ These data support that impaired small intestinal gas handling could be a mechanism of IBS or gas-bloating. Furthermore, a gas challenge test in healthy subjects during blocked rectal gas outflow showed that abdominal distension by girth measurement was similar in the jejunal and rectal infusion experiments, whereas abdominal symptoms including bloating were more significant in jejunal group.⁶² These data indicate that gas related symptom perception is determined by intraluminal gas distribution, whereas abdominal distension depends on the volume of intestinal gas. Besides, the patients with IBS or FB are considered to evacuate intestinal gas less effectively, so that they are more likely to have symptoms of abdominal distension.^{63,64} This aspect of bloating's mechanism has not been considered to be very relevant, but some researchers are interested in this view owing to the observations of anorectal function, especially in patients with constipation. Constipated patients with bloating plus distension exhibited a greater degree of anorectal dysfunction than those without distension. Moreover, self-restrained anal evacuation also increased symptom perception, while impaired gut propulsion caused by intravenous glucagon did not.^{63,64}

Taken together, ineffective anorectal evacuation as well as impaired gas handling may be possible mechanisms of abdominal distension and bloating. However, the data on the link between altered food transit of gut and bloating are not consistent, although they probably account for bloating in some of the IBS patients.^{56,57}

Abnormal Abdominal-diaphragmatic Reflexes

The abdominal cavity is determined by the placement of the walls of abdominal cavity including diaphragm, vertebral column and abdominal wall musculature. Even if there is no increase in

intra-abdominal volume, a change of the position of abdominal cavity components may produce abdominal distension.⁷ Thus, there have been some efforts to evaluate the relationship between bloating and lumbar lordosis or weakened abdominal muscles. In one classic report, Sullivan suggested that the patients with bloating have weak abdominal muscles and frequently had recently gained weight than controls.⁶⁵ But another study measuring upper and lower abdominal wall activities using surface electromyography has suggested that there were no differences in abdominal muscle activities between the patients and the controls.⁶⁶ Moreover, in an early CT study, some IBS patients showed a tendency of lumbar lordosis but not consistent, and a change in lumbar lordosis did not correlate in any way with the changes in abdominal girth. Also, there were no noticeable changes in position of the diaphragm.⁶⁷

Tremolaterra et al⁶⁸ reported that intestinal gas load was associated with a significant increment in abdominal wall muscle activity in healthy subjects. In contrast, the response to gas infusion was impaired in patients with bloating, and rather a paradoxical relaxation of the internal oblique muscles was noted.⁶⁸ Further study using modern CT analysis with electromyography from Barcelona group has provided a novel concept of abnormal abdomino-phrenic reflexes for abdominal bloating and distension. Since then, several studies have demonstrated that abdomino-phrenic dyssynergia is one of main factors for abdominal distension and bloating. In healthy adults, colonic gas infusion increases anterior wall tone and relaxes the diaphragm at the same time. On the contrary, patients with bloating have shown diaphragmatic contraction (descent) and relaxation of the internal oblique muscle with the same gas load.^{4,69,70}

Visceral Hypersensitivity

The sensation of bloating may originate from abdominal viscera in patients with FGIDs, in whom normal stimuli or small variations of gas content within the gut may be perceived as bloating. Indeed, it has been well recognized that the patients with IBS have lower visceral perception threshold than healthy controls,^{1,71} and it has been speculated that this process might be associated with the sensation of bloating. Kellow et al⁷² revealed that threshold for perception of small bowel contraction was lower than normal in some patients with IBS. Also, altered rectal perception assessed by phasic balloon distension has been reported in IBS patients.⁷³ In addition, a gas challenge test proved a role of sensory disturbances in IBS patients,⁶⁰ and recent clinical experiment has demonstrated that bloating without visible distension is

associated with visceral hypersensitivity.⁷⁴

The autonomic nervous system may also contribute to modulation of the visceral sensitivity. Sympathetic activation is known to increase the perception of intestinal distention in FD patients; likewise, autonomic dysfunction could affect the visceral sensitivity in IBS patients.⁷⁵⁻⁷⁷ This mechanism may play a role in bloating. Moreover, it has been proposed that visceral perception may be influenced by cognitive mechanism. That is, IBS patients with bloating pay more attention to their abdominal symptoms, which is a kind of hyper-vigilance.⁷⁸ Also, a report indicated that female patients with IBS had worsening of abdominal pain and bloating during their peri-menstrual phase, at which time heightened rectal sensitivity might have contributed to bloating, but not to distension.^{24,79} Taken together, altered sensory threshold combined with altered conscious perception may explain the mechanism of bloating.

Food Intolerance and Carbohydrate Malabsorption

It is well recognized that dietary habits may be responsible for abdominal symptoms, and there have been efforts to prove the relationship between diet and IBS symptoms. Fiber overload has long been regarded as worsening factor of IBS symptoms through decreased small bowel motility or intraluminal bulking.^{80,81} In addition, lactose intolerance may contribute to symptom development in IBS patients. In the small intestine, disaccharides are split by intestinal enzymes into monosaccharides which are then absorbed. If this process is not carried out, the disaccharide reaches the colon, in turn is split by bacterial enzymes into short chain carbonic acids and gases. Hence, malabsorption of lactose may produce the symptom of bloating in patients with IBS or FB.^{11,82-84} Additionally, a new hypothesis is proposed, by which excessive delivery of highly fermentable but poorly absorbed short chain carbohydrates and polyols (collectively termed FODMAPs; fermentable oligo-, di- and mono-saccharides and polyols) to the small intestine and colon may contribute to the development of GI symptoms. FODMAPs are small molecules that are osmotically active and very rapidly fermentable compared with long-chain carbohydrates. These molecules induce relatively selective bacterial proliferation, especially of *bifidobacterium*, and it has been demonstrated indirectly that these can lead to expansion of bacterial populations in distal small intestine.⁸⁵⁻⁸⁷ Thus, high FODMAP diet has demonstrated prolonged hydrogen production in the intestine, colonic distension by fermentation, increased colonic fluid delivery by osmotic load within the bowel lu-

men, and GI symptom generation.^{88,89}

Intraluminal Contents

Levitt et al⁹⁰ suggested that abdominal bloating might develop without gas retention, but by other gut contents. They had undertaken randomized, double-blind, crossover study of gaseous symptoms by observing the responses of healthy subjects to dietary supplement with lactulose or 2 types of fibers (psyllium or methylcellulose). In lactulose group, gas passages, subjective perception of rectal gas and breath hydrogen excretion were significantly increased, but not in fiber groups. However, the sensation of bloating was increased in all 3 groups. Thus, it has been proposed that increased intra-abdominal bulk, not gaseous filling, might be a cause of abdominal bloating.⁹⁰ In another study, bran accelerated small bowel transit and ascending colon clearance without causing symptom in controls, but small bowel transit has not further been accelerated in IBS patients with bloating. Thus, they speculated that bran might cause increased bulking effect in the colon, which led to the exacerbation of bloating in IBS patients.⁹¹ Francis and Whorwell⁸⁰ even proposed that use of the bran in IBS should be reconsidered, because excessive consumption of bran might give rise to symptoms such as bloating in IBS patients. Although more studies are needed for further understanding of their relationship, it could be possible that intraluminal bulking aggravates the bloating in some IBS patients.

Hard stool/Constipation

Many constipated patients complain of bloating.¹⁴ Also there is a tendency of its being more common in IBS-C patients than IBS-D patients, though it is not statistically significant in some studies.^{31,92-94} Distension of the rectum by retained feces slows small intestinal transit as well as colonic transit, probably explaining the aggravated bloating in constipated patients.^{15,56} Thus it seems reasonable that constipation or hard/lumpy stool induces alteration of gut motility and thus maybe increases bacterial fermentation. In addition, constipation may accelerate bloating by intraluminal bulking effect in the same manner as bran.

Psychological Aspects

Bloating is a frequent complaint of women with IBS. Park et al⁹⁵ proposed that there was a tendency to increase the index of psychological distress when the bloating was more severe. Also, patients with bloating revealed increased anxiety and depression, which allows the hypothesis that psychological distress may contribute to the perceived severity of bloating.⁹⁶ Additionally, in

large population surveys, bloating was significantly related with psychiatric dysfunction such as major depressive disorder, panic disorder and sleeping difficulties.^{97,98} Nevertheless, other studies have failed to demonstrate the relationships between psychological distress and either bloating or distension.^{14,99} However, it is unclear whether or not there is an actual relationship between bloating and psychosocial distress, and further studies are needed to demonstrate it.

Gender and Sex Hormones

In a population based study in USA, female gender was significantly associated with increased symptoms of bloating and distension in IBS, and similar findings have been reported so far.^{21,100-102} Although the question of the gender role in IBS has been raised from many studies, the mechanisms of gender differences in bloating and distension are unclear. Some studies have suggested that bloating is one of the frequent symptoms of menstruation as aforementioned.^{24,25} Hormonal effect has also been speculated, that is, the variation of reproductive hormones throughout the menstrual cycle and after the menopause may influence the gut motility and visceral perception.^{24,79,103} Additionally, difference in symptom expression by gender is presented as a potential explanation.³¹ Although more investigations regarding the underlying mechanisms for these disparities remain to be determined, it seems to be possible to speculate that the hormonal fluctuation may contribute to bloating in female IBS patients.

Treatment

Antibiotics

There has been an increasing acceptance of the use of the antibiotics to treat IBS symptoms, and it is plausible based on the presumption that altered gut flora or SIBO may contribute to gaseous distension or bloating symptom.^{44,104,105}

Although some questions have been raised regarding the validity of the lactulose breath test in diagnosis of SIBO and the possibility of overdiagnosis,¹⁰⁶ much more data support the clinical use of antibiotics in this condition. Specifically, rifaximin, a rifamycin derivative, has largely been studied, and it showed superiority to placebo in relieving bloating in IBS or in patients who were diagnosed as SIBO (Table 2). As rifaximin is a non-absorbable antimicrobial agent, the risk of side effects or emergence of resistant organisms is expected to be low; therefore it is suitable for chronic administration.^{44-46,107,108} Recently, a phase 3 multi-

Table 2. Studies for Rifaximin Treatment in Irritable Bowel Syndrome Patients

Author (yr)	Study design	Diagnostic criteria	IBS subtypes	Female ratio (n) rifaximin vs. placebo	Mean age / Female ratio (n) rifaximin vs. placebo	Drug dosage	Treatment duration (days)	RR for global symptoms (rifaximin, %)	RR for bloating
Sharara et al ¹⁵ (2006)	Double-blind, placebo-controlled, single center	70 met Rome II	All (IBS-D 20.0%, IBS-C 38.3%, IBS-M 41.7%)	42.2/52.4% (63) vs. 38.9/57.4% (61)	400 mg b.i.d.	10	41.3	NA (bloating score; 24.4 → 20.8) (P = 0.001)	
Pimentel et al ¹⁰⁹ (2011)	Double-blind, placebo-controlled, multi-center (TARGET 1)	Rome II	Excluded IBS-C	46.2/76.1% (309) vs. 45.5/70.7% (314)	550 mg t.i.d.	14	40.8	39.5%	
Pimentel et al ¹⁰⁹ (2011)	Double-blind, placebo-controlled, multi-center (TARGET 2)	Rome II	Excluded IBS-C	45.9/72.1% (315) vs. 46.3/70.3% (320)	550 mg t.i.d.	14	40.6	41.0%	
Peralta et al ¹¹¹ (2009)	Observational analysis, single arm, single center	Rome II	All (IBS-D 35.2%, IBS-C 20.4%, IBS-M 44.4%)	NA (54)	1,200 mg/day	7	NA	NA (symptom score; 2.3 → 0.8) (P = 0.003)	
Yang et al ¹¹³ (2008)	Retrospective study, single center	Rome I	NA	NA (84)	1,200 mg/day	NA (follow-up duration; median 11 months)	69.0 (other antibiotics, 44.0)	NA	
Pimentel et al ⁴⁴ (2006)	Double-blind, randomized, placebo-controlled study, 2 centers	Rome I	All (%; NA)	39.1/67.4% (43) vs. 38.2/65.9% (44)	400 mg t.i.d.	10	36.40	NA	
Jolley et al ¹¹² (2011)	Retrospective study, single center	Rome III	All (IBS-D 28.0%, IBS-C 20.0%, IBS-M 15.0%; not reported in 37.0% 1,200 mg/d group) (30.0%, 20.0%, 14.0% and 37.0% respectively; high dose group)	58.0/77.2% (162) vs. 60.0/72.8% (81) (high dose group)	1,200 mg/day vs. 2,400 mg/day (high dose group)	10	49.0 vs. 47.0 (high dose group)	NA	

IBS, irritable bowel syndrome; RR, response rate; NA, not available; IBS-D, IBS with diarrhea; IBS-C, IBS with constipation; IBS-M, mixed IBS.

center trial proved that rifaximin provided significant improvement of IBS symptoms including abdominal pain and bloating in non-constipated IBS patients.¹⁰⁹ Pimentel et al¹¹⁰ also suggested that neomycin normalized lactulose breath test and it contributed to the reduction of bloating in IBS patients. Besides, there have been several retrospective or observational studies, which support the efficacy of rifaximin treatment in IBS patients. Specifically, rifaximin turned the lactulose breath test to negative and significantly reduced the overall symptom scores.¹¹¹ Also, high dose rifaximin treatment (2,400 mg/day) was proved to be effective in patients who had incomplete response to usual dose of rifaximin.¹¹² Another retrospective study has revealed that rifaximin is superior to other antibiotics, such as neomycin, doxycycline, amoxicillin/clavulanate and ciprofloxacin in patients with IBS.¹¹³ Therefore, antibiotics like rifaximin could be considered as a short course therapeutic regimen for bloating, mainly in IBS without constipation. Further studies are needed to determine how long these antibiotics should be given and whether drug resistance will be a problem.

Probiotics

Alteration in gut microbiota may produce or perpetuate the symptoms of bloating or distension, therefore many researchers postulated that modification of the gut microflora could improve gas related symptoms.³⁹ One placebo-controlled study conducted in IBS patients revealed a beneficial effect of *Bifidobacterium infantis* and they suggested immune-modulating role of that organism.¹¹⁴ Another multicenter, clinical trial in women with IBS also showed that *B. infantis* relieved many of the symptoms of IBS, but just at a specific dosage (1×10^8 CFU/mL).¹¹⁵ In addition, more recent experiments have shown that some probiotic strains significantly alleviate the bloating as well as overall symptoms.^{116,117} One study from Korea has shown that multi-species probiotics given to IBS patients are effective in the relief of bloating, albeit not statistically significant over placebo.¹¹⁸ In the most recent meta-analysis of probiotics for lower GI symptoms, specific probiotics are recommended in the management of bloating in IBS patients as moderate grade of evidence along with 70% level of agreement.¹¹⁹

On the contrary, many other studies have failed to prove favorable effects of the probiotics. Kim et al¹²⁰ evaluated the effectiveness of VSL#3, a composite probiotic containing *Bifidobacterium*, *Lactobacillus* and *Streptococcus* in IBS patients. VSL#3 reduced flatulence scores and retarded colonic transit without altering bowel function, but there was no significant reduction in

bloating score with VSL#3.¹²⁰ Some experimental studies from Korea showed a trend towards amelioration of bloating, but failed to prove beneficial effect over placebo.^{121,122} In addition, several other studies using *Lactobacillus* strains reported unfavorable effect on bloating in IBS (Table 3).^{41,123,124} Most of the studies were relatively small and there have been inconsistent results regarding the efficacy of probiotics on bloating. Hence, larger and well-designed trials are needed to prove whether the probiotics are reasonable to treat patients with bloating.

Prokinetics

Prokinetics have been used in the treatment of bloating in FD traditionally, in spite of the weak evidence for correlation between symptoms and underlying pathophysiological mechanisms. A number of studies have shown the beneficial effect of prokinetics such as dopamine antagonist, muscarinic antagonist, and serotonergic agents in FD, but studies conducted in IBS patients are relatively rare (Table 4). Several studies have suggested that cisapride, a 5-hydroxytryptamine 4 (5-HT₄) receptor agonist, significantly improves postprandial bloating in FD patients.^{125,126} Levosulpiride turned out to be as effective as cisapride in the treatment of FD symptoms, such as bloating.¹²⁷ Acotiamide, a novel prokinetic agent, also provided relief of bloating in FD patients in a small study.¹²⁸

Some researchers tried to investigate the efficacy of tegaserod, a selective 5-HT₄ partial agonist, in patients with IBS-C whose main symptom was not diarrhea, and they suggested significant relief in bloating with tegaserod.¹²⁹⁻¹³² However, tegaserod was withdrawn from the market in 2007 due to possible adverse cardiovascular effects. Additionally, neostigmine, a potent prokinetic drug, also exhibited significant effect in reducing objective abdominal distension as well as bloating in IBS or FB patients.¹³³ On the other hand, some other studies do not agree with the favorable action of prokinetics in IBS. One double-blind trial suggested that cisapride was not superior to placebo in the treatment of bloating and other abdominal symptoms of IBS, but it reduced difficulty of stool passage.¹³⁴ However, cisapride was also removed from the market due to the side effect. Another study conducted in IBS patients to evaluate the efficacy of domperidone showed no significant improvement of bloating.¹³⁵ In a small experimental study, pyridostigmine reduced the severity of bloating, but it did not reach the statistical difference across groups.³ Although there are conflicting evidences regarding the effect of prokinetics on bloating, some of the prokinetics could be a treatment option for bloating.

Table 3. Summary of Studies for Probiotics in Irritable Bowel Syndrome

Author (yr)	Study design	Criteria	IBS subtypes	Sample size	Probiotic strains (daily dose)	Duration (weeks)	Results
Nobaek et al ¹¹ (2000)	RCT	Rome I	All (IBS-C, IBS-D, IBS-M)	60	<i>L. plantarum</i> DSMI 9843 (299V) (5×10^7 CFU/mL)	4	Flatulence; improved in test group ($P < 0.05$) Pain, bloating; no benefit over placebo
O'Mahony et al ¹⁴ (2005)	RCT	Rome II	All	75	<i>L. salivarius</i> UCC 4331 or <i>B. infantis</i> 35624	8	Abdominal pain, bowel movement difficulty; significantly improved in <i>B. infantis</i> group (all $P < 0.05$) Bloating; improved in <i>B. infantis</i> group ($P < 0.05$) No benefit in <i>L. salivarius</i> group
Whorwell et al ¹⁵ (2006)	RCT	Rome II	All	362	<i>B. infantis</i> 35624 (3 groups; 1×10^6 , 1×10^8 or 1×10^{10} CFU/mL)	4	Abdominal pain, bloating, incomplete evacuation, straining, passage of gas; improved only in 1×10^8 group (all $P < 0.05$)
Kim et al ¹²⁰ (2005)	RCT	Rome II	All	48	VSL#3	4-8	Flatulence; improved in test group ($P < 0.01$) Failed to show improvement in bloating
Niv et al ¹²⁴ (2005)	RCT	Rome II	All	54	<i>L. reuteri</i> ATCC 55730 (1×10^8 CFU/tablet, twice a day)	26	Abdominal pain, bloating, gases, visible abdominal swellings, GSS; improved, but no benefit over placebo
Guglielmetti et al ¹¹⁶ (2011)	RCT	Rome III	All	122	<i>B. bifidum</i> MIMBb/75 (1×10^8 CFU/capsule, once a day)	4	Pain, distension/bloating, GSS; significantly reduced in test group (all $P < 0.0001$)
Choi et al ¹²¹ (2011)	RCT	Rome II	IBS-D, IBS-M	67	<i>S. boulardii</i> (2×10^{11} cells/day)	4	Quality of life; significant improvement in test group ($P < 0.05$)
Kicha et al ¹²² (2012)	RCT	Rome III	IBS-D	50	A mixture of <i>L. acidophilus</i> , <i>L. plantarum</i> , <i>L. rhamnosus</i> , <i>B. breve</i> , <i>B. lactis</i> , <i>B. longum</i> and <i>S. thermophilus</i> (1×10^{10} CFU/day)	8	Adequate relief of overall IBS symptoms in test group ($P < 0.05$) Bloating; no benefit over placebo
Ducrotté et al ¹¹⁷ (2012)	RCT	Rome III	All	214	<i>L. plantarum</i> DSMI 9843 (299V) (1×10^8 CFU/day)	4	Abdominal pain, bloating; improved in test group (all $P < 0.05$)
Yoon et al ¹¹⁸ (2013)	RCT	Rome III	All	49	A mixture of <i>B. longum</i> , <i>B. bifidum</i> , <i>B. lactis</i> , <i>L. acidophilus</i> , <i>L. rhamnosus</i> and <i>S. thermophilus</i> (5×10^9 cells/capsule, twice daily)	4	GSS; significantly relieved in test group ($P = 0.03$) Abdominal pain, bloating; improved, but no statistical significance over placebo

IBS, irritable bowel syndrome; RCT, randomized controlled trial; IBS-C, IBS with constipation; IBS-D, IBS with diarrhea; IBS-M, mixed IBS; *L. plantarum*, *Lactobacillus plantarum*; *L. salivarius*, *Lactobacillus salivarius*; *B. infantis*, *Bifidobacterium infantis*; *L. reuteri*, *Lactobacillus reuteri*; *B. bifidum*, *Bifidobacterium bifidum*; *S. boulardii*, *Saccharomyces boulardii*; *L. acidophilus*, *Lactobacillus acidophilus*; *L. rhamnosus*, *Lactobacillus rhamnosus*; *B. lactis*, *Bifidobacterium lactis*; *B. longum*, *Bifidobacterium longum*, *S. thermophilus*, *Streptococcus thermophilus*; GSS, global symptom score.

Table 4. Summary of Studies for Prokinetics in Irritable Bowel Syndrome

Author (yr)	Study design	Diagnostic Criteria	IBS subtypes	Sample size	Prokinetics used (daily dose)	Duration (wk)	Results
Schütze et al ¹³⁴ (1997)	RCT	Rome I	IBS-C	96	Cisapride (5 mg t.i.d., titrated to 10 mg t.i.d. if no response after 4 wk)	12	Bloating, GSS; not superior to placebo Difficulty of stool passage; significant improvement in test group ($P < 0.05$)
Müller-Lissner et al ¹²⁰ (2001)	RCT	Rome I	All (IBS-D, IBS-C, IBS-M)	881	Tegaserod (2 mg or 6 mg b.i.d.)	12	Abdominal pain/discomfort; improved in both test groups ($P < 0.05$), more consistent efficacy over time in higher dose group Bloating; favorable trend in reduction in both test groups
Novick et al ¹³⁰ (2002)	RCT	Rome I	All	1,519 (all female)	Tegaserod (6 mg b.i.d.)	12	Abdominal pain, bloating, stool consistency, GSS; improved in test group (all $P < 0.05$)
Kellow et al ¹³¹ (2003)	RCT	Rome II	Excluded IBS-D	520	Tegaserod (6 mg b.i.d.)	12	GSS; improved in test group ($P < 0.0001$) Abdominal pain, bloating, hard stools; improved in test group (all $P < 0.05$)
Tack et al ¹³² (2005)	RCT	Rome II	IBS-C	2,660 (all female)	Tegaserod (6 mg b.i.d.)	4	Abdominal pain, bloating, constipation; improved in test group (all $P < 0.0001$)
Chey et al ¹⁶⁹ (2008)	RCT	Rome II	IBS-C, IBS-M	661 (all female)	Tegaserod (6 mg b.i.d.)	4	Overall symptom; relieved in test group ($P < 0.001$) Bloating; no benefit over placebo
George et al ¹⁷⁰ (2008)	RCT	Rome II	IBS-C	510	Renzapride (1 mg, 2 mg or 4 mg o.d.)	12	Stool frequency, stool consistency; improved in 2 mg and 4 mg o.d. groups (all $P < 0.05$) Bloating; reduction in 1 mg o.d. group ($P = 0.01$)

RCT, randomized controlled trial; GSS, global symptom score; IBS, irritable bowel syndrome; IBS-D, IBS with diarrhea; IBS-C, IBS with constipation; IBS-M, mixed IBS.

Table 5. Summary of Studies for Spasmolytics in Irritable Bowel Syndrome

Author (yr)	Study design	Diagnostic criteria	IBS subtypes	Sample size	Spasmolytics used (daily dose)	Duration (weeks)	Results
Battaglia et al ¹⁴¹ (1998)	RCT	Drossman's criteria for IBS	NA	325	Otilonium bromide (40 mg t.i.d.)	15	Abdominal pain, distension; significant reduction (all $P < 0.05$)
Dobrilla et al ¹⁷¹ (1990)	RCT	Clinical diagnosis and investigations	NA	70	Cimetropium (50 mg t.i.d.)	12	Severity and frequency of abdominal pain; significantly decreased ($P = 0.0005$ and 0.001 , respectively) Abdominal distension; decreased, but not statistically significant ($P = 0.055$)
Glende et al ¹⁴³ (2002)	RCT	Rome I	All (IBS-D, IBS-C, IBS-M)	378	Otilonium bromide (40 mg t.i.d.)	15	Abdominal pain, distension; improved in test group (all $P < 0.05$)
Mitchel et al ¹⁷² (2002)	RCT	Rome II	All	107	Alverine (150 mg t.i.d.)	12	Abdominal pain, bloating, general well-being; failed to show benefit over placebo
Clave et al ¹⁴⁰ (2011)	RCT	Rome II	All	356	Otilonium bromide (40 mg t.i.d.)	15	Abdominal pain ($P = 0.03$), bloating ($P = 0.02$), global efficacy ($P = 0.047$); significant benefit over placebo
Chang et al ¹⁷³ (2011)	RCT	Rome II	All	117	Otilonium bromide (40 mg t.i.d.) Ornimeverine (100 mg t.i.d.)	8	Abdominal pain, flatulence, bloating; global assessment; relieved in both treatment group (all $P < 0.05$)

NA, not available; RCT, randomized controlled trial; IBS, irritable bowel syndrome; IBS-D, IBS with diarrhea; IBS-C, IBS with constipation; IBS-M, mixed IBS.

simethicone significantly relieved the frequency and severity of gas-related symptoms in patients with FGID. Holtmann¹⁵⁴ also conducted a randomized, placebo-controlled trial of simethicone, and suggested that simethicone was significantly better than placebo for overall symptom control in FD patients, in spite of unfavorable effect for bloating. More recently, prospective, multi-center trial to demonstrate a favorable action of activated charcoal-simethicone combination therapy revealed that the severity of fullness and bloating was significantly decreased in the therapy group compared with placebo (Table 7).¹⁵⁵

Stimulants of Fluid Secretion

Lubiprostone and linaclotide are novel agents recently approved by the USA Food and Drug Administration, that enhance fluid secretion into the gut lumen and accelerate intestinal transit. These properties are considered to play a role in treatment of constipation, thus a number of clinical trials focusing in the chronic constipation or IBS-C have been conducted (Table 8). In 2 phase III trials, lubiprostone significantly improved the overall IBS symptoms including bloating in IBS-C.^{156,157} Several multi-center, randomized trials of linaclotide in chronic constipation or IBS-C also demonstrated the beneficial effect in relieving abdominal bloating.^{158,159} Thus so far, these 2 novel drugs offer a reasonable therapeutic approach for bloating mainly in IBS-C and functional constipation patients.

Antidepressants

Antidepressants such as selective serotonin reuptake inhibitor (SSRI) or tricyclic antidepressant (TCA) are believed to alleviate symptoms in FGIDs on the basis of their visceral analgesic properties as well as psychological aspects. However, one small study conducted in IBS patients with visceral hypersensitivity revealed that fluoxetine, one of the SSRI, was effective only in abdominal pain, not in other symptoms such as bloating.¹⁶⁰ Paroxetine was also evaluated in IBS patients who did not respond to high fiber diet. Overall well-being sensation was improved more with paroxetine than with placebo, but abdominal bloating was not.¹⁶¹ However, there were also some positive results. That is, the SSRI, citalopram significantly improved abdominal bloating compared with placebo, though the therapeutic effect was independent of the effect on anxiety, depression and colonic sensitivity (Table 9).¹⁶² Taken together, the results for the treatment of bloating and distension with antidepressants are partly contradictory, and there were few studies which explained the effect of TCA on bloating. Hence, larger, well-designed trials

with SSRIs and TCAs are warranted to identify the efficacy of these drugs on bloating and distension.

Opioid Agents

There have been a few reports that propose the usefulness of opioid agents in IBS patients (Table 10). The kappa receptor agonist, fedotozine has been shown to increase the threshold of perception to colonic distension and reduce visceral sensation.¹⁶³ It has also demonstrated its superiority to placebo in relieving postprandial fullness and bloating in FD patients.¹⁶⁴ In a phase II trial, asimadoline, a novel kappa-opioid agonist, has yielded excellent efficacy results on pain and bloating in IBS-D patients.¹⁶⁵ A small study has suggested that naloxone is beneficial in reducing the bloating score in IBS-C or IBS-M patients, but there were no significant differences in the results with naloxone and placebo.¹⁶⁶ Though a recent review also makes a suggestion of the use of opioid agonists in IBS-D patients,¹⁶⁷ their role in bloating is uncertain to date.

Summary

Abdominal bloating is a frequent and bothersome, but poorly understood clinical problem. The terms of bloating and distension are often confused, but these 2 symptoms should be considered to be separate, as they probably have different pathophysiological mechanisms. The possible mechanisms of bloating are complex and maybe various mechanisms are combined in symptom generation. Important mechanisms of bloating are impaired gas handling and hypersensitivity. Also, recent evidences are beginning to emphasize that patients with bloating may have an altered bacterial flora, SIBO, and abdomino-phrenic dyssynergia. Other less-established factors for bloating are food intolerance, intraluminal bulking and psychological factors (Fig. 2).

On approaching to the treatment of abdominal bloating, clinicians should consider a heterogeneous condition produced by a combination of various mechanisms. Currently, there is no treatment which has indisputably proven to be effective for bloating. Treatment strategy for bloating may include pharmacologic approach, dietary modification, and psychological therapy. Taken together, 5-HT₄ agonists, antibiotics such as rifaximin, some probiotics, and also novel agents, lubiprostone and linaclotide are substantiated to be effective in some degree in the treatment of bloating. Dietary intervention with low FODMAP is also newly qualified treatment option. Though the evidence is weak, antifoaming agents and antidepressants could be consid-




Figure 2. Potential mechanisms behind bloating and visible distention in functional gastrointestinal disorders. Modified from Simrén.¹⁶⁸ CNS, central nervous system; ENS, enteric nervous system; GI, gastrointestinal.

ered in some patients. Of course a careful history and physical examination should be the first step and reassurance and education are likewise important.

Though the whole mechanism and treatment strategies are yet to be fully elucidated, this article proposes a framework for assessing and managing the patients with bloating.

References

1. Bradette M, Delvaux M, Staumont G, Fioramonti J, Bueno L, Frexinos J. Evaluation of colonic sensory thresholds in IBS patients using a barostat. Definition of optimal conditions and comparison with healthy subjects. *Dig Dis Sci* 1994;39:449-457.
2. Issa B, Wafaei NA, Whorwell PJ. Abdominal bloating and distension: what is the role of the microbiota. *Dig Dis Sci* 2012;57:4-8.
3. Accarino A, Perez F, Azpiroz F, Quiroga S, Malagelada JR. Intestinal gas and bloating: effect of prokinetic stimulation. *Am J Gastroenterol* 2008;103:2036-2042.
4. Accarino A, Perez F, Azpiroz F, Quiroga S, Malagelada JR. Abdominal distention results from caudo-ventral redistribution of contents. *Gastroenterology* 2009;136:1544-1551.
5. Zar S, Benson MJ, Kumar D. Review article: bloating in functional bowel disorders. *Aliment Pharmacol Ther* 2002;16:1867-1876.
6. Houghton LA, Whorwell PJ. Towards a better understanding of abdominal bloating and distension in functional gastrointestinal disorders. *Neurogastroenterol Motil* 2005;17:500-511.
7. Azpiroz F, Malagelada JR. Abdominal bloating. *Gastroenterology* 2005;129:1060-1078.
8. Agrawal A, Whorwell PJ. Review article: abdominal bloating and distension in functional gastrointestinal disorders-epidemiology and exploration of possible mechanisms. *Aliment Pharmacol Ther* 2008;27:2-10.
9. Schmulson M, Chang L. Review article: the treatment of functional abdominal bloating and distension. *Aliment Pharmacol Ther* 2011;33:1071-1086.
10. Lacy BE, Gabbard SL, Crowell MD. Pathophysiology, evaluation, and treatment of bloating: hope, hype, or hot air? *Gastroenterol Hepatol (NY)* 2011;7:729-739.
11. Fernández-Bañares F, Rosinach M, Esteve M, Forné M, Espinós JC, María Viver J. Sugar malabsorption in functional abdominal bloating: a pilot study on the long-term effect of dietary treatment. *Clin Nutr* 2006;25:824-831.
12. Azpiroz F, Levitt M. Intestinal Gas. In: Feldman M, Friedman LS, Brand LJ, eds. *Sleisenger and Fordtran's. Gastrointestinal and liver disease: pathophysiology, diagnosis, management*. Philadelphia: Saunders 2010:233-240.
13. Páramo Hernández DB. Bloating and abdominal distention: Just gas? A look in the direction of physiology. *Rev Col Gastroenterol* 2011;26:269-272.
14. Houghton LA, Lea R, Agrawal A, Reilly B, Whorwell PJ. Relationship of abdominal bloating to distention in irritable bowel syndrome and effect of bowel habit. *Gastroenterology* 2006;131:1003-1010.
15. Agrawal A, Houghton LA, Reilly B, Morris J, Whorwell PJ. Bloating and distension in irritable bowel syndrome: the role of gastrointestinal transit. *Am J Gastroenterol* 2009;104:1998-2004.
16. Jiang X, Locke GR 3rd, Choung RS, Zinsmeister AR, Schleck CD, Talley NJ. Prevalence and risk factors for abdominal bloating and visible distention: a population-based study. *Gut* 2008;57:756-763.
17. Thompson WG, Longstreth GF, Drossman DA, Heaton KW, Irvine EJ, Müller-Lissner SA. Functional bowel disorders and

- functional abdominal pain. *Gut* 1999;45(suppl 2):II43-II47.
18. Drossman DA, Corazziari E, Tally NJ, Thompson W, Whitehead W. Rome II. The functional gastrointestinal disorders. Diagnosis, pathophysiology and treatment: amultinational consensus, 2nd ed. McLean, VA: Degnon Associates Inc 2000.
 19. Longstreth GF, Thompson WG, Chey WD, Houghton LA, Mearin F, Spiller RC. Functional bowel disorders. *Gastroenterology* 2006;130:1480-1491.
 20. Alvarez WC. Hysterical type of nongaseous abdominal bloating. *Arch Intern Med (Chic)* 1949;84:217-245.
 21. Ringel Y, Williams RE, Kalilani L, Cook SF. Prevalence, characteristics, and impact of bloating symptoms in patients with irritable bowel syndrome. *clin Gastroenterol Hepatol* 2009;7:68-72.
 22. Sandler R, Stewart W, Liberman J, Ricci J, Zorich N. Abdominal pain, bloating, and diarrheain the united states. *Dig Dis Sci* 2000;45:1166-1171.
 23. Ho KY, Kang JY, Seow A. Prevalence of gastrointestinal symptoms in a multiracial Asian population, with particular reference to reflux-type symptoms. *Am J Gastroenterol* 1998;93:1816-1822.
 24. Heitkemper MM, Cain KC, Jarrett ME, Burr RL, Hertig V, Bond EF. Symptoms across the menstrual cycle in women with irritable bowel syndrome. *Am J Gastroenterol* 2003;98:420-430.
 25. Huerta-Franco MR, Malacara JM. Association of physical and emotional symptoms with the menstrual cycle and life-style. *J Reprod Med* 1993;38:448-454.
 26. Delgado-Aros S, Locke GR 3rd, Camilleri M, et al. Obesity is associated with increased risk of gastrointestinal symptoms: a population-based study. *Am J Gastroenterol* 2004;99:1801-1806.
 27. Talley NJ, Quan C, Jones MP, Horowitz M. Association of upper and lower gastrointestinal tract symptoms with body mass index in an Australian cohort. *Neurogastroenterol Motil* 2004;16:413-419.
 28. Maxton DG, Morris JA, Whorwell PJ. Ranking of symptoms by patients with the irritable bowel syndrome. *BMJ* 1989;299:1138.
 29. Chang L, Lee OY, Naliboff B, Schmulson M, Mayer EA. Sensation of bloating and visible abdominal distension in patients with irritable bowel syndrome. *Am J Gastroenterol* 2001;96:3341-3347.
 30. Houghton LA. Bloating in constipation: relevance of intraluminal gas handling. *Best Pract Res Clin Gastroenterol* 2011;25:141-150.
 31. Talley NJ, Dennis EH, Schettler-Duncan VA, Lacy BE, Olden KW, Crowell MD. Overlapping upper and lower gastrointestinal symptoms in irritable bowel syndrome patients with constipation or diarrhea. *Am J Gastroenterol* 2003;98:2454-2459.
 32. Balsari A, Ceccarelli A, Dubini F, Fesce E, Poli G. The fecal microbial population in the irritable bowel syndrome. *Microbiologica* 1982;5:185-194.
 33. Malinen E, Rinttilä T, Kajander K, et al. Analysis of the fecal microbiota of irritable bowel syndrome patients and healthy controls with real-time PCR. *Am J Gastroenterol* 2005;100:373-382.
 34. Lee BJ, Bak YT. Irritable bowel syndrome, gut microbiota and probiotics. *J Neurogastroenterol Motil* 2011;17:252-266.
 35. Parkes GC, Brostoff J, Whelan K, Sanderson JD. Gastrointestinal microbiota in irritable bowel syndrome: their role in its pathogenesis and treatment. *Am J Gastroenterol* 2008;103:1557-1567.
 36. Kassinen A, Krogius-Kurikka L, Mäkiyuokko H, et al. The fecal microbiota of irritable bowel syndrome patients differs significantly from that of healthy subjects. *Gastroenterology* 2007;133:24-33.
 37. Ponnusamy K, Choi JN, Kim J, Lee SY, Lee CH. Microbial community and metabolomic comparison of irritable bowel syndrome faeces. *J Med Microbiol* 2011;60(Pt 6):817-827.
 38. Tana C, Umesaki Y, Imaoka A, Handa T, Kanazawa M, Fukudo S. Altered profiles of intestinal microbiota and organic acids may be the origin of symptoms in irritable bowel syndrome. *Neurogastroenterol Motil* 2010;22:512-519, e114-e115.
 39. King TS, Elia M, Hunter JO. Abnormal colonic fermentation in irritable bowel syndrome. *Lancet* 1998;352:1187-1189.
 40. Collins SM, Denou E, Verdu EF, Bercik P. The putative role of the intestinal microbiota in the irritable bowel syndrome. *Dig Liver Dis* 2009;41:850-853.
 41. Nobaek S, Johansson M-L, Molin G, Ahrné S, Jeppsson B. Alteration of intestinal microflora is associated with reduction in abdominal bloating and pain in patients with irritable bowel syndrome. *Am J Gastroenterol* 2000;95:1231-1238.
 42. Kajs TM, Fitzgerald JA, Buckner RY, et al. Influence of a methanogenic flora on the breath H₂ and symptom response to ingestion of sorbitol or oat fiber. *Am J Gastroenterol* 1997;92:89-94.
 43. Vernia P, Camillo MD, Marinaro V, Caprilli R. Effect of predominant methanogenic flora on the outcome of lactose breath test in irritable bowel syndrome patients. *Eur J Clin Nutr* 2003;57:1116-1119.
 44. Pimentel M, Park S, Mirocha J, Kane SV, Kong Y. The effect of a nonabsorbed oral antibiotic (rifaximin) on the symptoms of the irritable bowel syndrome: a randomized trial. *Ann Intern Med* 2006;145:557-563.
 45. Sharara AI, Aoun E, Abdul-Baki H, Mounzer R, Sidani S, ElHajj I. A randomized double-blind placebo-controlled trial of rifaximin in patients with abdominal bloating and flatulence. *Am J Gastroenterol* 2006;101:326-333.
 46. Di Stefano M, Strocchi A, Malservisi S, Veneto G, Ferrieri A, Corazza GR. Non-absorbable antibiotics for managing intestinal gas production and gas-related symptoms. *Aliment Pharmacol Ther* 2000;14:1001-1008.
 47. Walters B, Vanner SJ. Detection of bacterial overgrowth in IBS using the lactulose H₂ breath test: comparison with ¹⁴C-D-xylose and healthy controls. *Am J Gastroenterol* 2005;100:1566-1570.
 48. Posserud I, Stotzer PO, Björnsson ES, Abrahamsson H, Simrén M. Small intestinal bacterial overgrowth in patients with irritable bowel syndrome. *Gut* 2007;56:802-808.
 49. Riordan SM, McIver CJ, Duncombe VM, Thomas MC, Nagree A, Bolin TD. Small intestinal bacterial overgrowth and the irritable bowel syndrome. *Am J Gastroenterol* 2001;96:2506-2508.
 50. Haderstorfer B, Psycholgin D, Whitehead WE, Schuster MM. Intestinal gas production from bacterial fermentation of undigested carbohydrate in irritable bowel syndrome. *Am J Gastroenterol* 1989;84:375-378.
 51. Grace E, Shaw C, Whelan K, Andreyev HJ. Review article: small intestinal bacterial overgrowth-prevalence, clinical features, current and developing diagnostic tests, and treatment. *Aliment Pharmacol Ther* 2013;38:674-688.
 52. Chami TN, Schuster MM, Bohlman ME, Pulliam TJ, Kamal N, Whitehead WE. A simple radiologic method to estimate the quantity of bowel gas. *Am J Gastroenterol* 1991;86:599-602.

53. Koide A, Yamaguchi T, Odaka T, et al. Quantitative analysis of bowel gas using plain abdominal radiograph in patients with irritable bowel syndrome. *Am J Gastroenterol* 2000;95:1735-1741.
54. Lasser RB, Bond JH, Levitt MD. The role of intestinal gas in functional abdominal pain. *N Engl J Med* 1975;293:524-526.
55. Drossman DA, Camilleri M, Mayer EA, Whitehead WE. AGA technical review on irritable bowel syndrome. *Gastroenterology* 2002;123:2108-2131.
56. Cann PA, Read NW, Brown C, Hobson N, Holdsworth CD. Irritable bowel syndrome: relationship of disorders in the transit of a single solid meal to symptom patterns. *Gut* 1983;24:405-411.
57. Kellow JE, Phillips SF. Altered small bowel motility in irritable bowel syndrome is correlated with symptoms. *Gastroenterology* 1987;92:1885-1893.
58. Lee OY. Asian motility studies in irritable bowel syndrome. *J Neurogastroenterol Motil* 2010;16:120-130.
59. Tornblom H, Van Oudenhove L, Sadik R, Abrahamsson H, Tack J, Simrén M. Colonic transit time and IBS symptoms: what's the link? *Am J Gastroenterol* 2012;107:754-760.
60. Serra J, Azpiroz F, Malagelada JR. Impaired transit and tolerance of intestinal gas in the irritable bowel syndrome. *Gut* 2001;48:14-19.
61. Salvioli B, Serra J, Azpiroz F, et al. Origin of gas retention and symptoms in patients with bloating. *Gastroenterology* 2005;128:574-579.
62. Harder H, Serra J, Azpiroz F, Passos MC, Aguadé S, Malagelada JR. Intestinal gas distribution determines abdominal symptoms. *Gut* 2003;52:1708-1713.
63. Serra J, Azpiroz F, Malagelada JR. Mechanisms of intestinal gas retention in humans: impaired propulsion versus obstructed evacuation. *Am J Physiol Gastrointest Liver Physiol* 2001;281:G138-G143.
64. Shim L, Prott G, Hansen RD, Simmons LE, Kellow JE, Malcolm A. Prolonged balloon expulsion is predictive of abdominal distension in bloating. *Am J Gastroenterol* 2010;105:883-887.
65. Sullivan SN. A prospective study of unexplained visible abdominal bloating. *N Z Med J* 1994;107:428-430.
66. McManis PG, Newall D, Talley NJ. Abdominal wall muscle activity in irritable bowel syndrome with bloating. *Am J Gastroenterol* 2001;96:1139-1142.
67. Maxton DG, Martin DF, Whorwell PJ, Godfrey M. Abdominal distension in female patients with irritable bowel syndrome: exploration of possible mechanisms. *Gut* 1991;32:662-664.
68. Tremolaterra F, Villoria A, Azpiroz F, Serra J, Aguadé S, Malagelada JR. Impaired viscerosomatic reflexes and abdominal-wall dystonia associated with bloating. *Gastroenterology* 2006;130:1062-1068.
69. Villoria A, Azpiroz F, Burri E, Cisternas D, Soldevilla A, Malagelada JR. Abdomino-phrenic dyssynergia in patients with abdominal bloating and distension. *Am J Gastroenterol* 2011;106:815-819.
70. Villoria A, Azpiroz F, Soldevilla A, Perez F, Malagelada JR. Abdominal accommodation: a coordinated adaptation of the abdominal wall to its content. *Am J Gastroenterol* 2008;103:2807-2815.
71. Bouin M, Plourde V, Boivin M, et al. Rectal distention testing in patients with irritable bowel syndrome: sensitivity, specificity, and predictive values of pain sensory thresholds. *Gastroenterology* 2002;122:1771-1777.
72. Kellow JE, Eckersley CM, Jones MP. Enhanced perception of physiological intestinal motility in the irritable bowel syndrome. *Gastroenterology* 1991;101:1621-1627.
73. Mertz H, Naliboff B, Munakata J, Niazi N, Mayer EA. Altered rectal perception is a biological marker of patients with irritable bowel syndrome. *Gastroenterology* 1995;109:40-52.
74. Agrawal A, Houghton LA, Lea R, Morris J, Reilly B, Whorwell PJ. Bloating and distention in irritable bowel syndrome: the role of visceral sensation. *Gastroenterology* 2008;134:1882-1889.
75. Park DI, Rhee PL, Kim YH, et al. Role of autonomic dysfunction in patients with functional dyspepsia. *Dig Liver Dis* 2001;33:464-471.
76. Iovino P, Azpiroz F, Domingo E, Malagelada JR. The sympathetic nervous system modulates perception and reflex responses to gut distention in humans. *Gastroenterology* 1995;108:680-686.
77. Azpiroz F, Bouin M, Camilleri M, et al. Mechanisms of hypersensitivity in IBS and functional disorders. *Neurogastroenterol Motil* 2007;19(1 suppl):62-88.
78. Accarino AM, Azpiroz F, Malagelada JR. Attention and distraction: effects on gut perception. *Gastroenterology* 1997;113:415-422.
79. Houghton LA, Lea R, Jackson N, Whorwell PJ. The menstrual cycle affects rectal sensitivity in patients with irritable bowel syndrome but not healthy volunteers. *Gut* 2002;50:471-474.
80. Francis CY, Whorwell PJ. Bran and irritable bowel syndrome: time for reappraisal. *Lancet* 1994;344:39-40.
81. Friedman G. Diet and the irritable bowel syndrome. *Gastroenterol Clin North Am* 1991;20:313-324.
82. Böhmer CJ, Tuynman HA. The clinical relevance of lactose malabsorption in irritable bowel syndrome. *Eur J Gastroenterol Hepatol* 1996;8:1013-1016.
83. Dapoigny M, Stockbrügger RW, Azpiroz F, et al. Role of alimentation in irritable bowel syndrome. *Digestion* 2003;67:225-233.
84. Fernández-Bañares F, Esteve-Pardo M, de Leon R, et al. Sugar malabsorption in functional bowel disease: clinical implications. *Am J Gastroenterol* 1993;88:2044-2050.
85. Roberfroid MB, Van Loo JA, Gibson GR. The bifidogenic nature of chicory inulin and its hydrolysis products. *J Nutr* 1998;128:11-19.
86. O'Leary C, Quigley EM. Small bowel bacterial overgrowth, celiac disease, and IBS: what are the real associations? *Am J Gastroenterol* 2003;98:720-722.
87. Tursi A, Brandimarte G, Giorgetti G. High prevalence of small intestinal bacterial overgrowth in celiac patients with persistence of gastrointestinal symptoms after gluten withdrawal. *Am J Gastroenterol* 2003;98:839-843.
88. Barrett JS, Geary RB, Muir JG, et al. Dietary poorly absorbed, short-chain carbohydrates increase delivery of water and fermentable substrates to the proximal colon. *Aliment Pharmacol Ther* 2010;31:874-882.
89. Ong DK, Mitchell SB, Barrett JS, et al. Manipulation of dietary short chain carbohydrates alters the pattern of gas production and genesis of symptoms in irritable bowel syndrome. *J Gastroenterol*

- Hepatol 2010;25:1366-1373.
90. Levitt MD, Furne J, Olsson S. The relation of passage of gas an abdominal bloating to colonic gas production. *Ann Intern Med* 1996;124:422-424.
 91. Hebden JM, Blackshaw E, D'Amato M, Perkins AC, Spiller RC. Abnormalities of GI transit in bloated irritable bowel syndrome: effect of bran on transit and symptoms. *Am J Gastroenterol* 2002; 97:2315-2320.
 92. Schmulson M, Chang L, Naliboff B, Lee OY, Mayer EA. Correlation of symptom criteria with perception thresholds during rectosigmoid distension in irritable bowel syndrome patients. *Am J Gastroenterol* 2000;95:152-156.
 93. Su AM, Shih W, Presson AP, Chang L. Characterization of symptoms in irritable bowel syndrome with mixed bowel habit pattern. *Neurogastroenterol Motil* 2013 Published Online First: 29 Aug 2013. doi: 10.1111/nmo.12220.
 94. Ringel Y, Williams RE, Kalilani L, Cook SF. Prevalence, characteristics, and impact of bloating symptoms in patients with irritable bowel syndrome. *Clin Gastroenterol Hepatol* 2009;7:68-72.
 95. Park HJ, Jarrett M, Cain K, Heitkemper M. Psychological distress and GI symptoms are related to severity of bloating in women with irritable bowel syndrome. *Res Nurs Health* 2008;31:98-107.
 96. Song JY, Merskey H, Sullivan S, Noh S. Anxiety and depression in patients with abdominal bloating. *Can J Psychiatry* 1993;38:475-479.
 97. Johnsen R, Jacobsen BK, Førde OH. Associations between symptoms of irritable colon and psychological and social conditions and lifestyle. *Br Med J (Clin Res Ed)* 1986;292:1633-1635.
 98. Walker EA, Katon WJ, Jemelka RP, Roy-Bryne PP. Comorbidity of gastrointestinal complaints, depression, and anxiety in the Epidemiologic Catchment Area (ECA) Study. *Am J Med* 1992; 92:26S-30S.
 99. Heitkemper MM, Cain KC, Jarrett ME, Burr RL, Crowell MD, Woods NF. Relationship of bloating to other GI and menstrual symptoms in women with irritable bowel syndrome. *Dig Dis Sci* 2004;49:88-95.
 100. Smith RC, Greenbaum DS, Vancouver JB, et al. Gender differences in manning criteria in the irritable bowel syndrome. *Gastroenterology* 1991;100:591-595.
 101. Schmulson M, Lee OY, Chang L, Naliboff B, Mayer EA. Symptom differences in moderate to severe IBS patients based on predominant bowel habit. *Am J Gastroenterol* 1999;94:2929-2935.
 102. Anbardan SJ, Daryani NE, Fereshtehnejad SM, Taba Taba Vakili S, Keramati MR, Ajdarkosh H. Gender role in irritable bowel syndrome: A comparison of irritable bowel syndrome module (Rome III) between male and female patients. *J Neurogastroenterol Motil* 2012;18:70-77.
 103. Altman G, Cain KC, Motzer S, Jarrett M, Burr R, Heitkemper M. Increased symptoms in female IBS patients with dysmenorrhea and PMS. *Gastroenterol Nurs* 2006;29:4-11.
 104. Spiller RC. Role of infection in irritable bowel syndrome. *J Gastroenterol* 2007;42(suppl 17):41-47.
 105. Pimentel M, Lezzano S. Irritable bowel syndrome: bacterial overgrowth-what's known and what to do. *Curr Treat Options Gastroenterol* 2007;10:328-337.
 106. Yu D, Cheeseman F, Vanner S. Combined oro-caecal scintigraphy and lactulose hydrogen breath testing demonstrate that breath testing detects oro-caecal transit, not small intestinal bacterial overgrowth in patients with IBS. *Gut* 2011;60:334-340.
 107. Corazza GR, Ventrucci M, Strocchi A, et al. Treatment of small intestine bacterial overgrowth with rifaximin, a non-absorbable rifamycin. *J Int Med Res* 1988;16:312-316.
 108. Scarpellini E, Gabrielli M, Lauritano CE, et al. High dosage rifaximin for the treatment of small intestinal bacterial overgrowth. *Aliment Pharmacol Ther* 2007;25:781-786.
 109. Pimentel M, Lembo A, Chey WD, et al. Rifaximin therapy for patients with irritable bowel syndrome without constipation. *N Engl J Med* 2011;364:22-32.
 110. Pimentel M, Chow EJ, Lin HC. Normalization of lactulose breath testing correlates with symptom improvement in irritable bowel syndrome. a double-blind, randomized, placebo-controlled study. *Am J Gastroenterol* 2003;98:412-419.
 111. Peralta S, Cotton C, Doveri T, Almasio PL, Craxi A. Small intestine bacterial overgrowth and irritable bowel syndrome-related symptoms: experience with Rifaximin. *World J Gastroenterol* 2009;15:2628-2631.
 112. Jolley J. High-dose rifaximin treatment alleviates global symptoms of irritable bowel syndrome. *Clin Exp Gastroenterol* 2011;4:43-48.
 113. Yang J, Lee HR, Low K, Chatterjee S, Pimentel M. Rifaximin versus other antibiotics in the primary treatment and retreatment of bacterial overgrowth in IBS. *Dig Dis Sci* 2008;53:169-174.
 114. O'Mahony L, McCarthy J, Kelly P, et al. *Lactobacillus* and *bifidobacterium* in irritable bowel syndrome: symptom responses and relationship to cytokine profiles. *Gastroenterology* 2005;128:541-551.
 115. Whorwell PJ, Altringer L, Morel J, et al. Efficacy of an encapsulated probiotic *Bifidobacterium infantis* 35624 in women with irritable bowel syndrome. *Am J Gastroenterol* 2006;101:1581-1590.
 116. Guglielmetti S, Mora D, Gschwendner M, Popp K. Randomised clinical trial: *Bifidobacterium bifidum* MIMBb75 significantly alleviates irritable bowel syndrome and improves quality of life - a double-blind, placebo-controlled study. *Aliment Pharmacol Ther* 2011;33:1123-1132.
 117. Ducrotté P, Sawant P, Jayanthi V. Clinical trial: *Lactobacillus plantarum* 299v (DSM 9843) improves symptoms of irritable bowel syndrome. *World J Gastroenterol* 2012;18:4012-4018.
 118. Yoon JS, Sohn W, Lee OY, et al. Effect of multi-species probiotics on irritable bowel syndrome: a randomized, double-blind, placebo-controlled trial. *J Gastroenterol Hepatol* Published Online First: 5 Jul 2013. doi: 10.1111/jgh.12322.
 119. Hungin AP, Mulligan C, Pot B, et al. Systematic review: probiotics in the management of lower gastrointestinal symptoms in clinical practice-an evidence-based international guide. *Aliment Pharmacol Ther* Published Online First: 27 Aug 2013. doi: 10.1111/apt.12460.
 120. Kim HJ, Vazquez Roque MI, Camilleri M, et al. A randomized controlled trial of a probiotic combination VSL# 3 and placebo in irritable bowel syndrome with bloating. *Neurogastroenterol Motil* 2005;17:687-696.
 121. Choi CH, Jo SY, Park HJ, Chang SK, Byeon JS, Myung SJ. A randomized, double-blind, placebo-controlled multicenter trial of *Saccharomyces boulardii* in irritable bowel syndrome: effect on quality of life. *J Clin Gastroenterol* 2011;45:679-683.
 122. Ki Cha B, Mun Jung S, Hwan Choi C, et al. The effect of a multi-

- species probiotic mixture on the symptoms and fecal microbiota in diarrhea-dominant irritable bowel syndrome: a randomized, double-blind, placebo-controlled trial. *J Clin Gastroenterol* 2012;46:220-227.
123. O'Sullivan MA, O'Morain CA. Bacterial supplementation in the irritable bowel syndrome. A randomised double-blind placebo-controlled crossover study. *Dig Liver Dis* 2000;32:294-301.
 124. Niv E, Naftali T, Hallak R, Vaisman N. The efficacy of *Lactobacillus reuteri* ATCC 55730 in the treatment of patients with irritable bowel syndrome - a double blind, placebo-controlled, randomized study. *Clin Nutr* 2005;24:925-931.
 125. Champion MC, MacCannell KL, Thomson AB, et al. A double-blind randomized study of cisapride in the treatment of nonulcer dyspepsia. The Canadian Cisapride Nud Study Group. *Can J Gastroenterol* 1997;11:127-134.
 126. Chen JD, Ke MY, Lin XM, Wang Z, Zhang M. Cisapride provides symptomatic relief in functional dyspepsia associated with gastric myoelectrical abnormality. *Aliment Pharmacol Ther* 2000;14:1041-1047.
 127. Mearin F, Rodrigo L, Pérez-Mota A, et al. Levosulpiride and cisapride in the treatment of dysmotility-like functional dyspepsia: a randomized, double-masked trial. *Clin Gastroenterol Hepatol* 2004;2:301-308.
 128. Tack J, Masclee A, Heading R, et al. A dose-ranging, placebo-controlled, pilot trial of acotiamide in patients with functional dyspepsia. *Neurogastroenterol Motil* 2009;21:272-280.
 129. Muller-Lissner SA, Fumagalli I, Bardhan KD, et al. Tegaserod, a 5-HT₄ receptor partial agonist, relieves symptoms in irritable bowel syndrome patients with abdominal pain, bloating and constipation. *Aliment Pharmacol Ther* 2001;15:1655-1666.
 130. Novick J, Miner P, Krause R, et al. A randomized, double-blind, placebo-controlled trial of tegaserod in female patients suffering from irritable bowel syndrome with constipation. *Aliment Pharmacol Ther* 2002;16:1877-1888.
 131. Kellow J, Lee OY, Chang FY, et al. An Asia-Pacific, double blind, placebo controlled, randomised study to evaluate the efficacy, safety, and tolerability of tegaserod in patients with irritable bowel syndrome. *Gut* 2003;52:671-676.
 132. Tack J, Müller-Lissner S, Bytzer P, et al. A randomised controlled trial assessing the efficacy and safety of repeated tegaserod therapy in women with irritable bowel syndrome with constipation. *Gut* 2005;54:1707-1713.
 133. Caldarella MP, Serra J, Azpiroz F, Malagelada JR. Prokinetic effects in patients with intestinal gas retention. *Gastroenterology* 2002;122:1748-1755.
 134. Schütze K, Brandstatter G, Dragosics B, Judmaier G, Hentschel E. Double-blind study of the effect of cisapride on constipation and abdominal discomfort as components of the irritable bowel syndrome. *Aliment Pharmacol Ther* 1997;11:387-394.
 135. Cann PA, Read NW, Holdsworth CD. Oral domperidone: double blind comparison with placebo in irritable bowel syndrome. *Gut* 1983;24:1135-1140.
 136. McKee DP, Quigley EM. Intestinal motility in irritable bowel syndrome: is IBS a motility disorder? Part 1. Definition of IBS and colonic motility. *Dig Dis Sci* 1993;38:1761-1772.
 137. Dumitrascu DL, Stănculete M. The effect of trimebutine on the psychosocial adjustment to illness in the irritable bowel syndrome. *Rom J Intern Med* 2006;44:273-280.
 138. Ford AC, Talley NJ, Spiegel BM, et al. Effect of fibre, antispasmodics, and peppermint oil in the treatment of irritable bowel syndrome: systematic review and meta-analysis. *BMJ* 2008;337:a2313.
 139. Poynard T, Regimbeau C, Benhamou Y. Meta-analysis of smooth muscle relaxants in the treatment of irritable bowel syndrome. *Aliment Pharmacol Ther* 2001;15:355-361.
 140. Clave P, Acalovschi M, Triantafyllidis JK, et al. Randomised clinical trial: otilonium bromide improves frequency of abdominal pain, severity of distention and time to relapse in patients with irritable bowel syndrome. *Aliment Pharmacol Ther* 2011;34:432-442.
 141. Battaglia G, Morselli-Labate AM, Camarri E, et al. Otilonium bromide in irritable bowel syndrome: a double-blind, placebo-controlled, 15-week study. *Aliment Pharmacol Ther* 1998;12:1003-1010.
 142. Luttecke K. A three-part controlled study of trimebutine in the treatment of irritable colon syndrome. *Curr Med Res Opin* 1980;6:437-443.
 143. Glende M, Morselli-Labate AM, Battaglia G, Evangelista S. Extended analysis of a double-blind, placebo-controlled, 15-week study with otilonium bromide in irritable bowel syndrome. *Eur J Gastroenterol Hepatol* 2002;14:1331-1338.
 144. Liu JH, Chen GH, Yeh HZ, Huang CK, Poon SK. Enteric-coated peppermint-oil capsules in the treatment of irritable bowel syndrome: a prospective, randomized trial. *J Gastroenterol* 1997;32:765-768.
 145. Cappello G, Spezzaferro M, Grossi L, Manzoli L, Marzio L. Peppermint oil (Mintoil) in the treatment of irritable bowel syndrome: a prospective double blind placebo-controlled randomized trial. *Dig Liver Dis* 2007;39:530-536.
 146. Darvish-Damavandi M, Nikfar S, Abdollahi M. A systematic review of efficacy and tolerability of mebeverine in irritable bowel syndrome. *World J Gastroenterol* 2010;16:547-553.
 147. Choi YK, Kraft N, Zimmerman B, Jackson M, Rao SS. Fructose intolerance in IBS and utility of fructose-restricted diet. *J Clin Gastroenterol* 2008;42:233-238.
 148. Barrett JS, Gibson PR. Clinical ramifications of malabsorption of fructose and other short-chain carbohydrates. *Pract Gastroenterol* 2007;31:51-65.
 149. Shepherd SJ, Parker FC, Muir JG, Gibson PR. Dietary triggers of abdominal symptoms in patients with irritable bowel syndrome: randomized placebo-controlled evidence. *Clin Gastroenterol Hepatol* 2008;6:765-771.
 150. Gibson PR, Shepherd SJ. Personal view: food for thought-western lifestyle and susceptibility to Crohn's disease. The FODMAP hypothesis. *Aliment Pharmacol Ther* 2005;21:1399-1409.
 151. de Roest RH, Dobbs BR, Chapman BA, et al. The low FODMAP diet improves gastrointestinal symptoms in patients with irritable bowel syndrome: a prospective study. *Int J Clin Pract* 2013;67:895-903.
 152. Rider JA. Experience with the use of a defoaming agent in the treatment of gastrointestinal gas. *Ann N Y Acad Sci* 1968;150:170-177.
 153. Bernstein JE, Kasich AM. A double-blind trial of simethicone in functional disease of the upper gastrointestinal tract. *J Clin*

- Pharmacol 1974;14:617-623.
154. Holtmann G, Gschossmann J, Mayr P, Talley NJ. A randomized placebo-controlled trial of simethicone and cisapride for the treatment of patients with functional dyspepsia. *Aliment Pharmacol Ther* 2002;16:1641-1648.
 155. Lecuyer M, Cousin T, Monnot MN, Coffin B. Efficacy of an activated charcoal-simethicone combination in dyspeptic syndrome: results of a randomized prospective study in general practice. *Gastroenterol Clin Biol* 2009;33:478-484.
 156. Johanson JF, Ueno R. Lubiprostone, a locally acting chloride channel activator, in adult patients with chronic constipation: a double-blind, placebo-controlled, dose - ranging study to evaluate efficacy and safety. *Aliment Pharmacol Ther* 2007;25:1351-1361.
 157. Drossman DA, Chey WD, Johanson JF, et al. Clinical trial: lubiprostone in patients with constipation-associated irritable bowel syndrome-results of two randomized, placebo-controlled studies. *Aliment Pharmacol Ther* 2009;29:329-341.
 158. Quigley EM, Tack J, Chey WD, et al. Randomised clinical trials: linaclotide phase 3 studies in IBS-C - a prespecified further analysis based on European Medicines Agency-specified endpoints. *Aliment Pharmacol Ther* 2013;37:49-61.
 159. Lembo AJ, Schneider HA, Shiff SJ, et al. Two randomized trials of linaclotide for chronic constipation. *N Engl J Med* 2011;365:527-536.
 160. Kuiken SD, Tytgat GN, Boeckxstaens GE. The selective serotonin reuptake inhibitor fluoxetine does not change rectal sensitivity and symptoms in patients with irritable bowel syndrome: a double blind, randomized, placebo-controlled study. *Clin Gastroenterol Hepatol* 2003;1:219-228.
 161. Tabas G, Beaves M, Wang J, Friday P, Mardini H, Arnold G. Paroxetine to treat irritable bowel syndrome not responding to high-fiber diet: a double-blind, placebo-controlled trial. *Am J Gastroenterol* 2004;99:914-920.
 162. Tack J, Broekaert D, Fischler B, Van Oudenhove L, Gevers AM, Janssens J. A controlled crossover study of the selective serotonin reuptake inhibitor citalopram in irritable bowel syndrome. *Gut* 2006;55:1095-1103.
 163. Corazzari E. Role of opioid ligands in the irritable bowel syndrome. *Can J Gastroenterol* 1999;13(suppl A):71A-75A.
 164. Fraitag B, Homerin M, Hecketsweiler P. Double-blind dose-response multicenter comparison of fedotozine and placebo in treatment of nonulcer dyspepsia. *Dig Dis Sci* 1994;39:1072-1077.
 165. Mangel A, Bornstein J, Hamm L, et al. Clinical trial: asimadoline in the treatment of patients with irritable bowel syndrome. *Aliment Pharmacol Ther* 2008;28:239-249.
 166. Hawkes ND, Rhodes J, Evans BK, Rhodes P, Hawthorne AB, Thomas GA. Naloxone treatment for irritable bowel syndrome - a randomized controlled trial with an oral formulation. *Aliment Pharmacol Ther* 2002;16:1649-1654.
 167. Camilleri M. Current and future pharmacological treatments for diarrhea-predominant irritable bowel syndrome. *Expert Opin Pharmacother* 2013;14:1151-1160.
 168. Simrén M. Bloating and abdominal distention: not so poorly understood anymore! *Gastroenterology* 2009;136:1487-1490.
 169. Chey WD, Paré P, Viegas A, Ligozio G, Shetzline MA. Tegaserod for female patients suffering from IBS with mixed bowel habits or constipation: a randomized controlled trial. *Am J Gastroenterol* 2008;103:1217-1225.
 170. George AM, Meyers NL, Hickling RI. Clinical trial: renzapride therapy for constipation-predominant irritable bowel syndrome - multicentre, randomized, placebo-controlled, double-blind study in primary healthcare setting. *Aliment Pharmacol Ther* 2008;27:830-837.
 171. Dobrilla G, Imbimbo BP, Piazza L, Bensi G. Longterm treatment of irritable bowel syndrome with cimetropium bromide: a double blind placebo controlled clinical trial. *Gut* 1990;31:355-358.
 172. Mitchell SA, Mee AS, Smith GD, Palmer KR, Chapman RW. Alverine citrate fails to relieve the symptoms of irritable bowel syndrome: results of a double-blind, randomized, placebo-controlled trial. *Aliment Pharmacol Ther* 2002;16:1187-1195.
 173. Chang FY, Lu CL, Luo JC, Chen TS, Chen MJ, Chang HJ. The evaluation of otilonium bromide treatment in asian patients with irritable bowel syndrome. *J Neurogastroenterol Motil* 2011;17:402-410.
 174. Wittmann T, Paradowski L, Ducrotté P, Bueno L, Andro Delestrain MC. Clinical trial: the efficacy of alverine citrate/simethicone combination on abdominal pain/discomfort in irritable bowel syndrome - a randomized, double-blind, placebo-controlled study. *Aliment Pharmacol Ther* 2010;31:615-624.
 175. Vahedi H, Merat S, Rashidioon A, Ghoddoosi A, Malekzadeh R. The effect of fluoxetine in patients with pain and constipation-predominant irritable bowel syndrome: a double-blind randomized-controlled study. *Aliment Pharmacol Ther* 2005;22:381-385.
 176. Vahedi H, Merat S, Momtahan S, et al. Clinical trial: the effect of amitriptyline in patients with diarrhoea-predominant irritable bowel syndrome. *Aliment Pharmacol Ther* 2008;27:678-684.
 177. Read NW, Abitbol JL, Bardhan KD, Whorwell PJ, Fraitag B. Efficacy and safety of the peripheral kappa agonist fedotozine versus placebo in the treatment of functional dyspepsia. *Gut* 1997;41:664-668.
 178. Szarka LA, Camilleri M, Burton D, et al. Efficacy of on-demand asimadoline, a peripheral kappa-opioid agonist, in females with irritable bowel syndrome. *Clin Gastroenterol Hepatol* 2007;5:1268-1275.