Heme Oxygenase-1 via Transcriptional Activation of Nrf2 Attenuates Apoptosis in Lung Cancer Cells

Purpose: Increasing evidences have indicated the critical role of HO-1 in cytoprotection and more diverse biological functions. HO-1 has been reported to stimulate tumor cell growth and proliferation. And several tumors, including renal cell carcinoma, prostate tumor, hepatoma, and sarcoma, express a high level of HO-1. Indeed, inhibition of HO-1 by using specific HO inhibitors demonstrated *in vivo* antitumor activity. However, the precise mechanism of HO-1 induction and signals in lung cancer is not clearly known yet. We aimed to analyze the role of HO-1, characterize the mechanism of HO-1 induction, the role of Nrf2 in the induction, and investigated whether inhibition of HO-1 may induce apoptosis in lung cancer cells. **Materials and Methods:** Western blot and immunostaining analyses were performed to ascertain whether HO-1, Nrf2, and NFkB were expressed or not in various lung cancer cell lines. Apoptosis by HO-1 inhibition through siRNA transfection was measured by flow cytometric analysis and Western blot. And the expression of HO-1 by siRNA of Nrf2 and NFkB was examined by ARE-driven luciferase activity and Western blot. **Results:** We demonstrated that HO-1 was expressed highly in A549 cells than other lung cancer cells. And A549 cells were transfected by HO-1 small interfering RNA (siRNA) induced apoptosis. Nrf2 siRNA, next, resulted in a decrease of HO-1 expression. However, NFkB siRNA had no influence on the expression of HO-1. **Conclusion:** Increasing HO-1 expression in A549 cells may be resulted from the transcriptional activation of Nrf2, and inhibition of Nrf2-HO-1 pathway induces apoptosis. Therefore it provides new important insights into the possible molecular mechanism of the antitumor therapy. (J Lung Cancer 2006;5(1):39 – 46)

Key Words: Heme oxygenase-1, Nrf2, Apoptosis, Lung cancer

Hak-Rvul Kim, M.D.¹
Eun-Jung Kim, M.S.¹
Ki-Eun Hwang, M.D.¹
So-Young Kim, M.D.¹
Seong-Hoon Park, M.D.²
Jung-Hyun Park, M.D.¹
Hwi-Jung Kim, M.D.¹
Sei-Hoon Yang, M.D.¹ and
Eun-Taik Jeong, M.D.¹

Departments of ¹Internal Medicine
and ²Radiology, Wonkwang University
School of Medicine, Iksan, Korea

Received: May 15, 2006
Accepted: May 30, 2006

Address for correspondence
Eun-Taik Jeong, M.D.
Department of Internal Medicine, Wonkwang University School of Medicine,
344-2, Shinyong-dong, Iksan 570-749,
Korea
Tel: 82-63-850-1073
Fax: 82-63-855-2025
E-mail: jetpul@wonkwang.ac.kr

INTRODUCTION

Heme oxygenase (HO) is the rate-limiting enzyme in heme catabolism, which leads to the generation of biliverdin, free iron and carbon monoxide (CO)(1). Biliverdin is subsequently converted to bilirubin, potent antioxidant, by cytosolic biliverdin reductase, and free iron is promptly sequestered into ferritin(2). Three distinct isoforms of HO have been identified and cloned(3). These isozymes are products of different genes and differ markedly in their tissue distribution.

HO-1, an inducible isoform of HO, is a member of the family of heat shock proteins (HSP32). It is found at low levels in most mammalian tissues but is highly expressed in liver and

spleen(4), and its expression is triggered by diverse stress-inducing stimuli including hypoxia(5), heat shock(6), UV irradiation(7), heavy metals(8,9), reactive oxygen species (ROS) such as hydrogen peroxide (H₂O₂)(10,11), and reactive nitrogen oxide such as nitric oxide (NO)(12). The expression of HO-1 gene is primarily regulated at the transcriptional level although there are interspecies variations in the regulation of HO-1(13~15). The transcriptional factor Nrf2 plays a key role in HO-1 gene activation(16). The biological functions of HO-1 are thus believed to be associated with a fundamental adaptive and defense response against oxidative stress and cellular stress. Recent findings indicate that HO-1 possess anti-inflammatory and antiapoptotic functions, and it exert a role in controlling growth

and cell proliferation in a cell-specific manner(17~20). Elevated HO-1 expression and activity was found in various tumors such as renal cell carcinoma, prostate tumors, hepatoma, and lymphosarcomas. Indeed, inhibition of HO-1 by using specific HO inhibitors such as zinc protoporphyrin (ZnPP) demonstrated *in vivo* antitumor activity(7,21~23). We, therefore, hypothesized that inhibition of HO-1 expression and activity may sensitize lung cancer cells to anticancer treatment modalities. In this study, we aimed to analyze the role of HO-1, characterize the mechanism of HO-1 induction, the role of Nrf2 in the induction, and investigated whether inhibition of HO-1 may induce apoptosis in lung cancer cells.

MATERIALS AND METHODS

1) Cell culture

A549, NCI-H23, NCI-H157, and NCI-H460 human lung cancer cells were obtained from Korean Cell Line Bank (Seoul, Korea) and grown in RPMI 1,640 containing 100 units/ml penicillin, 0.1 mg/ml streptomycin, and 10% FBS, and maintained in a humidified atmosphere of 5% CO₂ in air at 37°C and maintained in log phase.

2) Western blotting

Cell extracts were separated by 10% SDS-PAGE under reduced conditions and transferred onto nitrocellulose membrane. The membrane was blocked with 5% skim milk in TBS-T (25 mM Tris, pH 7.6, 138 mM NaCl and 0.05% Tween-20) for 1 hour and probed with primary antibodies (1 : 1,000~1 : 5,000). After a series of washes, the membrane was further incubated with secondary antibody (1 : 2,000~1 : 10,000) conjugated with horse radish peroxidase (HRP). The immunoreactive signal was detected with enhanced chemiluminescent (ECL) detection system (Amersham Co, England).

3) Immunocytochemistry

After removal of the culture medium, untreated lung cancer cells were washed three times with PBS, and fixed with 4% paraformaldehyde in PBS for 1 hour at room temperature (RT). Cells were then rinsed twice with PBS, preincubated in the same buffer containing 0.1% Triton X-100, and followed by three washes. Cells were blocked with 0.1% BSA in PBS for 1 hour and then respectively incubated with primary anti-HO-1,

anti-Nrf2, and anti-NFkB (Santa Cruz Biotech Inc, Santa Cruz, CA) in PBS containing 3% BSA for 18~20 hour at 4°C. After three washes with PBS, cells were incubated with goat anti-rabbit Texas Red-tagged secondary antibody at a dilution of 1 : 500 in PBS for 1 hour at RT. Cells were then washed in PBS and further incubated with 10 µM Hoechst 33,342 in PBS at RT for 30 min. Subcellular localization of HO-1, Nrf2, and NFkB were observed under fluorescent microscope equipped with digital camera (IX71, Olympus, Japan). The fluorescent images were captured using appropriate filters.

4) Flow cytometric analysis

The degree of apoptosis was determined by measuring the number of cells showing below the G₀/G₁ phase of DNA content from Flow cytometric analysis after staining of cells with propidium iodide (PI) as originally described by Crissman and Steinkamp(24). Cell cycle analysis was performed with FACScan equipped with a Cell Quest software (Becton Dickinson, San Jose, CA).

5) Preparation of cytosolic and nuclear extracts

Cells were washed with ice-cold PBS, scraped, and centrifuged at 1,000×g for 5 min at 4°C. The cell pellet was resuspended in 200 µl of lysis buffer (10 mM HEPES, pH 7.9, 1.5 mM MgCl₂, 10 mM KCl, 0.5 mM phenylmethylsulfonyl fluoride, and 0.5 mM dithiothreitol) and incubated on ice for 15 min. At the end of this incubation, 10 µl of 10% NP-40 was added and the tube was vortexed for 10 s. After centrifugation at 13,000×g for 1 min at 4°C, supernatants (cytosolic extracts) were collected and stored at 80°C, whereas the pellets were further processed to obtain nuclear extracts. The pellet was resuspended in extraction buffer (5 mM HEPES, pH 7.9, 1.5 mM MgCl₂, 0.5 mM phenylmethylsulfonyl fluoride, 0.2 mM EDTA, 0.5 mM dithiothreitol, and glycerol 25% vol/vol) and incubated for 30 min at 4°C. Nuclear extracts were isolated by centrifugation at 13,000×g for 30 min at 4°C. The supernatant was aliquoted and stored at 80°C until used for western blot analysis. Protein concentration was determined by the Lowry method.

6) Transfection of HO-1, Nrf2, and NFkB small interfering RNA (siRNA) into A549 cells

To study the effects of HO-1, Nrf2, and NFkB inhibition,

their expressions were specifically suppressed by introduction of 21-nucleotide duplex small interfering RNA (siRNA), which targets nucleotides of the HO-1, Nrf2, and NFkB mRNA coding sequence(25). Briefly, A549 cells (10^5 cells/well) were plated in 6-well plates and transiently transfected with 2 μ g per well of HO-1, Nrf2, and NFkB siRNA (Santa Cruz Biotechnology, Santa Cruz, CA) mixed with X-tremeGENE siRNA transfection reagent (Roche Applied Science, Penzberg, Germany) according to the manufacturer's directions. The Silencer Negative Control siRNA (Roche Applied Science, Penzberg, Germany) was used as a negative control and was introduced into the cells using the same protocol.

7) Reporter gene activity assay

Cells were transfected with ARE-luciferase (ARE-Luc) reporter plasmid. ARE-Luc was generated by transferring the enhancer 2 (E2) and minimal promoter (MP) sequence 40 into luciferase reporter plasmid pGL3-Basic. E2 and MP were generated by PCR of mouse cDNA with oligonucleotide pairs E2 SacI: 5'-AGC TGA GCT CCT TAA GCA ATC CAT TAG

GAA TCC G-3' and E2 ApaI: 5'-AGC TGG GCC CCC GCT CTC TTG CCA GAC TC-3' and MP ApaI: 5'-TTA CAG GCA GGG CCC GCC T-3' and MP XhoI: 5'-TAT GCT CGA GAC GGC TCT GC-3', respectively. For transfection of reporter plasmid, cells were seeded on 24-well plates at a density of 3×10^4 cells/well on the day before transfection. A total of 750 ng of plasmid DNA including 200 ng of luciferase reporter, Nrf2 and NFkB siRNA, and 50 ng of pcDNA3- β -gal was transfected into cells by TfxTM-50 reagent (Promega, Madison, WI). The amount of transfected DNA was held constant to 750 ng by the addition of empty vector DNA where necessary. After 48 h of transfection, cells were washed twice with PBS buffer and then lysed in reporter lysis buffer. Luciferase activity was measured with a luciferase assay system (Promega) according to the manufacturer's instructions. Luciferase activity was measured in triplicate, averaged, and then normalized with β -galactosidase activity using the galactosidase assay system (Galacto-Light, Tropix Inc., MA) according to the manufacturer's instructions.

Fig. 1. Expression of HO-1 in endogenous lung cancer cells. (A) Expression levels of HO-1 protein were measured by Western blotting using anti-HO-1 antibody. Lysates of untreated lung cancer cells were subjected to 12.5% SDS-PAGE to measure the expression of HO-1 protein. (B) Untreated lung cancer cells were reacted with Hoechst and anti-HO-1 antibody. Then, cells were visualized under fluorescent microscope.

8) Statistical analysis

Each experiment was performed at least three times, and all values are represented as means $\pm$ SD of triplicates. Student's t-test was used to analyze a statistical significance of the results. Values of $p < 0.05$ were considered as statistically significant.

RESULTS

1) HO-1 expression in endogenous lung cancer cells

To study of the role of HO-1 on apoptosis, Western blot and immunostaining analyses were performed to ascertain whether HO-1 was expressed or not in various lung cancer cell lines. Untreated lung cancer cells were used to carry out Western blot for HO-1 (Fig. 1A). The data revealed that HO-1 protein was expressed highly in A549 cells than other lung cancer cell lines. Consistent with proteins level of HO-1 expressed in Fig. 1A, immunostaining analyses demonstrated that HO-1 was strongly located in cytoplasm of A549 cells, whereas a little proportion of other lung cancer cells showed positive staining (Fig. 1B).

2) Induction of apoptosis by HO-1 siRNA transfection in A549 cells

To further elucidate the functional role of HO-1 inhibition, we suppressed HO-1 mRNA through siRNA transfection of a 21-nucleotide duplex HO-1 siRNA. The Silencer Negative Control siRNA 1 was used as a negative control. Suppression of HO-1 levels after siRNA treatment for 48 hours was confirmed by RT-PCR analysis for HO-1 mRNA of A549 cells. siRNA transfection resulted in targeted knockdown of HO-1 mRNA expression (data not shown). We, next, examined whether HO-1 siRNA may induce apoptosis in A549 cells. DNA fragmentation by apoptosis phenomenon was analyzed as the sub-G₀/G₁ fraction of cell cycle analysis by flow cytometry in A549 cells stained with PI; contrast to control and nonsense siRNA groups, the sub-G₀/G₁ fraction after HO-1 siRNA groups significantly increased up to 22.4% (Fig. 2A). To gain the down-stream event of apoptosis, we further tested the activation of caspase-3 by Western blot analysis. As shown in Fig. 2B, degradation of pro-caspase-3 occurred in HO-1 siRNA groups.

Fig. 2. Apoptosis in A549 cells after HO-1 siRNA transfection. Non-sense indicates the result for cells treated with Silencer Negative Control siRNA. Cells were transfected with 0.5 μ g siRNA of HO-1 and control plasmid. (A) Cellular DNA was stained with PI and cell cycle was analyzed by flow cytometry. The data represent the mean $\pm$ S.D. of triplicate. * $p < 0.05$. (B) Cells were transfected with 0.5 μ g siRNA of HO-1 and control plasmid. After 48 hrs, lysate was subjected to 12.5% SDS-PAGE to measure the expression of pro-caspase-3.

3) Expression of Nrf2 and NFkB in endogenous lung cancer cells

Many studies have reported that HO-1 expression is mainly regulated by NFkB, p38 MAP kinase signaling pathways, and Nrf2-dependent antioxidant response element (ARE) activation in several cell types(26~28). Therefore, to further provide the exact mechanism of up-regulation of HO-1 gene expression in A549 cells, Western blot and immunostaining analyses were performed to ascertain whether Nrf2 and NFkB were expressed or not in lung cancer cell lines. Untreated lung cancer cells were used to carry out Western blot in both cytosolic and nuclear fractions for Nrf2 and NFkB (Fig. 3A). The data demonstrated that highly expressed Nrf2 and NFkB in nucleus and degraded IkB in cytoplasm presented in A549 cells than other lung cancer cell lines. Consistent with proteins level expressed in Fig. 3A, immunostaining analyses demonstrated that the many proportion of Nrf2 and NFkB translocated into the nucleus in endogenous A549 cells (Fig. 3B).

Fig. 3. Expression of Nrf2 and NFkB in endogenous lung cancer cells. (A) Expression levels in both cytosolic and nuclear fractions of Nrf2 and NFkB protein were measured by Western blotting. Lysate was fractionated into cytosolic and nuclear portions and proteins were separated on 15% SDS-PAGE to immunoblot for Nrf2 and NFkB in nucleus and IκB in cytoplasm. (B) Untreated A549 cells were reacted with Hoechst and anti-HO-1 antibody. Then, cells were visualized under fluorescent microscope.

4) Effect of Nrf2 and NFkB siRNA transfection on HO-1 promotor activity

To confirm the involvement of Nrf2 and NFkB in pathways of up-regulation of HO-1 gene expression, siRNA of Nrf2 and NFkB were transfected in A549 cells. Suppression of Nrf2 and NFkB levels after siRNA treatment for 48 hours was respectively confirmed by RT-PCR analysis for Nrf2 and NFkB mRNA of A549 cells (data not shown). And we measured indirectly the expression level of HO-1 through using the ARE-driven luciferase activity. As shown in Fig. 4, knock down the transcriptional expression of Nrf2 by using siRNA significantly suppressed the transcriptional activation of ARE up to 11.7-fold of controls ($p < 0.05$). However, NFkB siRNA only partial involved the inhibition of HO-1 signalling. These results suggest that up-regulation of HO-1 was obligatory required in the pathway of Nrf2/HO-1.

5) Effect of Nrf2 and NFkB siRNA transfection on HO-1 expression

We further examined that transfection of siRNAs apparently

Fig. 4. Expression of Nrf2 and NFkB siRNA transfection on HO-1 promotor activity. In A549 cells, Nrf2 and NFkB expression vectors were respectively cotransfected with ARE-luciferase reporter gene for 24 hrs, and then luciferase activity was measured at 18 h later. The data represent the mean $\pm$ S.D. of triplicate. * $p < 0.05$.

knocked down the expression of HO-1 gene at the translational level. siRNA of Nrf2 and NFkB were transfected in A549 cells, and suppression of Nrf2 and NFkB levels after siRNA treatment for 48 hours was respectively confirmed by Western blot. Consistent with previous results of transcriptional level, the

Fig. 5. Expression of Nrf2 and NFkB siRNA transfection on HO-1 protein expression. Nonsense indicates the result for cells treated with Silencer Negative Control siRNA. (A) Cells were transfected with 0.5 μ g siRNA of Nrf2 and control plasmid. Suppression of Nrf2 levels after siRNA treatment after 48 hrs was respectively confirmed by Western blot. Simultaneously lysates were subjected to 12.5% SDS-PAGE to measure the expression of HO-1 protein. (B) Cells were transfected with 0.5 μ g siRNA of NFkB and control plasmid. Suppression of NFkB levels after siRNA treatment after 48 hrs was respectively confirmed by Western blot. Simultaneously lysates were subjected to 12.5% SDS-PAGE to measure the expression of HO-1 protein.

expression of HO-1 protein was completely disappeared by transfection of Nrf2 siRNA (Fig. 5A). However, NFkB siRNA did not affect the total amount of HO-1 protein expression level (Fig. 5B). These data indicate that the expression of HO-1 is indeed regulated through Nrf2-driven transcription activation, whereas it is independent of NFkB signaling pathway.

DISCUSSION

HO-1 has been reported to stimulate cell growth and the proliferation of various types of cells. The transfection of HO-1 gene into coronary endothelial cells promoted the formation of capillary-like tubular structures(29), high expression of HO-1 was observed in the hyperproliferating epidermis during cutaneous wound repair and in psoriatic skin lesions(30). HO-1 was also extensively expressed in various tumor cells, which are the most rapidly proliferating cells, and different human cancers express high levels of HO-1, which may provide them a growth advantage(31). Therefore, inhibition of HO-1 might lead to reduced tumor growth *in vitro* and *in vivo*(7,32,33). In our la-

boratory, HO-1 expression is significantly up-regulated in A549 cells than other lung cancer cell lines by Western blot and immunostaining analyses.

Because tumor growth is based on augmented cell growth and prolonged cell survival, it is possible that the strong HO-1 positivity in A549 cells develops its tumor-promoting activity by increasing cell proliferation and protection towards apoptosis. It is now well known that HO-1 has an important role in rapid tumor growth because of its antioxidative and anti-apoptotic effects(7,34). Fang et al demonstrated that transfection of SW480 cells with siRNA for HO-1, to reduce HO-1 expression, resulted in a similar induction of apoptosis(32). In this study, we showed that A549 cells transfected by HO-1 siRNA induced apoptosis, demonstrating increased sub-G₀/G₁ fraction and activation of caspase-3.

Nrf2 is a basic leucine zipper transcription factor, which transcriptionally regulates many genes including HO-1(35), NAD(P)H:quinine oxidoreductase-1 (NQO-1), γ -glutamylcysteine synthase, and glutathione S-transferase. Inactive Nrf2 is localized in the cytoplasm, in part or totally, as a consequence of binding to cytoskeleton-associated protein Keap1(36). Increased Nrf2 in the nucleus enhances its binding activity to ARE, followed by an increase of HO-1 gene transcription and protein synthesis. NFkB, an antiapoptotic transcription factor, is located in the cytosol as a three-subunit complex. When suitably stimulated, the inhibitory subunit I κ B will dissociate from the other two prototypical subunits so that it enables the p50-p65 dimer to translocate to the nucleus, where it activate relevant genes (37). Liu et al reported that the expression of HO-1 by HO-1 stimuli was reduced to the control level when human gastric cancer cells were treated with dominant-negative NFkB inhibitor, NFkB, thus, plays a critical role in the HO-1(38). In this study, we showed that highly expressed Nrf2 and NFkB in nucleus and degraded I κ B in cytoplasm presented in A549 cells. Immunostaining analyses also demonstrated that the many proportion of Nrf2 and NFkB translocated into the nucleus in endogenous A549 cells.

To further provide the exact mechanism how the expression of HO-1 was increased in A549 cells, we examined two signaling pathways, including Nrf2 and NFkB. Our data demonstrated that Nrf2 siRNA could significantly suppress the Nrf2-driven transcriptional activation of ARE-luciferase, but NFkB siRNA was not prominent compared to cells transfected Nrf2

siRNA. Our results by using the Western blot also suggested that Nrf2/HO-1 signaling was directly involved the increased expression of HO-1 in A549 cells. On the contrary, Nrf2 siRNA had no influence on the expression of HO-1.

In conclusion, we have demonstrated that HO-1 was expressed highly in A549 cells, and knock down of HO-1 mRNA by using the siRNA induced apoptosis. We have further elucidated that Nrf2/HO-1 signaling is directly involved in the mechanism of HO-1 expression in A549 cells. Taken together, these results suggest that high HO-1 levels via transcriptional activation of Nrf2 in A549 lung cancer cells, at least partly, be responsible for their resistance, which may be a potential new therapeutic targets.

ACKNOWLEDGEMENT

This study was supported by a grant from Wonkwang University in 2006.

REFERENCES

1. Tenhunen R, Marver HS, Schmid R. The enzymatic catabolism of hemoglobin: stimulation of microsomal heme oxygenase by hemin. *J Lab Clin Med* 1970;75:410-421.
2. Baranano DE, Rao M, Ferris CD, Snyder SH. Biliverdin reductase: a major physiologic cytoprotectant. *Proc Natl Acad Sci USA* 2002;99:16093-16098.
3. Maines MD. The heme oxygenase system: a regulator of second messenger gases. *Annu Rev Pharmacol Toxicol* 1997;37:517-554.
4. Tenhunen R, Marver HS, Schmid R. The enzymatic catabolism of hemoglobin: stimulation of microsomal heme oxygenase by hemin. *J Lab Clin Med* 1970;75:410-421.
5. Motterlini R, Foresti R, Bassi R, Calabrese V, Clark JE, Green CJ. Endothelial heme oxygenase-1 induction by hypoxia. Modulation by inducible nitric-oxide synthase and S-nitrosothiols. *J Biol Chem* 2000;275:13613-13620.
6. Stuhlmeier KM. Activation and regulation of Hsp32 and Hsp70. *Eur J Biochem* 2000;267:1161-1167.
7. Doi K, Akaike T, Fujii S, et al. Induction of haem oxygenase-1 nitric oxide and ischaemia in experimental solid tumours and implications for tumour growth. *Br J Cancer* 1999;80:1945-1954.
8. Elbirt KK, Whitmarsh AJ, Davis RJ, Bonkovsky HL. Mechanism of sodium arsenite-mediated induction of heme oxygenase-1 in hepatoma cells. Role of mitogen-activated protein kinases. *J Biol Chem* 1998;273:8922-8931.
9. Eyssen-Hernandez R, Ladoux A, Frelin C. Differential regulation of cardiac heme oxygenase-1 and vascular endothelial growth factor mRNA expressions by hemin, heavy metals, heat shock and anoxia. *FEBS Lett* 1996;382:229-233.
10. Keyse SM, Tyrrell RM. Heme oxygenase is the major 32-kDa stress protein induced in human skin fibroblasts by UVA radiation, hydrogen peroxide, and sodium arsenite. *Proc Natl Acad Sci USA* 1989;86:99-103.
11. Lautier D, Luscher P, Tyrrell RM. Endogenous glutathione levels modulate both constitutive and UVA radiation/hydrogen peroxide inducible expression of the human heme oxygenase gene. *Carcinogenesis* 1992;13:227-232.
12. Hara E, Takahashi K, Takeda K, et al. Induction of heme oxygenase-1 as a response in sensing the signals evoked by distinct nitric oxide donors. *Biochem Pharmacol* 1999;58:227-236.
13. Rizzardini M, Terao M, Falciani F, Cantoni L. Cytokine induction of haem oxygenase mRNA in mouse liver. Interleukin 1 transcriptionally activates the haem oxygenase gene. *Biochem J* 1993;290:343-347.
14. Agarwal A, Shiraishi F, Visner GA, Nick HS. Linoleyl hydroperoxide transcriptionally upregulates heme oxygenase-1 gene expression in human renal epithelial and aortic endothelial cells. *J Am Soc Nephrol* 1998;9:1990-1997.
15. Fogg S, Agarwal A, Nick HS, Visner GA. Iron regulates hyperoxia-dependent human heme oxygenase 1 gene expression in pulmonary endothelial cells. *Am J Respir Cell Mol Biol* 1999;20:797-804.
16. Alam J, Stewart D, Touchard C, Boinapally S, Choi AM, Cook JL. Nrf2, a Cap'n'Collar transcription factor, regulates induction of the heme oxygenase-1 gene. *J Bio Chem* 1999;274:26071-26078.
17. Wagner M, Cadetg P, Ruf R, Mazzucchelli L, Ferrari P, Redaelli CA. Heme oxygenase-1 attenuates ischemia/reperfusion-induced apoptosis and improves survival in rat renal allografts. *Kidney Int* 2003;63:1564-1573.
18. Choi BM, Pae HO, Chung HT. Nitric oxide priming protects nitric oxide-mediated apoptosis via heme oxygenase-1 induction. *Free Radic Biol Med* 2003;34:1136-1145.
19. Amon M, Menger MD, Vollmar B. Heme oxygenase and nitric oxide synthase mediate cooling-associated protection against TNF-alpha-induced microcirculatory dysfunction and apoptotic cell death. *F ASEB J* 2003;17:175-185.
20. Ke B, Shen XD, Zhai Y, et al. Heme oxygenase 1 mediates the immunomodulatory and antiapoptotic effects of interleukin 13 gene therapy in vivo and in vitro. *Hum Gene Ther* 2002;13:1845-1857.
21. Goodman AI, Choudhury M, da Silva JL, Schwartzman ML, Abraham NG. Overexpression of the heme oxygenase gene in renal cell carcinoma. *Proc Soc Exp Biol Med* 1997;214:54-61.
22. Maines MD, Abrahamsson PA. Expression of heme oxygenase-1 (HSP32) in human prostate: normal, hyperplastic, and tumor tissue distribution. *Urology* 1996;47:727-733.
23. Sahoo SK, Sawa T, Fang J, et al. Pegylated zinc protoporphyrin: a water-soluble heme oxygenase inhibitor with tumor-targeting capacity. *Bioconjug Chem* 2002;13:1031-1038.
24. Crissman HA, Steinkamp JA. Cell cycle-related changes in

- chromatin structure detected by flow cytometry using multiple DNA fluorochromes. *Eur J Histochem* 1993;37:129-138.
25. Elbashir SM, Harborth J, Lendeckel W, Yalcin A, Weber K, Tuschl T. Duplexes of 21-nucleotide RNAs mediate RNA interference in cultured mammalian cells. *Nature* 2001;411:494-498.
26. Alam J, Wicks C, Stewart D, et al. Mechanism of heme oxygenase-1 gene activation by cadmium in MCF-7 mammary epithelial cells. Role of p38 kinase and Nrf2 transcription factor. *J Biol Chem* 2000;275:27694-27702.
27. He CH, Gong P, Hu B, et al. Identification of activating transcription factor 4 (ATF4) as an Nrf2-interacting protein. Implication for heme oxygenase-1 gene regulation. *J Biol Chem* 2001;276:20858-20865.
28. Banning A, Brigelius-Flohe R. NF-kappaB, Nrf2, and HO-1 interplay in redox regulated VCAM-1 expression. *Antioxid Redox Signal* 2005;7:889-899.
29. Deramaudt BM, Braunstein S, Remy P, Abraham NG. Gene transfer of human heme oxygenase into coronary endothelial cells potentially promotes angiogenesis. *J Cell Biochem* 1998;68:121-127.
30. Hanselmann C, Mauch C, Werner S. Heme oxygenase-1: a novel player in cutaneous wound repair and psoriasis? *Biochem J* 2001;353:459-466.
31. Fang J, Akaike T, Maeda H. Antiapoptotic role of heme oxygenase (HO) and the potential of HO as a target in anticancer treatment. *Apoptosis* 2004;9:27-35.
32. Fang J, Sawa T, Akaike T, et al. *In vivo* antitumor activity of pegylated zinc protoporphyrin: targeted inhibition of heme oxygenase in solid tumor. *Cancer Res* 2003;63:3567-3574.
33. Sunamura M, Duda DG, Ghattas MH, et al. Heme oxygenase-1 accelerates tumor angiogenesis of human pancreatic cancer. *Angiogenesis* 2003;6:15-24.
34. Tanaka S, Akaike T, Fang J, et al. Antiapoptotic effect of haem oxygenase-1 induced by nitric oxide in experimental solid tumor. *Br J Cancer* 2003;88:902-909.
35. Alam J, Stewart D, Touchard C, Boinapally S, Choi AM, Cook JL. Nrf2, a Cap'n'Collar transcription factor, regulates induction of the heme oxygenase-1 gene. *J Biol Chem* 1999;274:26071-26078.
36. Zipper LM, Mulcahy RT. The Keap1 BTB/POZ dimerization function is required to sequester Nrf2 in cytoplasm. *J Biol Chem* 2002;277:36544-36552.
37. Barkett M, Gilmore TD. Control of apoptosis by Rel/NF-kappa B transcription factors. *Oncogene* 1999;18:6910-6924.
38. Liu ZM, Chen GG, Ng EK, Leung WK, Sung JJ, Chung SC. Upregulation of heme oxygenase-1 and p21 confers resistance to apoptosis in human gastric cancer cells. *Oncogene* 2004;23:503-513.