



The Role of Circulating Slit2, the One of the Newly Batokines, in Human Diabetes Mellitus

Yea Eun Kang¹, Sorim Choung², Ju Hee Lee¹, Hyun Jin Kim¹, Bon Jeong Ku^{1,2}

Departments of ¹Endocrinology and Metabolism, ²Medical Science, Chungnam National University School of Medicine, Daejeon, Korea

Background: Slit2 is a new secreted protein from adipose tissue that improves glucose hemostasis in mice; however, there is no study about the serum levels and precise role of Slit2 in human. The aim of this study is to explore the serum level of Slit2 in human, and to identify the role of Slit2 in diabetes mellitus (DM).

Methods: The participants of this study consist of 38 subjects with newly diagnosed DM, and 75 healthy subjects as a control group. Serum Slit2 levels were measured using an enzyme-linked immunosorbent assay. Relationship between circulating Slit2 and diabetic related factors was investigated in diabetic group compared with non-diabetic group. Additionally, the correlations between the serum level of Slit2 and diverse metabolic parameters were analyzed.

Results: Circulating Slit2 level was more decreased in diabetic group than in control group, but there was no significant difference statistically. Interestingly, serum levels of Slit2 were significantly negatively correlated to the serum concentrations of fasting glucose (coefficient $r=-0.246$, $P=0.008$), the serum concentrations of postprandial glucose (coefficient $r=-0.233$, $P=0.017$), and glycosylated hemoglobin (HbA1c; coefficient $r=-0.357$, $P<0.001$).

Conclusion: From our study, the first report of circulating Slit2 levels in human, circulating Slit2 level significantly negatively correlated with serum glucose and HbA1c. Our results suggest that the circulating Slit2 may play a role in maintenance of glucose homeostasis in human, even though exact contribution and mechanism are not yet known.

Keywords: Slit2; Batokine; Adipose tissue, brown; Adipokines; Diabetes mellitus

INTRODUCTION

Recent experimental studies have revealed the association of brown adipose tissue (BAT) activity with prevention against metabolic disease, such as type 2 diabetes mellitus (T2DM), obesity and dyslipidemia [1]. BAT is traditionally known for consuming energy in response to cold [2], but also has a secretory role, which contribute to protection against chronic inflammation for thermogenesis, and the BAT-derived endocrine regu-

latory factors called batokines [1,3]. Slit2, a newly discovered batokines, was identified as a member of family of Slit homolog proteins, which are produced endogenously by beige fat cells under the control of cold exposure and PR domain containing 16 (PRDM16) [4]. Slits are known to have the role in axon guidance [5,6], and recently Svensson et al. [4] described the role of Slit2 in regulating glucose homeostasis in adipose tissue. C-terminal fragments of Slit2 are the active fragments in the regulation of beige adipocyte induction [4]. Additionally, increased

Received: 19 June 2017, Revised: 25 July 2017, Accepted: 4 August 2017

Corresponding author: Bon Jeong Ku

Department of Internal Medicine, Chungnam National University School of Medicine, 282 Munhwa-ro, Jung-gu, Daejeon 35015, Korea

Tel: +82-42-280-7148, Fax: +82-42-280-7995, E-mail: bonjeong@cnu.ac.kr

Copyright © 2017 Korean Endocrine Society

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

circulating Slit2-C can promote thermogenesis of adipose tissue, increase energy expenditure, and improve glucose homeostasis in obese mice [4]. Although expression of Slit2 protein in adipose tissue may be linked to activate a thermogenic protein kinase A pathway *in vivo* [4], the role of Slit2 in human is not yet fully understood, and circulating Slit2 has not been studied. For these reasons, the aim of our study was to investigate the relationship between circulating Slit2 levels and several metabolic parameters in participants with newly diagnosed T2DM and control participants.

METHODS

Subjects

We studied a total of 45 participants with newly diagnosed T2DM and 78 control participants from the outpatient clinic of the Division of Endocrinology and Metabolism of Chungnam National University Hospital from January 2015 to July 2016 were studied. The inclusion criteria were as follows: body mass index (BMI) >18.0 kg/m², age >18 years, absence of any clinical sign of infection or inflammation, no alcohol or drug abuse, systolic blood pressure (SBP) less than 140 mm Hg and diastolic blood pressure (DBP) less than 100 mm Hg, no pregnancy and no previous history of diabetes mellitus. Ten participants were excluded because of liver cirrhosis (one case), malignancies (five cases of thyroid cancer, one case of prostate cancer, and one case of pituitary adenoma), severe rheumatoid arthritis (one case), and low BMI 16.2 kg/m² (one case). Finally, 113 participants were assessed to diabetes mellitus status [7]. Our experimental protocol was performed in accordance with the Declaration of Helsinki [8].

Clinical parameters

On the day on which the study commenced, all participants underwent physical examinations. Height, body weight, waist circumference (WC), SBP, and DBP were recorded [9]. Blood pressure were measured after 10 minutes of rest from each patient's right arm in the seating position by using an Omron IntelliSense Automatic Blood Pressure Monitor (Omron, Kyoto, Japan) height and body weight were measured without shoes in the morning. BMI was calculated as weight in kilograms (kg) divided by height in meters squared (m²), and WC was measured midway between the superior iliac spine at the end of gentle expiration and the lower rib margin in a standing position.

Biochemical parameters measurements

All blood samples were collected in the morning after an overnight fast of more than 8 hours into tubes containing ethylenediaminetetraacetic acid (EDTA) [9]. We measured the levels of fasting glucose, C-peptide, insulin, triglycerides, total cholesterol (TC), low density lipoprotein cholesterol (LDL-C), high density lipoprotein cholesterol (HDL-C), apolipoprotein B, aspartate aminotransferase, alanine aminotransferase, blood urea nitrogen, creatinine, high-sensitivity C-reactive protein (hs-CRP), and glycosylated hemoglobin (HbA1c). Blood chemistry and lipid profiles were measured using a blood chemistry analyzer (Hitachi 747, Hitachi, Tokyo, Japan). HbA1c was measured by high-performance liquid chromatography (Bio-Rad, Hercules, CA, USA). Insulin and C-peptide were measured by radioimmunoassay (Roche, Penzberg, Germany). We also performed the 75-g oral glucose tolerance test. The homeostasis model assessment of insulin resistance (HOMA-IR) was calculated as follows: fasting insulin level (μU/mL)×fasting glucose level (in mmol/L)/22.5 and the homeostasis model assessment of β-cell index (HOMA-β) was calculated as follows: fasting insulin level (μU/mL)×20/fastest glucose level (in mmol/L)–3.5 [8,9].

Slit2 measurements

Serum Slit2 levels were measured using a quantitative sandwich enzyme-linked immunosorbent assay (ELISA) kit (LifeSpan BioSciences Inc., Seattle, WA, USA; ELISA kit for human Slit2, catalog No. LS-F21017). The ELISA quantitatively immunoassayed human Slit2 levels. The intra-assay coefficient of variation (CV) of the ELISA kit was <10% and the inter-assay CV of the ELISA Kit was <10%.

Statistical analysis

Continuous variables are expressed as the mean±standard deviation (SD), and the significance of between-group differences was evaluated using Student *t* test or the Mann-Whitney *U* test. Categorical variables are expressed as percentages, and between-group differences were compared with the chi-square test. To evaluate associations between Slit2 level and other variables, we subjected the data to Pearson and partial correlation analyses. A two-tailed *P* value <0.05 was regarded as statistically significant. All statistical analyses were performed with SPSS version 21.0 (IBM Co., Armonk, NY, USA) [8,9].

Ethics statement

The protocol for this research was approved by the Institutional Review Board of Chungnam National University Hospital (Reg.

No. CNUH 2014-12-013).

RESULTS

Clinical and biochemical characteristics of study subjects

Baseline characteristics of the subjects are shown in Table 1. Totally, 113 participants (mean age 49.6 ± 12.2 years, 43.4% male) completed this study. The mean BMI, WC, serum glucose,

and HOMA-IR values were 24.9 ± 4.1 kg/m², 85.3 ± 11.3 cm, 125.1 ± 54.3 mg/dL, and 3.24 ± 1.99 mg/dL, respectively. The T2DM group had significantly higher levels of fasting glucose, fasting C-peptide, postprandial glucose, HOMA-IR, hemoglobin, DBP, hs-CRP, and apolipoprotein B than did those without diabetes. The T2DM group also had lower levels of HOMA- β . Circulating Slit2 level values in the diabetic group were 1.855 ± 1.306 ng/dL, and values in the control group were

Table 1. Characteristics of Study Population

Variable	T2DM group (n=38)	Non-diabetes group (n=75)	P value ^a
Sex, male/female	17/21	32/43	0.836
Age, yr	50.3 ± 10.8	49.3 ± 13.0	0.683
BMI, kg/m ²	25.7 ± 4.6	24.4 ± 3.7	0.136
Waist circumferences, cm	87.0 ± 11.6	83.5 ± 10.9	0.345
Fasting glucose, mg/dL	176.9 ± 67.7	98.9 ± 9.7	<0.001
Fasting insulin, mIU/L	11.5 ± 4.3	9.8 ± 4.8	0.055
Fasting C-peptide, mIU/L	0.88 ± 0.38	0.71 ± 0.37	0.025
Postload 2-hour glucose, mg/dL	324.5 ± 124.9	118.3 ± 30.4	<0.001
Postload 2-hour insulin, mIU/L	50.2 ± 43.8	52.1 ± 53.3	0.846
Postload 2-hour C-peptide, mIU/L	2.76 ± 1.71	3.15 ± 1.49	0.283
HbA1c, %	8.7 ± 2.4	5.4 ± 0.4	<0.001
HOMA-IR	4.8 ± 2.2	2.4 ± 1.2	<0.001
HOMA- β	24.2 ± 15.4	32.1 ± 17.0	0.016
TG, mg/dL	163.1 ± 108.4	134.1 ± 103.1	0.184
LDL-C, mg/dL	130.8 ± 42.2	120.0 ± 32.9	0.182
HDL-C, mg/dL	50.8 ± 13.0	55.6 ± 13.7	0.084
TC, mg/dL	206.0 ± 48.2	195.2 ± 36.6	0.240
AST, IU/L	25.1 ± 15.4	21.9 ± 7.5	0.147
ALT, IU/L	29.5 ± 17.4	23.9 ± 16.5	0.110
eGFR (MDRD), mL/min	112.2 ± 26.7	105.1 ± 18.4	0.108
ACR (albumin/creatinin ratio)	51.1 ± 124.7	3.98 ± 3.45	0.056
Hemoglobin, g/dL	14.6 ± 1.4	13.8 ± 1.5	0.016
SBP, mm Hg	129.0 ± 11.7	125.0 ± 13.7	0.125
DBP, mm Hg	79.3 ± 8.6	75.4 ± 8.8	0.034
hs-CRP, mg/L	4.1 ± 9.9	1.2 ± 1.8	0.034
Apolipoprotein A, mg/dL	130.6 ± 30.6	126.1 ± 25.1	0.649
Apolipoprotein B, mg/dL	120.1 ± 32.0	87.2 ± 22.2	0.003
Slit2, ng/mL	1.855 ± 1.306	2.352 ± 1.864	0.103

Values are expressed as mean \pm SD.

T2DM, type 2 diabetes mellitus; BMI, body mass index; HbA1c, glycosylated hemoglobin; HOMA-IR, homeostasis model assessment of insulin resistance; HOMA- β , homeostasis model assessment of β -cell; TG, triglycerides; LDL-C, low density lipoprotein cholesterol; HDL-C, high density lipoprotein cholesterol; TC, total cholesterol; AST, aspartate aminotransferase; ALT, alanine aminotransferase; eGFR, estimated glomerular filtration rate; MDRD, Modification of Diet in Renal Disease; ACR, albumin creatinine ratio; SBP, systolic blood pressure; DBP, diastolic blood pressure; hs-CRP, high-sensitivity C-reactive protein.

^aP values were calculated using a paired Student *t* test on the means of continuous variables and the Mann-Whitney *U* test on the means of nonparametric variables.

2.352±1.864 ng/dL. Serum Slit2 levels were lower in the diabetic group than in the control group, but there was no significantly significant difference.

Associations between serum Slit2 level and various variables

To investigate the serum levels of Slit2 in human diabetes, we investigated correlations between serum circulating Slit2 level and multiple metabolic parameters, including BMI, WCs, fasting glucose, fasting C-peptide, HbA1c, HOMA-IR, HOMA-β, and lipid profiles (Table 2). Increased serum Slit2 level were negatively correlated with fasting plasma glucose level (coefficient $r=-0.246$, $P=0.008$), postprandial glucose (coefficient $r=-0.233$, $P=0.017$), and HbA1c (coefficient $r=-0.357$, $P<0.001$). In addition, positive correlations were evident between Slit2 level and HOMA-β ($r=0.203$, $P=0.033$). However, Slit2 level was not correlated with hs-CRP level or lipid profiles including TG, TC, LDL-C, and HDL-C.

Table 2. Correlation Analysis between Slit2 Level and Metabolic Parameters

Variable	Coefficient (r) ^a	P value
BMI, kg/m ²	0.143	0.144
Waist circumferences, cm	0.265	0.103
Fasting glucose, mg/dL	-0.246	0.008 ^b
Fasting insulin, mIU/L	0.055	0.572
Fasting C-peptide, mIU/L	0.042	0.666
Postload 2-hour glucose, mg/dL	-0.233	0.017 ^c
Postload 2-hour insulin, mIU/L	0.154	0.119
Postload 2-hour C-peptide, mIU/L	0.110	0.275
HbA1c, %	-0.357	<0.001 ^b
HOMA-IR	-0.136	0.157
HOMA-β	0.203	0.033 ^c
TG, mg/dL	0.021	0.831
LDL-C, mg/dL	-0.138	0.155
HDL-C, mg/dL	-0.032	0.742
TC, mg/dL	-0.170	0.078
hs-CRP, mg/dL	-0.084	0.486

BMI, body mass index; HbA1c, glycosylated hemoglobin; HOMA-IR, homeostasis model assessment of insulin resistance; HOMA-β, homeostasis model assessment of β-cell; TG, triglycerides; LDL-C, low density lipoprotein cholesterol; HDL-C, high density lipoprotein cholesterol; TC, total cholesterol; hs-CRP, high-sensitivity C-reactive protein.

^aCoefficients (r) were calculated using Spearman's method; ^b $P<0.01$; ^c $P<0.05$.

DISCUSSION

In our study, we demonstrated for the first time that the circulating Slit2 was linked to human serum glucose level and insulin secretory function. Our results showed that human circulating Slit2 was negatively correlated with serum glucose, HOMA-β, and HbA1c, although there was no significant difference statistically between diabetic group and control group.

Slit2 was first characterized in neuronal development [10] and was found to interact with Robo1 to mediate myogenesis [11], endothelial cell migration, and leukocyte chemotaxis [12]. The role of Slit2 has been shown to induce tumor angiogenesis and to regulate the lymphangiogenesis in various human cancers [13,14]. The full-length Slit2 protein is a secreted 200 kDa ligand that is cleaved into two fragments, a 140 kDa N-terminal product (Slit2-N), and a 50 to 60 kDa C-terminal product (Slit2-C) [15]. Many studies have revealed that N-Slit2 binds cell membrane via Robo receptor, whereas Slit2-C is diffusible [16]. However, few studies focused on the role of Slit2-N in human diabetes. Increased levels of advanced glycation end-products can increase the levels of Slit2 in human retinal pigment epithelium cells via the PI3K (phosphatidylinositol-4,5-bisphosphate 3-kinase)/Akt pathway in patients with diabetic retinopathy [17]. Additionally, Zhao et al. [18] reported the Slit2-N inhibited lipopolysaccharide-induced endothelial inflammation via binding with Robo4 receptor. However, the direct role of Slit2 in metabolic disease and glucose homeostasis is not fully understood.

Recently, Svensson et al. [4] demonstrated the functional specificity of circulating Slit2, the active C fragments of Slit2, in regulating thermogenesis. They identified that the Slit2 was the new secreted factor from beige adipose cells using PRDM16, the major transcriptional coregulator that controls the development of brown adipocytes [19], transgenic mice, and cold exposure experiments. Moreover, they demonstrated that the recombinant Slit2 protein induced a thermogenic program in mice and in cells, and observed the reduction of in the thermogenic gene expression and expression of mitochondrial genes in brown adipocytes in whole body Slit2 knockout mice. Notably, they identified that the Slit2-C, which is a circulating active form of Slit2, regulated whole-body energy expenditure and induced the improvement of glucose homeostasis in obese mice, which was distinct from previous studies of Slit2. Despite the fact that Slit2 protein expression in adipose tissue may be related to thermogenesis *in vivo* [4], the function of circulating Slit2 proteins in humans has not been studied. For these reasons, we evaluated

serum circulating Slit2 level in human for the first time.

The results of our study are consistent with previous studies on Slit2-C, since circulating Slit2 was negatively correlated with serum fasting glucose level, serum postprandial glucose level, and HbA1c level in human, although there is no significant difference of circulating Slit2 level between T2DM subjects and control subjects. Interestingly, HOMA- β index, an indicator for β -cell function, was significantly positively related with circulating Slit2 level. From our result, we suggest serum circulating Slit2 may have a specific role in the regulation of glucose homeostasis and in the maintenance of β -cell functions, although further large scaled studies are still required.

Even though our study clearly showed that serum circulating Slit2 levels are significantly related with glucose homeostasis and β -cell function in humans, as in previous studies [4], we could not evaluate the exact roles and mechanisms of Slit2. Further studies are necessary, since little is known about the exact receptor and the enzymatic processing of Slit2-C. Furthermore, Slit2-C has multiple domains, and the mechanism underlying these processes should be explored by research *in vivo* and *in vitro*, including through binding assays of Slit2-C and the expression of receptors specific for circulating Slit2-C.

In conclusion, these preliminary results suggest a possible association: circulating Slit2 may play a pivotal role in maintenance of glucose homeostasis in humans.

CONFLICTS OF INTEREST

No potential conflict of interest relevant to this article was reported.

ACKNOWLEDGMENTS

This research was supported by the National Research Foundation of Korea (NRF) funded by the Ministry of Science, ICT & Future Planning (NRF-2017R1D1A1B03027820), the National Research Foundation of Korea (NRF) grant funded by the Korean Government (NRF-2016R1D1A1A09916900), and the research fund of the Daejeon and Chungcheong Branch of Korean Endocrine Society (2017).

ORCID

Yea Eun Kang <https://orcid.org/0000-0002-2012-3716>
 Sorim Choung <https://orcid.org/0000-0001-7814-3034>
 Ju Hee Lee <https://orcid.org/0000-0001-5976-7175>

Hyun Jin Kim <https://orcid.org/0000-0002-6760-4963>

Bon Jeong Ku <https://orcid.org/0000-0002-3414-8949>

REFERENCES

- Villarroya F, Cereijo R, Villarroya J, Giralt M. Brown adipose tissue as a secretory organ. *Nat Rev Endocrinol* 2017;13:26-35.
- Cannon B, Nedergaard J. Brown adipose tissue: function and physiological significance. *Physiol Rev* 2004;84:277-359.
- Ku CR, Cho YH, Hong ZY, Lee H, Lee SJ, Hong SS, et al. The effects of high fat diet and resveratrol on mitochondrial activity of brown adipocytes. *Endocrinol Metab (Seoul)* 2016;31:328-35.
- Svensson KJ, Long JZ, Jedrychowski MP, Cohen P, Lo JC, Serag S, et al. A secreted Slit2 fragment regulates adipose tissue thermogenesis and metabolic function. *Cell Metab* 2016;23:454-66.
- Brose K, Tessier-Lavigne M. Slit proteins: key regulators of axon guidance, axonal branching, and cell migration. *Curr Opin Neurobiol* 2000;10:95-102.
- Mehlen P, Delloye-Bourgeois C, Chedotal A. Novel roles for Slits and netrins: axon guidance cues as anticancer targets? *Nat Rev Cancer* 2011;11:188-97.
- American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care* 2010;33 Suppl 1:S62-9.
- Kang YE, Kim JM, Joung KH, Lee JH, You BR, Choi MJ, et al. The roles of adipokines, proinflammatory cytokines, and adipose tissue macrophages in obesity-associated insulin resistance in modest obesity and early metabolic dysfunction. *PLoS One* 2016;11:e0154003.
- Shin MY, Kim JM, Kang YE, Kim MK, Joung KH, Lee JH, et al. Association between growth differentiation factor 15 (GDF15) and cardiovascular risk in patients with newly diagnosed type 2 diabetes mellitus. *J Korean Med Sci* 2016; 31:1413-18.
- Wong K, Park HT, Wu JY, Rao Y. Slit proteins: molecular guidance cues for cells ranging from neurons to leukocytes. *Curr Opin Genet Dev* 2002;12:583-91.
- Kramer SG, Kidd T, Simpson JH, Goodman CS. Switching repulsion to attraction: changing responses to slit during transition in mesoderm migration. *Science* 2001;292:737-40.
- Wu JY, Feng L, Park HT, Havlioglu N, Wen L, Tang H, et al. The neuronal repellent Slit inhibits leukocyte chemotaxis induced by chemotactic factors. *Nature* 2001;410:948-52.
- Wang B, Xiao Y, Ding BB, Zhang N, Yuan Xb, Gui L, et al.

- Induction of tumor angiogenesis by Slit-Robo signaling and inhibition of cancer growth by blocking Robo activity. *Cancer Cell* 2003;4:19-29.
14. Yang XM, Han HX, Sui F, Dai YM, Chen M, Geng JG. Slit-Robo signaling mediates lymphangiogenesis and promotes tumor lymphatic metastasis. *Biochem Biophys Res Commun* 2010;396:571-7.
 15. Brose K, Bland KS, Wang KH, Arnott D, Henzel W, Goodman CS, et al. Slit proteins bind Robo receptors and have an evolutionarily conserved role in repulsive axon guidance. *Cell* 1999;96:795-806.
 16. Kidd T, Brose K, Mitchell KJ, Fetter RD, Tessier-Lavigne M, Goodman CS, et al. Roundabout controls axon crossing of the CNS midline and defines a novel subfamily of evolutionarily conserved guidance receptors. *Cell* 1998;92:205-15.
 17. Zhou W, Yu W, Xie W, Huang L, Xu Y, Li X. The role of SLIT-ROBO signaling in proliferative diabetic retinopathy and retinal pigment epithelial cells. *Mol Vis* 2011;17:1526-36.
 18. Zhao H, Anand AR, Ganju RK. Slit2-Robo4 pathway modulates lipopolysaccharide-induced endothelial inflammation and its expression is dysregulated during endotoxemia. *J Immunol* 2014;192:385-93.
 19. Seale P, Conroe HM, Estall J, Kajimura S, Frontini A, Ishibashi J, et al. Prdm16 determines the thermogenic program of subcutaneous white adipose tissue in mice. *J Clin Invest* 2011;121:96-105.