

## Life-threatening airway obstruction after flapless implant placement in the anterior mandible

Jin-Hong Kim, Hee-Keun Park, Moon-Key Kim, Sang-Hoon Kang

Department of Oral and Maxillofacial Surgery, National Health Insurance Corporation Ilsan Hospital, Goyang, Korea

**Abstract** (J Korean Assoc Oral Maxillofac Surg 2012;38:310-3)

This paper reports a patient who had an airway obstruction caused by a hemorrhage of the mouth floor that occurred after she underwent flapless implantation in the region of the mandibular anterior teeth. The hemorrhage may have been caused by iatrogenic malpositioned fixture and patient's hypertension. The lingual periosteum was not dissected during the flapless procedure. Therefore, when hemorrhage occurred, the blood did not drain easily into the oral cavity but instead drained into the deep neck region, which might have been the cause of the airway obstruction.

**Key words:** Airway obstruction, Dental implants

[paper submitted 2011. 11. 24 / accepted 2012. 2. 20]

### I. Introduction

Complications may occur during dental implantation treatment including fixture fracture, nerve injury, and hemorrhage<sup>1</sup>. Flapless implantation provides the advantages of reduced pain, reduced edema, and shorter operation time but may result in fixture malpositioning since it is a blind method<sup>2</sup>. The anterior mouth floor (AMF) is rich in blood vessels, including the sublingual artery and submental artery<sup>3,4</sup>.

Thus, when a fixture is placed by penetrating the lingual cortex for implantation in the area of the mandibular anterior teeth, severe hemorrhage may occur if the surrounding soft tissue and blood vessels of the AMF area are damaged. Airway obstruction may also occur in rare cases<sup>5</sup>.

In this paper, we report the case of a patient who experienced airway obstruction due to AMF hemorrhage that occurred after she underwent flapless implantation in the region of the mandibular anterior teeth.

### II. Case Report

A 56-year-old female patient came to the emergency room (ER) with chief complaint of dyspnea due to swollen tongue and neck. The medical history was remarkable for hypertension. The dental history showed that, earlier on the day of her visit to the ER, the patient had undergone at a local dental clinic extraction of both lower central incisors followed immediately by flapless implantation. According to the patient's history of chief complaint, there was continuous oozing around the surgical site after the implant was installed at around 3 pm that day. Her dentist could not give her an acceptable explanation for her condition, so the patient visited the ER at 10:30 pm with the swelling aggregated around her neck area.

The patient said her symptoms of edema and dyspnea had begun on the AMF after the implantation. Clinical examination in the ER showed severe swelling of the AMF and tongue, accompanied by ecchymosis and orthopnea. The patient's respiration rate was 20 times per minute, and routine lab results including hemoglobin, T-CO<sub>2</sub>, and PT and aPTT were all within normal limits. Her SpO<sub>2</sub> was over 98%, and systolic and diastolic blood pressure (BP) in sitting position were 160 mmHg and 100 mmHg, respectively. Note, however, that the patient's vital signs worsened in the Semi-Fowler's and supine positions. A computed tomography (CT) of the neck could not be performed immediately after the

---

#### Sang-Hoon Kang

Department of Oral and Maxillofacial Surgery, National Health Insurance Corporation Ilsan Hospital, 100, Ilsan-ro, Ilsandong-gu, Goyang 410-719, Korea

TEL: +82-31-900-0267 FAX: +82-31-900-0343

E-mail: omfs1ksh@daum.net

©This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

patient arrived at the ER since she had difficulty breathing in the supine position. The neck lateral radiography showed that the airway had become obstructed by the swelling of the tongue at the level of the oropharynx.(Fig. 1) The patient was referred to the Oral and Maxillofacial Surgery Department for surgical site management and to the Ear, Nose, and Throat (ENT) Department for airway evaluation.

The patient continuously complained of dyspnea in sitting position until, two hours after her arrival at the ER, the otolaryngologists and ER doctors decided to perform tracheostomy. This was done under local anesthesia by an ENT doctor, with the patient in sitting position. The



**Fig. 1.** Neck lateral plain radiography showing tongue swelling and airway obstruction.

*Jin-Hong Kim et al: Life-threatening airway obstruction after flapless implant placement in the anterior mandible. J Korean Assoc Oral Maxillofac Surg 2012*




**Fig. 2.** Computed tomography axial view showing tongue swelling and airway obstruction.

*Jin-Hong Kim et al: Life-threatening airway obstruction after flapless implant placement in the anterior mandible. J Korean Assoc Oral Maxillofac Surg 2012*

subsequent neck CT showed that the oropharyngeal airway was completely obstructed.(Figs. 2, 3) The implants in the lower central incisor area had been implanted in labioversion in the wrong positions through the lingual cortex.(Fig. 4)

Since the patient's hypertension continued following treatment in the ER, antihypertensive medication was started after a cooperative diagnosis with the Cardiology Department. After tracheostomy, hemostasis was performed under general anesthesia. The hematomas that had spread under the mouth floor were eliminated, and a drainage tube was inserted through the neck. The intraoperative examination showed that the periosteum was tightly bound to the lingual side



**Fig. 3.** Three-dimensionally reconstructed computed tomography image showing airway obstruction.

*Jin-Hong Kim et al: Life-threatening airway obstruction after flapless implant placement in the anterior mandible. J Korean Assoc Oral Maxillofac Surg 2012*



**Fig. 4.** Malpositioned implant fixture penetrating the lingual bony cortex.

*Jin-Hong Kim et al: Life-threatening airway obstruction after flapless implant placement in the anterior mandible. J Korean Assoc Oral Maxillofac Surg 2012*

above where the two fixtures had been implanted. Ruptured genioglossus muscle and moderate bleeding were also observed. Following postoperative concentrated monitoring in the intensive care unit, the swelling of the mouth floor and the tongue showed marked reduction after the operation, and the drainage tube was removed on the second postoperative day. The swollen mouth floor and tongue returned to normal size in postoperative week one. The tracheal intubation was removed, and the patient was discharged. The ecchymosis had also improved to normal state by the second postoperative week.

### III. Discussion

Flapless surgery has several potential advantages including less swelling, pain, surgical time, and need for suturing. Other merits include the preservation of soft and hard tissues and maintenance of blood supply<sup>2,6,7</sup>. Despite these advantages, however, the flapless technique also has several potential shortcomings including the inability to visualize the anatomical landmarks and vital structures, potential thermal trauma to the bone due to limited external irrigation during the drilling osteotomy, and inability to check the vertical endpoint of the implant placement (too shallow/too deep). A flapless approach is also not advisable for the manipulation of circumferential tissues including alveolar bone and mucoperiosteum<sup>2,8,9</sup>.

In the case discussed here, fixtures were implanted using a flapless method immediately after extraction of the mandibular anterior labioversioned teeth. During this process, the implantation path may not have been easily controlled in the area of the labioversioned teeth. Additionally, the fixture - which was larger than the original teeth and was implanted using a blind method - may have damaged the lingual soft tissue since the lingual cortex was pierced by the fixture.

Perforating the lingual cortex of the anterior mandible can invade the floor of the mouth and damage structures in the sublingual area. Sublingual anatomical structures are bound by the mandible laterally and anteriorly, by the base of the tongue medially and posteriorly, by the mucosa superiorly, and by the mylohyoid muscle inferiorly. The lingual space has sublingual artery and vein, perforating submental artery and vein, submandibular ganglion, lingual nerve, hypoglossal nerve, sublingual gland, and submandibular duct.

As a major branch of the lingual artery, the sublingual artery is one of the main nutrient vessels to the AMF. Thus, the importance of the submental artery in supplying blood

to sublingual anatomical structures has been established<sup>3,4</sup>. In addition, the sublingual artery releases several alveolar branches for complementary blood supply to the lingual anterior cortical bone of the mandible<sup>3,4,10</sup>.

The submental artery is the largest cervical branch of the facial artery, coursing anteriorly on the mylohyoid muscle and branching to anastomose with the sublingual artery and with the mylohyoid artery deep into the mylohyoid muscle. At the anterior mandible, it divides into deep and superficial branches, anastomosing with the inferior labial and mental arteries<sup>3,4</sup>. In human cadaver studies, the sublingual artery is located in the mandibular anterior region, and the submental artery perforates the mylohyoid muscle into the anterior mandible<sup>4</sup>. The sublingual artery may enter the lingual cortex of the anterior mandible. The perforating submental artery was present in the anterior mandible, whereas the sublingual artery was small or was missing in the submental area of the mandible<sup>3</sup>.

Mechanical injury to the local arterial plexus may potentially lead to dangerous hemorrhage. Therefore, detailed knowledge of the regional fine arterial anatomy is both helpful and imperative for the surgeon performing implant surgery on the mandible.

In the case discussed here, the iatrogenic complication as well as the patient's medical history of hypertension may have caused the hemorrhage. Furthermore, the dyspnea caused by tongue swelling could be one of the factors that contributed to the patient's rising BP. In the flapless method that was used, the lingual periosteum was not dissected. Thus, when hemorrhage occurred, the blood did not drain easily into the oral cavity but drained into the deep AMF region instead. This may have contributed to the airway obstruction.

When a patient visits doctors due to bleeding after implant surgery, airway management must be the first consideration. Following airway management, hemostasis must follow, with compression with gauze<sup>11</sup>, hemostatic agent<sup>12</sup>, electric cauterization<sup>13</sup>, etc. For severe bleeding, the vessel could be tied or clipped, but it is very difficult to perform those procedures under local anesthesia without general anesthesia. In cases of difficult intraoral approaches, extraoral approaches should also be considered<sup>3,5</sup>. Vascular angiography may be helpful<sup>14</sup>. In the case discussed here, when airway management was completed, compression with gauze and hemostatic agent were initially used, and bleeding was controlled under general anesthesia.

Computer-guided navigation implant surgery affords the possibility of optimizing implant position by engaging

more bone, leading to better primary stability. In computer-navigated flapless implant surgery, postoperative pain and swelling can be reduced, postoperative soft tissue dehiscence can be eliminated, and intraoperative safety for anatomical structures at risk of injury is enhanced<sup>15</sup>. Despite the high cost of equipment<sup>16</sup>, computer-aided navigation technology in dental implantology has many medical benefits, and computer-guided implantation may be used as a predictable procedure in many cases<sup>17</sup>. Nonetheless, it has yet to be verified for use with all implant surgeries.

Airway obstruction due to hemorrhage of the mouth floor can occur after flapless implantation in the region of the mandibular anterior teeth. In the case described here, the iatrogenic malpositioned fixture and the patient's hypertension may have caused the hemorrhage. When hemorrhage occurred during the flapless procedure, the blood did not drain easily into the oral cavity but drained into the deep neck region instead, which may have contributed to the airway obstruction.

## References

1. Lamas Pelayo J, Penarrocha Diago M, Marti Bowen E, Penarrocha Diago M. Intraoperative complications during oral implantology. *Med Oral Patol Oral Cir Bucal* 2008;13:E239-43.
2. Brodala N. Flapless surgery and its effect on dental implant outcomes. *Int J Oral Maxillofac Implants* 2009;24(Suppl):118-25.
3. Bavitz JB, Harn SD, Homze EJ. Arterial supply to the floor of the mouth and lingual gingiva. *Oral Surg Oral Med Oral Pathol* 1994;77:232-5.
4. Hofschneider U, Tepper G, Gahleitner A, Ulm C. Assessment of the blood supply to the mental region for reduction of bleeding complications during implant surgery in the interforaminal region. *Int J Oral Maxillofac Implants* 1999;14:379-83.
5. Woo BM, Al-Bustani S, Ueek BA. Floor of mouth haemorrhage and life-threatening airway obstruction during immediate implant placement in the anterior mandible. *Int J Oral Maxillofac Surg* 2006;35:961-4.
6. Becker W, Goldstein M, Becker BE, Sennerby L. Minimally invasive flapless implant surgery: a prospective multicenter study. *Clin Implant Dent Relat Res* 2005;7(Suppl 1):S21-7.
7. Sclar AG. Guidelines for flapless surgery. *J Oral Maxillofac Surg* 2007;65(7 Suppl 1):20-32.
8. Warrer K, Buser D, Lang NP, Karring T. Plaque-induced peri-implantitis in the presence or absence of keratinized mucosa. An experimental study in monkeys. *Clin Oral Implants Res* 1995;6:131-8.
9. Oh TJ, Shotwell JL, Billy EJ, Wang HL. Effect of flapless implant surgery on soft tissue profile: a randomized controlled clinical trial. *J Periodontol* 2006;77:874-82.
10. Krenkel C, Holzner K, Poisel S. Hematoma of the mouth floor following oral surgery and its anatomical characteristics. *Dtsch Z Mund Kiefer Gesichtschir* 1985;9:448-51. [German]
11. ten Bruggenkate CM, Krekeler G, Kraaijenhagen HA, Foitzik C, Oosterbeek HS. Hemorrhage of the floor of the mouth resulting from lingual perforation during implant placement: a clinical report. *Int J Oral Maxillofac Implants* 1993;8:329-34.
12. Boyes-Varley JG, Lownie JF. Haematoma of the floor of the mouth following implant placement. *SADJ* 2002;57:64-5.
13. Darriba MA, Mendonca-Caridad JJ. Profuse bleeding and life-threatening airway obstruction after placement of mandibular dental implants. *J Oral Maxillofac Surg* 1997;55:1328-30.
14. Zimmerman RA, McLean G, Freiman D, Golestaneh Z, Perez M. The diagnostic and therapeutic role of angiography in lingual arterial bleeding. *Radiology* 1979;133:639-43.
15. Wittwer G, Adeyemo WL, Schicho K, Gigovic N, Turhani D, Enislidis G. Computer-guided flapless transmucosal implant placement in the mandible: a new combination of two innovative techniques. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2006;101:718-23.
16. Ewers R, Schicho K, Undt G, Wanschitz F, Truppe M, Seemann R, et al. Basic research and 12 years of clinical experience in computer-assisted navigation technology: a review. *Int J Oral Maxillofac Surg* 2005;34:1-8.
17. Wittwer G, Adeyemo WL, Schicho K, Figl M, Enislidis G. Navigated flapless transmucosal implant placement in the mandible: a pilot study in 20 patients. *Int J Oral Maxillofac Implants* 2007;22:801-7.