

Supplementary Table S1. Significant factors for recent smoking cessation before policy change (2013–2014) among women smokers

	Univariate analysis			Multivariate analysis		
	OR	95% CI	p-value	OR	95% CI	p-value
Age, yr						
19–29	2.97	0.71–12.39	0.135			
30–39	1.60	0.35–7.36	0.548			
40–49	1					
50–59	0.49	0.05–4.99	0.543			
≥60	1.18	0.23–6.09	0.839			
BMI	0.93	0.83–1.05	0.228			
Diabetes	0.06	0.01–0.50	0.009	0.08	0.01–0.76	0.027
Hypertension	0.66	0.19–2.31	0.518			
Married	0.28	0.11–0.76	0.011	0.35	0.11–1.08	0.068
Education level						
Below elementary	2.44	0.55–10.80	0.238	3.39	0.69–16.81	0.134
Middle school	3.67	0.59–23.01	0.162	4.58	0.72–29.06	0.106
High school	1			1		
Above university	11.62	3.17–43.01	<0.001	6.87	1.71–27.64	0.007
Type of residence						
General type	1					
Apartment	0.84	0.32–2.18	0.712			
Household income						
Lowest quartile (Q1)	0.61	0.17–2.20	0.447			
Low–middle quartile (Q2)	0.39	0.09–1.71	0.209			
High–middle quartile (Q3)	1.20	0.39–4.00	0.760			
Highest quartile (Q4)	1					

OR: odds ratio; CI: confidence interval; BMI: body mass index.