

Supplementary Table S3. Significant factors for recent smoking cessation before policy change (2013–2014) among men smokers

	Univariate analysis			Multivariate analysis		
	OR	95% CI	p-value	OR	95% CI	p-value
Age, yr						
19–29	4.10	2.06–8.13	<0.001	2.16	1.01–4.62	0.046
30–39	1.40	0.69–2.81	<0.001	1.08	0.52–2.25	0.831
40–49	1			1		
50–59	1.07	0.50–2.32	0.856	1.14	0.53–2.46	0.733
≥60	1.73	0.90–3.32	0.100	1.98	1.01–3.87	0.047
BMI	0.94	0.88–1.00	0.051			
Diabetes	0.78	0.38–1.58	0.484			
Hypertension	0.55	0.32–0.95	0.033	0.78	0.42–1.29	0.283
Married	0.32	0.20–0.50	<0.001	0.48	0.29–0.80	0.005
Education level						
Below elementary	0.77	0.36–1.63	0.493			
Middle school	1.24	0.57–2.69	0.586			
High school	1					
Above university	1.27	0.26–2.19	0.419			
Type of residence						
General type	1					
Apartment	0.86	0.54–1.35	0.502			
Household income						
Lowest quartile (Q1)	1.28	0.61–2.66	0.506			
Low–middle quartile (Q2)	1.00	0.55–1.81	0.995			
High–middle quartile (Q3)	0.69	0.39–1.23	0.213			
Highest quartile (Q4)	1					

OR: odds ratio; CI: confidence interval; BMI: body mass index.