Supplementary Table 1. Cox analysis of death according to total 9086 intracerebral hemorrhage patients for 4 years

| | 3-month | | 1-year | | 2-year | | 4-year | |
|-----------------------|-------------------------------|-------------|-------------------------------|-------------|-------------------------------|-------------|-------------------------------|-------------|
| | Mild | Severe | Mild | Severe | Mild | Severe | Mild | Severe |
| | HR (95% CI) | HR (95% CI) |
| Medical facility type | | | | | | | | |
| Low-volume hospital | 1 | | 1 | | 1 | | 1 | |
| High-volume hospital | 0.90 (0.83-0.98)* | 0.0121 | 0.89 (0.82-0.97)† | 0.0046 | 0.97 (0.89-1.05) | 0.3754 | 0.89 (0.81-0.98)* | 0.0181 |
| Surgery type | | | | | | | | |
| No surgery | 1 | | 1 | | 1 | | 1 | |
| Surgery | 1.30 (1.18-1.42) [‡] | < 0.0001 | 1.49 (1.36–1.63) [‡] | < 0.0001 | 1.61 (1.47–1.77) [‡] | < 0.0001 | 1.59 (1.42–1.78) [‡] | < 0.0001 |
| Severity | | | | | | | | |
| Mild | 1 | | 1 | | 1 | | 1 | |
| Severe | 2.05 (1.88–2.23) [‡] | < 0.0001 | 1.91 (1.76–2.07)‡ | < 0.0001 | 1.98 (1.83–2.15) [‡] | < 0.0001 | 2.35 (2.14-2.58) [‡] | < 0.0001 |
| Age | | | | | | | | |
| 18–45 years | 1 | | 1 | | 1 | | 1 | |
| 46–59 years | 1.39 (1.16–1.66) [‡] | 0.0004 | 1.25 (1.04–1.50)* | 0.0162 | 1.20 (1.00-1.44) | 0.0504 | 0.96 (0.77–1.19) | 0.7044 |
| 60–69 years | 1.86 (1.55–2.23) [‡] | < 0.0001 | 1.63 (1.35–1.95)‡ | < 0.0001 | 1.38 (1.15-1.44) [‡] | 0.0006 | 0.94 (0.75-1.17) | 0.5499 |
| ≥70 years | 3.33 (2.81–3.96) [‡] | < 0.0001 | 2.33 (1.96–2.77)‡ | < 0.0001 | 1.67 (1.41–1.99) [‡] | < 0.0001 | 1.11 (0.91–1.37) | 0.3134 |
| Gender | | | | | | | | |
| Male | 1 | | 1 | | 1 | | 1 | |
| Female | 0.83 (0.78-0.90)‡ | < 0.0001 | 0.89 (0.83-0.96) | 0.0025 | 1.00 (0.93-1.08) | 0.9701 | 0.96 (0.88-1.05) | 0.3831 |
| Arrival mode | | | | | | | | |
| EMS | 1 | | 1 | | 1 | | 1 | |
| No EMS | 0.81 (0.73-0.90)‡ | < 0.0001 | 0.85 (0.77-0.94)† | 0.0017 | 0.85 (0.77-0.94)† | 0.0011 | 0.86 (0.77-0.97)* | 0.0142 |
| Medical history | | | | | | | | |
| CCI score | | | | | | | | |
| 0 | 1 | | 1 | | 1 | | 1 | |
| 1 | 0.97 (0.86-1.09) | 0.6382 | 0.90 (0.80-1.02) | 0.0947 | 1.06 (0.94-1.20) | 0.3143 | 0.99 (0.86-1.15) | 0.9353 |
| 2 | 0.85 (0.75-0.96) | 0.0068 | 0.75 (0.67-0.85)‡ | < 0.0001 | 0.82 (0.73-0.92) | 0.0011 | 0.83 (0.72-0.96)* | 0.0127 |
| ≥3 | 0.93 (0.85-1.03) | 0.1734 | 0.68 (0.61-0.75)‡ | < 0.0001 | 0.67 (0.61-0.74)‡ | < 0.0001 | 0.62 (0.55-0.70)‡ | < 0.0001 |

Mild severity = NIHSS \leq 15, GCS \geq 10; severe severity = NIHSS \geq 16, GCS \leq 9. *p<0.05 significance. †p<0.01 significance. †p<0.001 significance. HR: hazard ratio, CI: confidence interval, EMS: emergency medical services, CCI: Charlson comorbidity index, NIHSS: National Institutes of Health Stroke Scale, GCS: Glasgow coma scale

Long-Term ICH Prognosis | Park SW, et al.