

The Prognostic Role of Mitotic Index in Hepatocellular Carcinoma Patients after Curative Hepatectomy

Sang Yun Ha, MD, PhD, Misun Choi, MD, Taebum Lee, MD, Cheol-Keun Park, MD, PhD

Department of Pathology and Translational Genomics, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, Korea

Supplementary Data

Supplementary Table 1. Association between mitotic index and treatment modality

Treatment modality	Low mitotic index	High mitotic index	Total	p-value
TACE	53 (54.1)	46 (59.0)	99 (56.3)	0.38
RFA	29 (29.6)	16 (20.5)	45 (25.6)	-
TACE and RFA	1 (1.0)	0	1 (0.6)	-
Surgery	5 (5.1)	3 (3.8)	8 (4.5)	-
Chemotherapy	1 (1.0)	4 (5.1)	5 (2.8)	-
Radiotherapy	2 (2.0)	2 (2.6)	4 (2.3)	-
PTBD	0	2 (2.6)	2 (1.1)	-
LT ^{a)}	7 (7.1)	5 (6.4)	12 (6.8)	-

Values are presented as number (%). TACE, transcatheter arterial chemoembolization; RFA, radiofrequency ablation; PTBD, percutaneous transhepatic biliary drainage; LT, liver transplantation. ^{a)}Among 12 patients, 7 received preoperative TACE and 4 received preoperative RFA.