

S2 Table. Cox regression analysis of disease-free survival in bulky cervical cancer (n=89)

Variable	No. (%)	HR	95% CI	p-value
FIGO stage				
IA1-IB1	0			
IB2-IIA	89 (100)			
Cell types				
Squamous cell carcinoma	76 (85.4)	1		
Adenocarcinoma	13 (14.6)	1.29	1.09-2.87	0.034
Lymph node status				
Negative	46 (51.7)	1		
Positive	43 (48.3)	1.84	1.11-3.02	0.017
Lymphovascular space invasion				
Negative	37 (41.6)	1		
Positive	52 (58.4)	1.28	0.47-3.45	0.632
Parametrial involvement				
Negative	67 (75.3)	1		
Positive	22 (24.7)	1.217	0.43-3.46	0.713
Vaginal resection margin with tumor				
Negative	82 (92.1)	1		
Positive	7 (7.9)	1.45	0.19-10.92	0.720
Deep stromal invasion				
Negative	12 (13.5)	1		
Positive	77 (86.5)	2.11	0.28-15.92	0.469
V-ATPase expression				
Negative	34 (38.2)	1		
Positive	55 (62.8)	3.8	1.40-10.31	0.005

HR, hazard ratio; CI, confidence interval; FIGO, International Federation of Gynecology and Obstetrics; V-

ATPase, vacuolar-type H⁺ ATPase.