

S3 Table. Univariable Cox proportional hazard model for clinicopathological variables

| | No. | Bladder recurrence-free survival (n=162, event=58) | | | Disease progression-free survival (n=162, event=48) | | | Overall survival (n=162, event=19) | | |
|---------------------------|-----|---|--------------------|---------|--|-------------------|---------|---------------------------------------|-------------------|---------|
| | | No. (%) | HR (95% CI) | p-value | No. (%) | HR (95% CI) | p-value | No. (%) | HR (95% CI) | p-value |
| Age | 162 | 58 (35.80) | 1.00 (0.98-1.03) | 0.859 | 48 (29.63) | 1.03 (1.00-1.06) | 0.100 | 19 (11.73) | 1.03 (0.98-1.08) | 0.223 |
| Intravesical instillation | | | | | | | | | | |
| No | 103 | 7 (6.80) | 1 (ref) | | 41 (39.81) | 1 (ref) | | 14 (13.59) | 1 (ref) | |
| Yes | 59 | 51 (86.44) | 22.08 (9.96-48.94) | < 0.001 | 7 (11.86) | 0.22 (0.10-0.49) | < 0.001 | 5 (8.47) | 0.43 (0.15-1.20) | 0.108 |
| ISUP | | | | | | | | | | |
| Low | 32 | 19 (59.38) | 1 (ref) | | 5 (15.63) | 1 (ref) | | 2 (6.25) | 1 (ref) | |
| High | 120 | 39 (32.50) | 0.57 (0.33-0.98) | 0.042 | 41 (34.17) | 3.02 (1.19-7.68) | 0.020 | 16 (13.33) | 2.99 (0.68-13.07) | 0.146 |
| pT category | | | | | | | | | | |
| Ta+T1+CIS only | 35 | 20 (57.14) | 1 (ref) | | 2 (5.71) | 1 (ref) | | 1 (2.86) | 1 (ref) | |
| T2+T3+T4 | 124 | 38 (30.65) | 0.49 (0.28-0.84) | 0.010 | 46 (37.1) | 8.72 (2.11-35.95) | 0.003 | 18 (14.52) | 6.83 (0.91-51.35) | 0.062 |

HR, hazard ratio; CI, confidence interval; ISUP, International Society of Urological Pathology; CIS, carcinoma *in situ*.