

**S4 Table.** Incidence of TEAEs and ADRs leading to fatal outcome with clofarabine use  
(n=60)

	Incidence of TEAEs	Incidence of ADRs
<b>Leading to fatal</b>	14 (23.3) [16]	6 (10.0) [7]
Infections and infestations	11 (18.3) [12]	5 (8.33) [5]
Septic shock	4 (6.7) [4]	1 (1.7) [1]
Sepsis	3 (5.0) [3]	2 (3.3) [2]
Escherichia sepsis	1 (1.7) [1]	-
Herpes simplex sepsis	1 (1.7) [1]	-
Lung infection	1 (1.7) [1]	1 (1.7) [1]
Pneumonia	1 (1.7) [1]	1 (1.7) [1]
Pneumonia fungal	1 (1.7) [1]	-
Blood and lymphatic system disorders	2 (3.3) [2]	2 (3.3) [2]
Febrile neutropenia	2 (3.3) [2]	2 (3.3) [2]
Nervous system disorders	1 (1.7) [1]	-
Cerebral infarction	1 (1.7) [1]	-
Respiratory, thoracic and mediastinal disorders	1 (1.7) [1]	-
Acute respiratory distress syndrome	1 (1.7) [1]	-

MedDRA ver. 20.1. Note: Denominator of percentage is the number of patients. Adverse events are displayed as 'number of patients (percentage of patients) [number of events]'. ADRs, adverse drug reactions; n, total patients analyzed; TEAEs, treatment-emergent adverse events.