S1 Table. Compared participants' characteristics for included and excluded due to missing answers of TV viewing time

| | TV data | TV missing | p for difference |
|---|-------------|-------------|------------------|
| No. at risk | 90,834 | 19,481 | |
| Age, mean±SD (yr) | 58 ± 10 | 58 ± 10 | 0.24 |
| BMI, , mean±SD (kg/m²) | 22.8±3.0 | 22.8±3.0 | 0.25 |
| BMI $\geq 25 \text{ kg/m}^2 (\%)$ | 19.7 | 19.2 | 0.07 |
| $BMI \ge 30 \text{ kg/m}^2 (\%)$ | 1.6 | 1.5 | 0.13 |
| Family history of colorectal cancer (%) | 2.2 | 0.5 | < 0.001 |
| Exercise ≥3 hr/wk (%) | 12.0 | 11.3 | 0.28 |
| Walking ≥ 30 min/day (%) | 70.6 | 63.5 | < 0.001 |
| College or higher education (%) | 10.9 | 3.2 | < 0.001 |
| Bowel movement ≥ 4 days (%) | 2.9 | 2.6 | 0.19 |
| Daily sleeping time (hr/day) | 7.3 | 7.3 | 0.99 |
| Current smokers (%) | 26.4 | 27.7 | < 0.001 |
| Current drinkers (%) | 45.2 | 51.8 | < 0.001 |
| Consumption of beef ≥ 3 times/wk (%) | 9.5 | 12.8 | < 0.001 |
| Consumption of pork ≥ 3 times/wk (%) | 20.7 | 29 | < 0.001 |

p for difference was adjusted for age.