

S2 Table. Clinical and pathological characteristics of 129 resectable PC patients in validation set

Characteristics	No. (%)
Sex	
Male	76 (58.9)
Female	53 (41.1)
Age (yr)	
≤ 65	109 (84.5)
> 65	20 (15.5)
Tumor location	
Head	91 (70.5)
Neck, body, and tail	38 (29.5)
Maximum tumor diameter (cm)	
≤ 2.9	43 (33.3)
> 2.9	86 (66.7)
Degree of differentiation	
Well-moderate	113 (87.6)
Poor	16 (12.4)
T category	
1	26 (20.2)
2	71 (55.0)
3	32 (24.8)
N category	
0	93 (72.1)
1	32 (24.8)
2	4 (3.1)
TNM	
I	73 (56.6)
II	52 (40.3)
III	4 (3.1)
CEA (ng/mL)	
≤ 4.42	74 (57.4)
> 4.42	55 (42.6)
CA199 (U/mL)	
≤ 49.77	31 (24.0)
> 49.77	98 (76.0)
NLR	
≤ 3.09	95 (73.6)
> 3.09	34 (26.4)
PLR	
≤ 139.63	77 (59.7)
> 139.63	52 (40.3)
MLR	
≤ 0.45	101 (78.3)
> 0.45	28 (21.7)

FAR	
≤ 0.09	68 (52.7)
> 0.09	61 (47.3)

CA199, serum carbohydrate antigen 199; CEA, carcinoembryonic antigen; FAR, fibrinogen-to-albumin ratio; MLR, monocyte-to-lymphocyte ratio; NLR, neutrophil-to-lymphocyte ratio; PC, pancreatic cancer; PLR, platelet-to-lymphocyte ratio.