

S4 Table. Clinical and pathological characteristics of 273 locally and metastatic pancreatic cancer patients in validation set

Characteristics	No. (%)
Sex	
Male	177 (64.8)
Female	96 (35.2)
Age (yr)	
≤ 52	24 (8.8)
> 52	249 (91.2)
Tumor location	
Head	157 (57.5)
Neck, body, and tail	116 (42.5)
Distant metastasis	
Yes	155 (56.8)
Locally advanced	118 (43.2)
Aggressive treatments	
Yes	118 (43.2)
No	155 (56.8)
CEA (ng/mL)	
≤ 25.98	221 (81.0)
> 25.98	52 (19.0)
CA199 (U/mL)	
≤ 582.9	138 (50.5)
> 582.9	135 (49.5)
NLR	
≤ 2.61	101 (37.0)
> 2.61	172 (63.0)
PLR	
≤ 170.73	200 (73.7)
> 170.73	73 (26.7)
MLR	
≤ 0.36	164 (60.1)
> 0.36	109 (39.9)
FAR	
≤ 0.079	131 (48.0)
> 0.079	142 (52.0)

CA199, serum carbohydrate antigen 199; CEA, carcinoembryonic antigen; FAR, fibrinogen-to-albumin ratio; MLR, monocyte-to-lymphocyte ratio; NLR, neutrophil-to-lymphocyte ratio; PLR, platelet-to-lymphocyte ratio.