

S7 Table. Risk factors for EFS for high and very high risk patients (n=231)

	Patients (events)	10-Year EFS (\pm SE, %)	p-value
Age at diagnosis (yr)			
< 1	21 (12)	42.9 \pm 10.8	0.001
1 to < 10	100 (31)	65.8 \pm 5.3	
\geq 10	110 (31)	71.5 \pm 4.5	
WBC at diagnosis ($\times 10^9/L$)			
< 50	135 (39)	68.2 \pm 4.5	0.021
\geq 50 and < 100	40 (10)	74.1 \pm 7.1	
\geq 100	56 (25)	56.5 \pm 6.7	
Initial CNS involvement			
No	224 (69)	67.4 \pm 3.4	0.010
Yes	6 (4)	33.3 \pm 19.2	
Prephase steroid response			
Good	196 (58)	67.8 \pm 3.7	0.030
Poor	35 (16)	57.1 \pm 8.4	
Genetic abnormality			
<i>ETV6-RUNX1</i>	26 (9)	61.5 \pm 10.8	0.007
<i>E2A-PBX1</i>	30 (4)	86.7 \pm 6.2	
<i>BCR-ABL1</i>	31 (15)	50.5 \pm 9.2	
<i>KMT2A</i>	22 (10)	54.5 \pm 10.6	
High hyperdiploidy	34 (4)	88.2 \pm 5.5	
Hypodiploidy	2 (1)	0	
Normal	28 (11)	64.3 \pm 9.1	
Others	56 (18)	65.0 \pm 6.9	

CNS, central nervous system; EFS, event-free survival; SE, standard error; WBC, white blood cell.