

S3 Table. Metabolic parameters measured by FDG PET/CT

		All (N=15)	SD group (N=8)	PD group (N=7)	p-value ^{a)}
Pre-treatment	SUVmax	9.05 (3.76)	9.03 (4.06)	9.08 (3.70)	0.867
	SUVavg	5.33 (2.44)	5.40 (2.67)	5.24 (2.36)	0.908
	MTV	53.83 (67.45)	24.62 (24.19)	87.21 (86.58)	0.072
	TLG	244.78 (270.26)	119.12 (129.05)	388.39 (325.40)	0.054
Post-treatment	SUVmax	8.81 (3.76)	8.35 (4.40)	9.33 (3.12)	0.463
	SUVavg	5.11 (2.38)	4.86 (2.69)	5.41 (2.14)	0.397
	MTV	75.54 (84.30)	27.46 (27.67)	130.50 (95.29)	0.009
	TLG	354.67 (429.97)	107.09 (105.50)	637.62 (493.21)	0.006
Change	SUVmax	-0.24 (1.88)	-0.67 (1.65)	0.25 (2.14)	0.463
	SUVavg	-0.21 (1.08)	-0.54 (0.69)	0.16 (1.35)	0.536
	MTV	21.71 (37.36)	2.84 (6.90)	43.28 (46.73)	0.040
	TLG	109.90 (291.27)	-12.03 (33.41)	249.23 (392.67)	0.006

Values are presented as mean (standard deviation). FDG, 18F-Fludeoxyglucose; PD, progressive disease; PET/CT, positron emission tomography/computed tomography; SD, stable disease. ^{a)}The Mann-Whitney test was used as a statistical comparison method. Significant probabilities ($p < 0.05$).