

**S5 Table.** Results of multivariate Cox PH analysis of age variables categorized according to the GELTAMO-IPI criterion in DLBCL-NOS subgroup (n=559)

Parameter	OS			PFS		
	HR	95% CI	p-value	HR	95% CI	p-value
<b>Age-adjusted IPI risk</b>						
Low	1	-	-	1	-	-
Low-int.	1.53	0.80-2.92	0.195	1.83	1.14-2.95	0.013
High-int.	3.96	2.28-6.89	<0.0001	3.11	1.99-4.85	< 0.001
High	4.28	1.94-9.40	0.0003	3.62	1.88-6.98	0.0001
<b>ARDI</b>						
< 85%	1.04	0.68-1.60	0.856	1.05	0.73-1.51	0.802
Presence of B symptoms	1.56	1.02-2.40	0.041	1.45	1.00-2.10	0.050
<b>Age (yr)</b>						
< 65	1	-	-	1	-	-
65-80	1.48	0.95-2.31	0.083	1.45	1.00-2.11	0.049
≥ 80	4.49	1.59-12.68	0.005	2.59	0.93-7.18	0.068

ARDI, average relative dose intensity; CI, confidence interval; DLBCL, diffuse large B-cell lymphoma; GELTAMO, Grupo Español de Linfomas/Trasplante Autólogo de Médula Ósea; HR, hazard ratio; int., intermediate; IPI, international prognostic index; NOS, not otherwise specified; OS, overall survival; PFS, progression-free survival; PH, proportional hazard.