	Unknown BMI
	(n=5,058)
Age (yr)	39.6±9.4
20-39	2,943 (58.2)
40-64	2,065 (40.8)
$\geq$ 65	50 (1.0)
Male sex	4,396 (86.9)
WC (cm)	87.3±13.3
Rural residence	3,588 (70.9)
Low income <sup>a)</sup>	623 (12.3)
Smoking	
Never	1,925 (38.1)
Former	977 (19.3)
Current	2,156 (42.6)
Alcohol	
None	1,724 (34.1)
Mild <sup>b)</sup>	2,800 (55.4)
Heavy <sup>c)</sup>	534 (10.6)
Regular exercise	3,331 (65.9)
Diabetes	505 (10.0)
Hypertension	1,350 (26.7)
Dyslipidemia	963 (19.0)

S1 Table. Baseline characteristics of those who with unknown BMI

Values are presented as mean±SD or number (percentage excluding missing data). BMI, body mass index; WC, waist circumference. <sup>a)</sup>The lowest quartile range of yearly income of those under National Health Insurance Service and the income of those under Medical Aid which is for the recipients of National Basic Livelihood Security Program, <sup>b)</sup>Alcohol consumption < 30 g/day, <sup>c)</sup>Alcohol consumption  $\geq$  30 g/day.