

S5 Table. BOR as per investigator assessment in patients with brain metastasis

| | 450-mg Fed (n=13) | 600-mg Fed (n=8) | 750-mg Fed (n=7) |
|----------------------|----------------------|---------------------|---------------------|
| BOR, n (%) | | | |
| Complete response | 0 | 0 | 0 |
| Partial response | 8 (61.5) | 5 (62.5) | 5 (71.4) |
| Stable disease | 4 (30.8) | 2 (25.0) | 1 (14.3) |
| Progressive disease | 0 | 0 | 1 (14.3) |
| Unknown | 1 (7.7) | 1 (12.5) | 0 |
| ORR: CR+PR | | | |
| N (%) | 8 (61.5) | 5 (62.5) | 5 (71.4) |
| 95% CI | 31.6-86.1 | 24.5-91.5 | 29.0-96.3 |
| DCR: CR+PR+SD | | | |
| N (%) | 12 (92.3) | 7 (87.5) | 6 (85.7) |
| 95% CI | 64.0-99.8 | 47.3-99.7 | 42.1-99.6 |

BOR, best overall response; CI, confidence interval; CR, complete response; DCR, disease control rate; N, number of patients included in the analysis; ORR, overall response rate; PR, partial response; SD, stable disease.