

S4 Table. Primers used for quantitative reverse-transcription polymerase chain reaction analyses

Gene	Forward Primers (5'-3')	Reverse primers (5'-3')
CUL1	AGCTTTCTACGCCAGCCG	AGCAGGATAGCCATCTGGAA
CUL2	CTCCGTGCTGTGTCCACTG	CACAAATAGTGTTGGCATGT
CUL3	AGTCGTGTGCCAAATGGT	GGATATAGTCAACAGGAT
CUL4A	ACCATTGATGGAATCCTACTGC	ACCTTTGGCCTTCGGCAGCATA
CUL4B	ACTTCGTGCAGGCAACAAAGA	CAGCATCTACAGATGCACTCT
CUL5	TGCAAGAGATAAGGCGTATA	GCTCAGGGCATT TTTGATTC
CUL7	GGCACCTGATCTCGGAG	TGGGCTCACATCTGCCAGCT
CUL9	AGCCAGGGATGCGAGTGCG	CTGCCAGAAAACCTGCACA
FBXL1	CAGCTTGTTATGGGTCCAA	CAGTGGGTGTAAACAGCATT
HIF1A	TGGAGGACTGTGACTGCGCG	GCCTCGGTGCAGCCCATGA
MEN1	AGCACATGGACCTATCGAAC	GACAATGGGATCCGAAAGC
P65	CAAGGTGCAGAAAGAGGACA	GGCTGGGGTCTGCGTAGGGA
P50	TCCTGGACTCTTGGTGCACCC	CTCCTTGGTCTGCTGCAGAGC
c-MYC	GACCGAGCTGCTGGGAGGAGA	AGAAGCCGCTCCACATACAGT