

S3 Table. Interactive association of eGFR and dipstick proteinuria on MM incidence

	eGFR (categorized into five different levels) (mL/min/1.73 m ²)									
	< 30		30-60		60-90		90-120		≥ 120	
Relative risk, HR (95% CI) ^{a)}										
Absent	1.22	0.91-1.63	1.21	1.08-1.36	1.00	Referent	0.92	0.85-1.00	1.02	0.88-1.18
Trace	1.72	0.55-5.33	1.50	0.93-2.42	1.32	1.04-1.68	1.58	1.12-2.23	1.04	0.43-2.49
1+	4.13	1.96-8.67	2.04	1.38-3.00	2.19	1.78-2.70	1.94	1.35-2.78	2.03	0.97-4.27
2+	6.49	3.09-13.64	3.21	2.10-4.89	3.01	2.25-4.02	2.23	1.26-3.93	4.32	1.80-10.39
3+ and 4+	7.45	2.79-19.85	3.67	1.97-6.84	3.91	2.45-6.22	4.11	1.84-9.16	0 (N/A)	
Incidence rate ^{b)}										
Absent	4.38		12.23		4.83		3.26		3.65	
Trace	7.11		14.25		6.97		6.06		4.10	
1+	23.71		19.96		12.79		8.13		8.83	
2+	47.25		30.71		18.71		9.97		20.12	
3+ and 4+	58.98		34.35		25.87		19.68		0 (N/A)	

CI, confidence interval; eGFR, estimated glomerular filtration rate; HR, hazard ratio; MM, multiple myeloma; N/A, not applicable. ^{a)}100,000 person-years, ^{b)}Adjusted for age, sex, income level, body mass index, smoking status, regular exercise, hypertension, diabetes and dyslipidemia.