



Testis	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Kidney	1.7	4.1	1999-2008	6.6 <sup>a)</sup>	2008-2019	2.4 <sup>a)</sup>	-	-	0.6	0.4	1999-2019	-0.9 <sup>a)</sup>	-	-	-	-
Bladder	1.6	1.4	1999-2003	2.6	2003-2012	-2.0 <sup>a)</sup>	2012-2019	0.1	0.5	0.4	1999-2001	11.2	2001-2019	-2.0 <sup>a)</sup>	-	-
Brain and CNS	2.6	2.6	1999-2019	0.1	-	-	-	-	1.7	1.4	1999-2003	3.9 <sup>a)</sup>	2003-2008	-4.6 <sup>a)</sup>	2008-2019	-0.6
Thyroid	10.7	65.6	1999-2011	22.1 <sup>a)</sup>	2011-2016	-13.3 <sup>a)</sup>	2016-2019	5.2	0.5	0.3	1999-2004	4.3	2004-2019	-4.8 <sup>a)</sup>	-	-
Hodgkin lymphoma	0.1	0.4	1999-2019	4.8 <sup>a)</sup>	-	-	-	-	0.0	0.0	1999-2003	47.9	2003-2019	-1.5	-	-
Non-Hodgkin lymphoma	3.3	5.0	1999-2019	2.6 <sup>a)</sup>	-	-	-	-	1.4	1.2	1999-2019	-0.7 <sup>a)</sup>	-	-	-	-
Multiple myeloma	0.8	1.5	1999-2006	6.1 <sup>a)</sup>	2006-2019	2.1 <sup>a)</sup>	-	-	0.4	0.6	1999-2005	9.8 <sup>a)</sup>	2005-2016	0.3	2016-2019	-6.9
Leukemia	4.0	4.7	1999-2019	0.7 <sup>a)</sup>	-	-	-	-	2.4	1.5	1999-2019	-2.1 <sup>a)</sup>	-	-	-	-
Other and ill-defined	12.4	20.0	1999-2010	3.4 <sup>a)</sup>	2010-2019	1.7 <sup>a)</sup>	-	-	7.1	3.3	1999-2005	-7.3 <sup>a)</sup>	2005-2019	-1.9 <sup>a)</sup>	-	-

APC was calculated using age-standardized incidence data based on the Segi's world standard population. APC, annual percentage change; CNS, central nervous system. <sup>a)</sup>

Significantly different from zero ( $p < 0.05$ ), <sup>b)</sup> Includes the gallbladder and other/unspecified parts of the biliary tract.