

S1 Table. Demographics and diagnostic characteristics of 103 NSCLC patients

Clinical characteristics	FFPE tissue samples (n=65)	cfDNA from blood samples (n=38)
Age (yr), median (min-max)	65 (43-91)	68 (45-93)
Sex		
Male	32 (49.23)	28 (47.37)
Female	33 (50.77)	20 (52.63)
Histology		
Adenocarcinoma	42 (64.62)	
Carcinoma	10 (15.38)	
Non-small cell carcinoma	8 (12.30)	
Squamous carcinoma	1 (1.54)	
Non-neuroendocrine carcinoma	2 (3.08)	
Malignant epithelial tumor	2 (3.08)	
<i>EGFR</i> mutational status		
19Del	11 (16.92)	8 (21.05)
19Del, T790M (c.2369C>T)	5 (7.69)	4 (10.53)
L858R (c.2573T>G)	13 (20)	6 (15.79)
T790M (c.2369C>T), L858R (c.2573T>G)	5 (7.69)	2 (5.26)
<i>EGFR</i> wild type	31 (47.70)	18 (47.37)

Values are presented as number (%). cfDNA, cell-free DNA; EGFR, epidermal growth factor receptor; FFPE, formalin-fixed, paraffin-embedded; NSCLC, non-small cell lung cancer.