Supplementary Table 1. Hearing thresholds according to groups

	Univariate				Multivariate			
	Group 1	Group 2	Group 3	p-value	Group 1	Group 2	Group 3	p-value
Baselines								
0.5 kHz	73.4 ± 26.4	71.1 ± 27.2	71.5 ± 27.0	0.912	73.8 ± 4.4	$\textbf{70.9} \pm \textbf{4.0}$	71.4 ± 3.8	0.885
1 kHz	76.2 ± 26.5	73.3 ± 26.9	75.4 ± 27.7	0.870	77.3 ± 4.4	72.8 ± 4.0	75.0 ± 3.8	0.763
2 kHz	74.3 ± 24.1	71.6 ± 27.6	73.1 ± 29.0	0.898	75.7 ± 4.5	$\textbf{70.9} \pm \textbf{4.0}$	72.5 ± 3.8	0.737
3 kHz	73.9 ± 24.3	71.1 ± 26.4	71.9 ± 30.9	0.887	75.0 ± 4.5	70.6 ± 4.1	71.5 ± 3.9	0.767
4 kHz	77.6 ± 24.0	72.6 ± 26.5	73.2 ± 31.0	0.656	78.2 ± 4.5	72.2 ± 4.0	72.9 ± 3.8	0.589
8 kHz	81.8 ± 20.7	81.3 ± 26.1	78.7 ± 25.7	0.793	80.9 ± 4.0	81.6 ± 3.5	79.0 ± 3.4	0.858
Day 10								
0.5 kHz	62.2 ± 27.6	58.6 ± 35.6	53.8 ± 33.0	0.454	61.8 ± 5.3	58.7 ± 4.8	53.9 ± 4.5	0.523
1 kHz	66.5 ± 26.8	61.9 ± 33.3	57.2 ± 33.4	0.374	67.0 ± 5.2	61.7 ± 4.7	57.0 ± 4.4	0.361
2 kHz	65.6 ± 24.7	60.5 ± 32.4	54.4 ± 36.0	0.241	66.6 ± 5.3	60.1 ± 4.7	54.0 ± 4.5	0.203
3 kHz	66.7 ± 26.6	63.8 ± 29.0	53.7 ± 36.7	0.107	67.5 ± 5.2	63.4 ± 4.7	$53.3 \pm 4.4^*$	0.097
4 kHz	70.6 ± 22.6	64.7 ± 30.2	57.4 ± 35.9	0.117	71.1 ± 5.0	64.4 ± 4.5	$57.2 \pm 4.3^*$	0.117
8 kHz	78.9 ± 21.7	74.1 ± 30.0	68.8 ± 31.1	0.227	78.8 ± 4.6	71.2 ± 4.1	68.8 ± 3.9	0.265
Day 90								
0.5 kHz	54.0 ± 27.0	51.0 ± 32.1	$36.0 \pm 27.8^{*\dagger}$	0.009	51.9 ± 4.8	51.9 ± 4.3	$36.9\pm4.3^{*\dagger}$	0.022
1 kHz	60.7 ± 25.3	54.3 ± 31.8	$42.8 \pm 32.8^*$	0.022	59.5 ± 5.0	54.8 ± 4.4	$43.4 \pm 4.5^*$	0.048
2 kHz	60.4 ± 22.6	55.0 ± 35.6	$44.7 \pm 34.0^*$	0.045	60.5 ± 4.9	54.9 ± 4.4	$44.6 \pm 4.4^*$	0.054
3 kHz	63.4 ± 24.7	59.4 ± 28.6	$45.7 \pm 35.0^{*\dagger}$	0.015	63.4 ± 5.0	59.4 ± 4.4	$45.6\pm4.5^{*\dagger}$	0.020
4 kHz	68.4 ± 22.1	62.2 ± 28.9	$50.3 \pm 34.1*$	0.014	68.5 ± 4.8	62.2 ± 4.3	$50.3 \pm 4.3^*$	0.018
8 kHz	76.8 ± 23.3	72.0 ± 28.7	62.5±31.1*	0.054	76.4 ± 4.5	72.2 ± 4.1	$62.7 \pm 4.1*$	0.072

The data are expressed as means \pm standard deviations in univariate analysis and means \pm standard errors in multivariate analysis. In univariate and multivariate analyses, the *p*-values were tested using a one-way analysis of variance followed by post-hoc Turkey comparisons, and an analysis of covariance followed by an LSD post-hoc test, respectively. Multivariate analyses were adjusted for age and duration from onset to treatment. *p<0.05 vs. Group 1, †p<0.05 vs. Group 2