

**Supplementary Table 1.** Subgroup analyses for the effect of prolonged DAPT on the primary end point

Subgroup	Prolonged DAPT	Clopidogrel	HR (95% CI)	DAPT better	Clopidogrel better	<i>P</i> value	<i>P</i> value for interaction
Age, yr							0.18
≥ 65	11/180	9/131	0.97 (0.40–2.35)			0.95	
< 65	6/191	1/135	1.49 (0.37–5.96)			0.57	
Acute coronary syndrome							0.19
Yes	11/178	5/131	1.76 (0.61–5.07)			0.30	
No	6/193	7/135	0.625 (0.22–1.94)			0.65	
STEMI							0.13
Yes	6/58	1/39	4.10 (0.49–34.06)			0.19	
No	11/313	11/227	0.72 (0.31–1.67)			0.45	
Diabetes							0.50
Yes	9/177	5/130	1.42 (0.48–4.24)			0.53	
No	8/194	7/136	0.87 (0.32–2.41)			0.80	
Chronic kidney disease							0.50
Yes	4/35	4/24	0.74 (0.19–2.98)			0.68	
No	13/336	8/242	1.28 (0.53–3.08)			0.59	
Total stent length ≥ 28 mm							0.95
Yes	7/171	7/165	1.08 (0.38–3.08)			0.89	
No	10/200	5/101	1.08 (0.37–3.16)			0.89	
Multivessel PCI							0.32
Yes	2/104	4/96	0.49 (0.09–2.69)			0.41	
No	15/267	8/170	1.31 (0.56–3.10)			0.53	
Everolimus stent							0.50
Yes	9/170	7/168	1.42 (0.53–3.80)			0.49	
Other DES	8/201	5/98	0.83 (0.27–2.54)			0.75	
Overall	371	266	0.87 (0.40–1.87)			0.71	

DAPT = dual antiplatelet therapy, STEMI = ST-segment elevation myocardial infarction, PCI = percutaneous coronary intervention, DES = drug-eluting stent, HR = hazard ratio, CI = confidence interval.