

Supplementary Table 1. Scores of the questionnaire for symptoms

	Mild intermittent	Mild persistent	Moderate to severe intermittent	Moderate to severe persistent	Total	p
Sneezing	0.48±0.57	0.89±0.66	2.22±1.45	2.38±1.61	2.00±1.59	<0.001*
Watery rhinorrhea	0.45±0.64	0.74±0.65	2.35±1.57	2.54±1.72	2.12±1.72	<0.001*
Nasal obstruction	0.60±0.78	1.32±0.75	3.48±1.24	4.06±1.15	3.29±1.67	<0.001*
Itchy nose or eyes	0.43±0.71	0.74±0.81	2.44±1.66	2.50±1.73	2.11±1.75	<0.001*
PND	0.28±0.55	1.11±1.33	1.69±1.55	2.57±1.80	1.93±1.78	<0.001*
Combined with eye symptoms	0.13±0.40	0.37±0.60	1.60±1.67	1.60±1.73	1.33±1.65	<0.001*
Throat discomfort	0.55±0.87	1.95±1.51	2.03±1.65	2.70±1.84	2.16±1.82	<0.001*
Chronic cough	0.18±0.58	0.32±0.58	0.67±1.07	1.05±1.47	0.79±1.29	<0.001*
Ear fullness	0.19±0.58	0.26±0.73	0.89±1.30	1.29±1.49	0.98±1.38	<0.001*
Headache	0.20±0.46	0.74±1.10	1.52±1.59	1.85±1.71	1.48±1.64	<0.001*
Difficulty concentrating	0.22±0.47	0.89±0.94	1.96±1.48	2.49±1.60	1.95±1.64	<0.001*
Sleep disturbance	0.24±0.57	0.58±0.69	2.09±1.65	2.80±1.66	2.15±1.78	<0.001*
Impairment of daily activities	0.20±0.48	0.95±0.78	2.41±1.45	3.24±1.45	2.48±1.71	<0.001*

Data are presented as mean±standard deviation. *statistical analysis was performed using the Kruskal–Wallis test. PND, post-nasal drip