

Supplementary Table 2. Primer sequences used for quantitative polymerase chain reaction analysis

Gene	Prime sequence	Tm (°C)
<i>Nestin</i>	5'-CCCTGAAGTCGAGGAGCTG-3'	57.4
	5'-CTGCTGCACCTCTAAGCGA-3'	57.3
<i>Dlx1</i>	5'-ATGCCAGAAAGTCTCAACAGC-3'	60.6
	5'-AACAGTGCATGGAGTAGTGCC-3'	62.4
<i>Dlx2</i>	5'-AAAGAAAGTCCGGAAACCACG-3'	60.1
	5'-TCTTCTTGAAGTGCATCGGC-3'	60.5
<i>Tuj1</i>	5'-TAGACCCCAGCGGCAACTAT-3'	58.2
	5'-GTTCCAGGTTCCAAGTCCACC-3'	58.0
<i>Gfap</i>	5'-CCCTGGCTCGTGTGGATT-3'	57.8
	5'-GACCGATACCACTCCTCTGTC-3'	56.5
<i>S100β</i>	5'-TGGTTGCCCTCATTGATGTCT-3'	61.4
	5'-CCCATCCCCATCTTCGTCC-3'	61.5
<i>Mbp</i>	5'-CTGCCGCTGTTTTGGATAATGA-3'	61.2
	5'-CATCGGGGAAGTCGAAACGG-3'	61.9
<i>Mog</i>	5'-AGCTGCTTCCTCTCCCTTCTC-3'	63.0
	5'-ACTAAAGCCCGGATGGGATAC-3'	61.1
<i>Mag</i>	5'-AATCGGCTCACAAGGGATTCA-3'	61.5
	5'-TCCTCCCAGCTTAAAGATTTGG-3'	60.2
<i>Gapdh</i>	5'-AGGTCGGTGTGAACGGATTG-3'	57.6
	5'-TGTAGACCATGTAGTTGAGGTCA-3'	55.1