Supplementary Table 1. The recovery of facial function and characteristics according to HBGS and May's classification (modified HBGS)

| May's classification | HB grading system (HBGS) | Characteristics | | | |
|----------------------|--------------------------|--|--|--|--|
| Complete | HBGS I | Normal function or complete facial function recovery | | | |
| Fair | HBGS II-III | Slight to moderate facial weakness and synkinesis | | | |
| Failed | HBGS IV-VI | Severe facial weakness, synkinesis, and contracture and/or spasm | | | |

HBGS, House-Brackmann grading system.

Supplementary Table 2. Cochrane collaboration's tool for assessing risk of bias

| Study | a | b | С | d | е | f | g |
|-----------------------|---|---|---|---|---|---|---|
| Li et al. (2016) [19] | + | - | _ | - | ? | + | ? |
| Brown (1982) [13] | + | - | - | _ | ? | + | ? |

a, random sequence generation; b, allocation concealment; c, blinding of participants and personnel; d, blinding of outcomes assessment; e, incomplete outcome data; f, selective reporting; g, other bias; +, low risk of bias; -, high risk of bias; ?, unclear risk of bias.

Supplementary Table 3. Newcastle-Ottawa scale of bias risk for the non-randomized studies

| Study | Adequate case definition | Representativeness | Selection of controls | Definition of controls | Comparability | Ascertainment of exposure | Method of ascertainment | Nonresponse rate | Score/10 |
|-------------------------------|--------------------------|--------------------|-----------------------|------------------------|---------------|---------------------------|-------------------------|------------------|----------|
| Kim et al. (2016) [8] | * | * | * | * | ** | * | * | | 8 |
| Yanagihara et al. (2001) [20] | * | * | * | * | ** | * | * | | 8 |
| Gantz et al. (1999) [21] | * | * | * | * | ** | * | * | | 8 |
| May et al. (1985) [22] | * | * | * | * | ** | * | * | | 8 |