

REFERENCES

1. Meng QX, Gao XH, Song JS, Li P, He L. Multi-factors analysis of the exposure of glottis area with suspend retaining laryngoscope. *Clin Med*. 2010 Jun;30(6):41-3.
2. Wang M, Xiao ZR, Yu JQ, Zeng RF, Tan GL. Related factors of difficult laryngeal exposure in suspension laryngoscopy under general anesthesia. *Med Innov China*. 2012 Mar;9(9):1-2.
3. Sun J, Zhang XG, Sun YH, Hu WL, Peng SD, Cui XB, et al. The related factors analysis of difficult laryngeal exposure under microlaryngoscopy. *J Inner Mongolia Med Univ*. 2015 Apr;37(2):179-81.
4. Wang J, Hu Y, Wang D, Zhao G, Li X, Li Y. The related factors analysis of difficult laryngeal exposure under retaining laryngoscope. *Lin Chuang Er Bi Yan Hou Tou Jing Wai Ke Za Zhi*. 2015 Sep;29(17):1519-21.
5. Huang C, Mo J. Study on affect factors related to laryngeal exposure in self-retaining microscopic surgery. *Chin J Otorhinolaryngol*. 2016 Aug;22(4):317-9.
6. Wa YL, Xu XL, Zhou L, Wang RQ, Zhuang PY. A study on the X-ray measurement predictors of difficult laryngeal exposure in patients undergoing microlaryngosurgery. *Lin Chuang Er Bi Yan Hou Tou Jing Wai Ke Za Zhi*. 2016 Jul;30(13):1042-6.
7. Paul RR, Varghese AM, Mathew J, Chandrasekharan R, Amalanathan S, Asif SK, et al. Difficult laryngeal exposure in microlaryngoscopy: can it be predicted preoperatively? *Indian J Otolaryngol Head Neck Surg*. 2016 Mar;68(1):65-70.
8. Jin XF, Fan GK. Analysis of the relevant factors for the difficult laryngeal exposure in patients undergoing suspension laryngoscopy. *Beijing Med J*. 2016 Apr;38(2):129-32.
9. Li JJ, Chen WX, Zhu ZF, Zhang JL, He FY, Wang YJ. Prospective study of risk factors of difficult laryngeal exposure in suspension laryngoscopy. *Lin Chuang Er Bi Yan Hou Tou Jing Wai Ke Za Zhi*. 2017 Apr;31(7):520-3.
10. Pinar E, Calli C, Oncel S, Selek B, Tatar B. Preoperative clinical prediction of difficult laryngeal exposure in suspension laryngoscopy. *Eur Arch Otorhinolaryngol*. 2009 May;266(5):699-703.
11. Liu YJ, Pang XH, Chu JS, Mao MR, Xu LY. An analysis of the related factors of difficult laryngeal exposure in microsurgical laryngoscope surgery. *China Med Pharm*. 2021 Dec;11(24):156-8.
12. Liu Y, Zhang Y, Chen Y, Yue L, Su T, Shi S. Sternum-mental angle: a new predictor of difficult laryngeal exposure in suspension microsurgery: an observational study. *Eur Ann Otorhinolaryngol Head Neck Dis*. 2022 Aug;139(4):202-7.
13. Chen FS, Zhang ZX, Chen J, Zhang L. Influencing factors of glottis exposure difficulty in laryngoscopic surgery under self-retaining laryngoscope. *Mod Instrum Med Treat*. 2019 Aug;25(1):17-20.
14. Cheng JW, Ye YH, Wu WJ, Zeng YL. Logistic analysis of influencing factors of glottic exposure difficulty in support laryngoscope. *Lingnan Mod Clin Surg*. 2020 Feb;20(1):93-7.
15. Hsiung MW, Pai L, Kang BH, Wang BL, Wong CS, Wang HW. Clinical predictors of difficult laryngeal exposure. *Laryngoscope*. 2004 Feb;114(2):358-63.
16. Wei W, Yan JH, Wang H. Analysis on related influencing factors of glottis exposure difficulty under self-retaining laryngoscope. *China Mod Med*. 2018 Dec;25(35):122-4.
17. Wang S, Wang XC, Wang YH. Influencing factors of difficult glottic exposure during low-temperature plasma resection under suspension laryngoscope in laryngeal carcinoma patients. *Clin Psychosom Dis*. 2021 May;27(3):138-41.
18. Kharrat I, Achour I, Trabelsi JJ, Trigui M, Thabet W, Mnejja M, et al. Prediction of difficulty in direct laryngoscopy. *Sci Rep*. 2022 Jun;12(1):10722.
19. Roh JL, Lee YW. Prediction of difficult laryngeal exposure in patients undergoing microlaryngosurgery. *Ann Otol Rhinol Laryngol*. 2005 Aug;114(8):614-20.