

SUPPLEMENTARY REFERENCES

1. Jaswal A, Jana AK, Sikder B, Nandi TK, Sadhukhan SK, Das A. Novel treatment of atrophic rhinitis: early results. *Eur Arch Otorhinolaryngol*. 2008 Oct;265(10):1211-7.
2. Salaheldin AH, Hussein A. Effect of platelet-rich plasma on nasal mucociliary clearance after submucous diathermy of inferior turbinate. *Egypt J Ear Nose Throat Allied Sci*. 2012 Jul;13(2):71-5.
3. Jang YJ, Kim JH, Song HY. Empty nose syndrome: radiologic findings and treatment outcomes of endonasal microplasty using cartilage implants. *Laryngoscope*. 2011 Jun;121(6):1308-12.
4. Saafan ME. Acellular dermal (alloderm) grafts versus silastic sheets implants for management of empty nose syndrome. *Eur Arch Otorhinolaryngol*. 2013 Feb;270(2):527-33.
5. Jung JH, Baguindali MA, Park JT, Jang YJ. Costal cartilage is a superior implant material than conchal cartilage in the treatment of empty nose syndrome. *Otolaryngol Head Neck Surg*. 2013 Sep;149(3):500-5.
6. Lee TJ, Fu CH, Wu CL, Lee YC, Huang CC, Chang PH, et al. Surgical outcome for empty nose syndrome: impact of implantation site. *Laryngoscope*. 2018 Mar;128(3):554-9.
7. Kim DH, Lee MH, Lee J, Song EA, Kim SW, Kim SW. Platelet-rich plasma injection in patients with atrophic rhinitis. *ORL J Otorhinolaryngol Relat Spec*. 2021;83(2):104-11.
8. Chang FY, Fu CH, Lee TJ. Outcomes of olfaction in patients with empty nose syndrome after submucosal implantation. *Am J Otolaryngol*. 2021 Jul-Aug;42(4):102989.