

Supplementary Table 2. Correlation matrix for audiological, sociodemographic, and psychometric variables in Mediterranean individuals with chronic tinnitus and unilateral SNHL (n=29)

Variable	MoCA score	THI score	GÜF score	AV.PTA	AV.4 KHz	AV.8 KHz	PHQ-9 score	Education level	Smoking status
THI score	-0.32								
GÜF score	-0.11	0.55**							
AV.PTA	-0.14	0.33	0.25						
AV.4 KHz	-0.08	-0.19	-0.34	-0.05					
AV.8 KHz	-0.10	-0.06	0.06	0.11	0.07				
PHQ-9 score	-0.20	0.54**	0.37*	-0.02	-0.25	-0.30			
Education level	0.24	-0.26	-0.31	-0.03	0.12	0.14	-0.41*		
Smoking status	-0.15	-0.22	-0.12	-0.21	0.10	0.04	-0.06	-0.05	
Age	-0.16	-0.26	-0.27	-0.66**	0.18	-0.05	0.07	-0.09	0.06

SNHL, sensorineural hearing loss; MoCA, Montreal Cognitive Assessment; THI, Tinnitus Handicap Inventory; GÜF, Hyperacusis Questionnaire; AV., average; PTA, pure tone average; PHQ-9, Patient Health Questionnaire.

* $P < 0.05$, ** $P < 0.01$.