

See “Comparison of efficacy of once daily multimatrix mesalazine 2.4 g/day and 4.8 g/day with other 5-aminosalicylic acid preparation in active ulcerative colitis: a randomized, double-blind study” on page 255.

Supplementary Information of Each Investigator

Satoshi Motoya, Sapporo-Kosei General Hospital; Tomohiro Tada, Tada Tomohiro Clinic; Toshihide Ohmori, Ohmori Toshihide Gastro-intestinal Clinic; Takehiro Arai, Tokatsu-Tsujinaka Hospital; Yukihiro Hamahata, Tsujinaka Hospital Kashiwanoha; Yasuo Suzuki, Toho University Sakura Medical Center; Toshiaki Terada, Terada Hospital; Jun Miwa, Toshiba General Hospital; Takashi Hiratsuka, Hiratsuka Gastroenterological Hospital; Yasuyoshi Takeo, Hachioji Digestive Disease Hospital; Toru Isono, Isono Clinic; Hitoshi Kaneko, Kaneko Clinic; Ryuzo Murai, Onaka Clinic; Nobuyuki Matsuhashi, NTT Medical Center Tokyo; Haruo Nishino, Matsushima Clinic; Shinichiro Suzuki, Fujisawa Shounandai Hospital; Koichiro Matsuda, Toyama Prefectural Central Hospital; Kazuhiko Kawakami, Matsuda Hospital; Mitsuki Miyata, Miyata Clinic; Akihiko Ohta, Ieda Hospital; Takayuki Yamamoto, Yokkaichi Social Insurance Hospital; Takataro Fukuhara, Hirakata Kohsai Hospital; Soken Sai, Sai Clinic; Tatsuhiko Usui, Usui Internal Medicine/Gastroenterology Department Clinic; Kentaro Tsuji, Tsuji Kentaro Clinic; Koji Sawada, Ikoma Digestive Tract Internal Medicine Clinic; Kenji Watanabe, Osaka City University Hospital; Shujiro Yazumi, the Tazuke Kofukai Medical Research Institute Kitano Hospital; Kazuhiko Nishimura, Nishimura Internal Medicine Clinic; Nobuo Aoyama, Aoyama Medical Clinic; Shiro Nakamura, Ken Fukunaga, Hyogo College of Medicine Hospital; Akira Chikuba, Chikuba Hospital for Gastrointestinal and Colorectal Surgery; Toshiaki Ochiai, Teppei Kabemura, Saiseikai Fukuoka General Hospital; Hisamitsu Hidaka, Hidaka Clinic; Keiichi Mitsuyama, Kurume University Hospital; Hirotada Akiho, Kitakyushu Municipal Medical Center; Tetsuya Ishida, Oita Red Cross Hospital; Naoto Ishikawa, Ishikawa Clinic; Naoya Shibata, Michihisa Fuchimoto, Ikime Coloproctological Clinic; Yukinori Sameshima, Sameshima Hospital; Takanori Kanai, Keio University Hospital; Naoki Yoshimura, Social Insurance Central General Hospital; Kiyonori Kobayashi, Kitasato University East Hospital; Takafumi Ando, Nagoya University Hospital; Tsuguhiro Kimura, Fujita Gastroenterological Hospital; Takashi Hisabe, Fukuoka University Chikushi Hospital; Toshiro Sugiyama, Toyama University Hospital; Hidenori Kurakata, Inoue Clinic; Shinichi Ogata, Saga Prefectural Medical Centre Koseikan; Masakazu Nagahori, Medical Hospital, Tokyo Medical and Dental University; Hirozumi Obata, Obata Medical Clinic; Yoshimasa Otani, Sagamidai Hospital; Hiroyuki Kurihara, Tokorozawa Proctologic Hospital; Ryoichi Suzuki, Kannai-Suzuki Clinic; Haruo Nishino, Matsushima Clinic.