

Supplementary Table 2. Comparison of Pediatric Inflammatory Bowel Disease Data with French Population-Based Registry (EPIMAD)

| Variable | VEOIBD, % (No./No.) | | | LO-PIBD, % (No./No.) | | |
|--------------------------|---------------------|----------------------|---------|----------------------|----------------------|---------|
| | India | EPIMAD ³¹ | P-value | India | EPIMAD ³¹ | P-value |
| No. of patients (%) | 22 (7.5) | 42 (3) | 0.003 | 270 (92.5) | 1,370 (97) | |
| CD:UC:IBD-U (%) | 59.1:36.4:4.5 | 60:33:7 | 0.90 | 60:39:1 | 73.5:24.0:2.5 | <0.001 |
| Male sex | 82 (18/22) | 52 (22/42) | 0.02 | 61 (165/270) | 52 (708/1,370) | 0.004 |
| Diagnostic delay > 6 mo | 40.9 (9/22) | 27 (11/42) | 0.23 | 78.8 (213/270) | 30 (407/1,370) | <0.001 |
| Family history of IBD | 13.6 (3/22) | 9.5 (4/42) | 1 | 9.2 (25/270) | 15 (210/1,370) | 0.009 |
| Diarrhea | 77.2 (17/22) | 76 (32/42) | 0.92 | 68.1 (184/270) | 66 (899/1,370) | 0.03 |
| Blood in stool | 68.2 (15/22) | 81 (34/42) | 0.25 | 52.6 (142/270) | 46 (624/1,370) | <0.001 |
| Mucus in stool | 50 (11/22) | 40 (17/42) | 0.47 | 33.7 (91/270) | 21 (281/1,370) | <0.001 |
| Weight loss | 36.4 (8/22) | 21 (9/42) | 0.19 | 50.37 (136/270) | 49 (670/1,370) | 0.66 |
| Abdominal pain | 50 (11/22) | 43 (18/42) | 0.58 | 67.04 (181/270) | 74 (1,013/1,370) | 0.15 |
| EIM | 22.7 (5/22) | 17 (7/42) | 0.74 | 21.8 (59/270) | 17 (231/1,370) | <0.049 |
| Perianal disease | 7.69 (1/13) | 8 (2/25) | | 7.4 (13/162) | 6 (59/1,032) | |
| Crohn's disease | | | | | | |
| Bleeding per rectum | 53.8 (7/13) | 68 (17/25) | 0.48 | 31.5 (51/162) | 30 (303/1,007) | 0.72 |
| Abdominal pain | 46.2 (6/13) | 48 (12/25) | 0.92 | 74.7 (121/162) | 74 (809/1,007) | 0.09 |
| Weight loss | 38.5 (5/13) | 20 (5/25) | 0.26 | 53.1 (86/162) | 56 (566/1,007) | 0.46 |
| EIM | 23.1 (3/13) | 20 (5/25) | 0.57 | 19.8 (32/162) | 21 (208/1,007) | 0.79 |
| Colonic disease L2 | 30.8 (4/13) | 39 (9/25) | 0.52 | 37.7 (61/162) | 14 (128/1,007) | <0.001 |
| B2+B3 disease in CD | 30.8 (4/13) | 13 (3/25) | 0.20 | 43.8 (71/162) | 22 (208/1,007) | <0.001 |
| Ulcerative colitis | | | | | | |
| Abdominal pain | 62.5 (5/8) | 21 (3/14) | 0.08 | 56.2 (59/105) | 55 (180/329) | 0.79 |
| EIMs | 25 (2/8) | 7 (1/14) | 0.53 | 25.7 (27/105) | 6 (19/329) | <0.001 |
| Proctitis (E1) | 25 (2/8) | 9 (1/11) | 0.32 | 11.4 (12/105) | 30 (92/303) | <0.001 |
| Procto-sigmoiditis (E2) | 12.5 (1/8) | 36 (4/11) | | 15.2 (16/105) | 26 (79/303) | |
| Upto hepatic flexure(E3) | 0 (0/8) | 27 (3/11) | | 22.9 (24/105) | 10 (30/303) | |
| Pancolitis (E4) | 62.5 (5/8) | 27 (3/11) | | 50.5 (53/105) | 33 (102/303) | |

EPIMAD, epidemiology of inflammatory bowel diseases: new insights from a French population-based registry; IBD, inflammatory bowel disease; VEOIBD, very early onset IBD; LO-PIBD, later onset pediatric IBD; CD, Crohn's disease; UC, ulcerative colitis; IBD-U, IBD undifferentiated; EIM, extraintestinal manifestations; L2, colonic CD; B2, stricturing disease; B3, fistulizing CD.