

Supplementary Table 2. Laboratory Data before and after ZAH Administration in CD Patients

| Variable | At 4 weeks | | | At 20 weeks | | |
|-----------------------------------|----------------------|----------------------|---------|----------------------|----------------------|---------|
| | Day 1 | 4 Weeks | P-value | Day 1 | 20 Weeks | P-value |
| WBC (/μL) | 6,475 (3,350–13,950) | 6,850 (4,110–13,750) | 0.367 | 6,430 (3,110–13,950) | 6,190 (2,680–16,200) | 0.634 |
| Hb (g/dL) | 12.6 (8.5–15.1) | 12.4 (8.0–16.1) | 0.702 | 12.5 (8.5–15.1) | 12.3 (9.2–15.5) | 0.593 |
| PLT ($\times 10^4/\mu\text{L}$) | 28.4 (11.7–42.7) | 27.9 (10.1–49.5) | 0.965 | 27.8 (11.7–41.1) | 28.6 (13.7–43.2) | 0.249 |
| CRP (mg/dL) | 0.31 (0.01–12.82) | 0.12 (0.01–9.61) | 0.213 | 0.18 (0.02–12.82) | 0.25 (0.01–5.44) | 0.182 |
| ALB (g/dL) | 3.4 (1.8–4.8) | 3.6 (1.9–4.4) | 0.141 | 3.5 (1.8–4.8) | 3.7 (2.9–4.6) | 0.105 |

Values are presented as median (range).

ZAH, zinc acetate hydrate; CD, Crohn's disease; WBC, white blood cell; Hb, hemoglobin; PLT, platelet; CRP, C-reactive protein; ALB, albumin.

P-values were calculated using the Wilcoxon matched-pairs signed-rank test for continuous variables.