

Supplementary Table 7. Comparative Analysis of Endoscopic Phenotype between Patients with and without Chronic Pouchitis (n = 50)

Variable	Category	Chronic pouchitis (-) (n = 26)	Chronic pouchitis (+) (n = 24)	P-value ^a
Normal	Yes	0	0	-
	No	26 (100)	24 (100)	
Afferent limb involvement	Yes	4 (15.4)	7 (29.2)	0.282
	No	13 (50.0)	9 (37.5)	
	NA	9 (34.6)	8 (33.3)	
Inlet limb involvement	Yes	7 (26.9)	10 (41.7)	0.523
	No	13 (50.0)	10 (41.7)	
	NA	6 (23.1)	4 (16.7)	
Diffuse inflammation	Yes	1 (3.8)	10 (41.7)	0.002
	No	25 (96.2)	14 (58.3)	
Focal inflammation	Yes	25 (96.2)	14 (58.3)	0.002
	No	1 (3.8)	10 (41.7)	
Cuffitis	Yes	19 (73.1)	17 (70.8)	1.000
	No	7 (26.9)	7 (29.2)	
Pouch fistula	Yes	0	7 (29.2)	0.003
	No	26 (100)	17 (70.8)	
Diffuse inflammation and/or pouch fistula	Yes	1 (3.8)	12 (50.0)	<0.001
	No	25 (96.2)	12 (50.0)	

Values are presented as number (%).

^aFisher exact test.