

Supplementary Table 2. Age of Onset, Onset-to-Diagnosis Interval, Longest Hospital Stays, Frequency of Hospitalization and Diagnosis-to-Surgery Interval for the 3 Cohorts

| | All periods (2007–2021) | <i>P</i> -value ^a | Cohort I (2007–2011) | <i>P</i> -value ^b | Cohort II (2012–2016) | <i>P</i> -value ^c | Cohort III (2017–2021) |
|---------------------------------------|----------------------------|------------------------------|-------------------------|------------------------------|--------------------------|------------------------------|---------------------------|
| Age at diagnosis (yr) | 39 (29–51) | 0.843 | 37 (29–45) | 0.580 | 39 (29–50) | 0.721 | 39 (29–53) |
| Interval from onset to diagnosis (mo) | 1 (1–12) | 0.033 | 2 (1–8) | 0.086 | 3 (1–15) | 0.028 | 2 (1–11) |
| Frequency of hospitalization (n) | 1 (1–3) | 0.019 | 2 (1–4) | 1.000 | 2 (1–4) | 0.026 | 1 (1–3) |
| Longest hospital stays (day) | 11 (7–19) | <0.001 | 13 (7–22) | 0.999 | 14 (8–22) | <0.001 | 9 (6–15) |
| Diagnosis-to-surgery interval (mo) | 66 (36–96) | <0.001 | 144 (132–157) | <0.001 | 87 (72–103) | <0.001 | 36 (22–51) |

Values are presented as median (interquartile range). All comparisons are performed by one-way analysis of variance, both overall and pairwise.

^aOverall comparison of all 3 study periods.

^bPairwise comparison of cohort i (2007–2011) and cohort II (2012–2016) (Bonferroni correction was used to adjust for multiple comparisons).

^cPairwise comparison of cohort ii (2012–2016) and cohort III (2017–2021) (Bonferroni correction was used to adjust for multiple comparisons).