

Supplementary Table 2. Patient Background Characteristics by Whether Bowel Urgency Resolved

Characteristic	Bowel urgency not resolved (n = 22)	Bowel urgency resolved (n = 31)	P-value
Sex			0.780 ^a
Male	9 (40.9)	15 (48.4)	
Female	13 (59.1)	16 (51.6)	
Age (yr), median (IQR)	42.5 (34.0–49.0)	42.0 (31.0–58.0)	0.545 ^b
Height (cm), median (IQR)	163.8 (160.0–168.0)	170.0 (156.0–173.0)	0.481 ^b
Body weight (kg), median (IQR)	56.7 (52.0–63.0)	59.0 (47.0–69.0)	0.588 ^b
Partial Mayo score, median (IQR)	5.0 (3.25–6.0)	5.0 (4.0–5.5)	0.627 ^b
Smoking habit			1.000 ^a
No	20 (90.9)	29 (93.5)	
Yes	2 (9.1)	2 (6.5)	
Duration of illness			0.254 ^a
< 12 wk	1 (4.5)	1 (3.2)	
12 wk to < 1 yr	5 (22.7)	2 (6.5)	
1 yr to < 5 yr	5 (22.7)	6 (19.4)	
≥ 5 yr	11 (50.0)	22 (71.0)	
First-onset and relapse in the active phase			0.638 ^a
Initial	3 (13.6)	2 (6.5)	
Relapse	19 (86.4)	29 (93.5)	
Previous medications (within 8 wk before the start of observation)			
5-ASA rectal formulation			0.638 ^a
No	19 (86.4)	29 (93.5)	
Yes	3 (13.6)	2 (6.5)	
5-ASA suppository formulation			0.505 ^a
No	22 (100.0)	29 (93.5)	
Yes	0	2 (6.5)	
5-ASA oral formulation			0.131 ^a
No	2 (9.1)	3 (9.7)	
< 3,600 mg/day	3 (13.6)	6 (19.4)	
3,600 to < 4,800 mg/day	10 (45.5)	5 (16.1)	
4,800 mg/day	7 (31.8)	17 (54.8)	
Immunomodulator			1.000 ^a
No	20 (90.9)	29 (93.5)	
Yes	2 (9.1)	2 (6.5)	
Disease pattern			0.026 ^c
Proctitis	5 (22.7)	0	
Left-sided colitis	8 (36.4)	16 (51.6)	
Pancolitis	9 (40.9)	15 (48.4)	
Disease severity			0.759 ^a
Mild	7 (31.8)	8 (25.8)	
Moderate	15 (68.2)	23 (74.2)	
Severe	0	0	

(Continued to the next page)

Supplementary Table 2. Continued

Characteristic	Bowel urgency not resolved (n = 22)	Bowel urgency resolved (n = 31)	P-value
Active lesion site			0.398 ^a
Rectum only	5 (22.7)	4 (18.2)	
Up to the sigmoid colon	7 (31.8)	13 (59.1)	
Up to the descending colon	4 (18.2)	2 (9.1)	
From the descending colon to the mouth	6 (27.3)	3 (13.6)	
No. of defecations per day (normal frequency)			0.386 ^b
No.	22	31	
Median (IQR)	1.0 (1.0–2.0)	2.0 (1.0–3.0)	
History of administration of budesonide rectal foam			0.753 ^a
No	16 (72.7)	24 (77.4)	
Yes	6 (27.3)	7 (22.6)	

Values are presented as number (%) unless otherwise indicated.

^aFisher exact test.

^bWilcoxon rank sum test.

IQR, interquartile range; 5-ASA, 5-aminosalicylic acid.