
<Clinical High Risk for Psychosis>

A-1 0.5-1 year


Study	Mean	SD	Total	Mean	TAU		Weight	Mean Difference IV, Random, 95% C		Mean Difference IV, Random, 95% CI				1
Amminger 2010	17.7	2.3	41	7.2	2.3	40	30.9%	10.50 [9.50	, 11.50]					-
Addington 2011	5.1	13.8	19	2.7	10.6	16	20.0%	2.40 [-5.69,	10.49]					
Morrison 2012	8.3	14.3	97	10.5	13.3	98	27.6%	-2.14 [-6.02	, 1.74]		-	-		
McGorry 2013	10.3	11.5	26	7.0	12.5	18	21.5%	3.30 [-3.97,	10.57]		100			_
Total (95% CI)			183			172	100.0%	3.84 [-2.20,	9.89]		ī	-		_
Heterogeneity: Tau	$u^2 = 30.5$	21; Ch	$ni^2 = 44$	11, df =	3 (P <	.01); I2	= 93%		-		1	l.	1	- 1
Test for overall effe					110	"				-10	-5	0	5	10

A-2 1.5-2 years




<Recent Onset Psychosis>


B-1 0.5-1 year



B-2 1.5-2 years



B-3 5 years



Supplementary Fig. 6. Mean difference of GAF score change between early intervention and treatment-as-usual groups over different follow-up periods.