
A 1 year

Study van der Gaag 2012	Mean 10.4	SD 10.3		BEG VA	TAU SD	76.250		Mean Difference IV, Random, 95% C		Mean Difference IV, Random, 95% CI					ı
								-1.00 [-4.64, 2.64]		-					
McGorry 2017	14.7	19.1	153	14.3	16.8	151	1.7%	0.40 [-3.64, 4.44]		-				_	
McGorry 2023	3.4	1.4	56	4.0	1.5	57	95.1%	-0.60 [-1.14, -0.06]			- 1				
Bechdolft 2023	13.3	16.3	96	15.7	14.5	55	1.1%	-2.40 [-7.44, 2.64]	-		•		_		
Total (95% CI)			376			332	100.0%	-0.61 [-1.14, -0.08]				•			
Heterogeneity: Tau ² = 0 Test for overall effect: 2	0; Chi ² = 2 = -2.27	0.77, (P = .	df = 3 (1 02)	P = .86)	$ \mathbf{l}^2 = 0$	%		•	-6	-4	-2	0	2	4	6

B 3-4 years



Supplementary Fig. 7. Mean difference of SOFAS score changes between early intervention and treatment as usual group over difference periods (1 year, 3-4 years) in clinical high risk for psychosis.