

Supplementary Table 1. Baseline characteristics among propensity score matched population

Variable	HLA one-way mismatch (n=14)	Control (n=70)	p-value
Age (yr)	52 (40–61)	52 (44–55)	0.727
Sex, female	4 (28.6)	12 (17.1)	0.534
BMI (kg/m²)	22.9±3.1	23.1±3.9	0.849
Diabetes mellitus	3 (21.4)	15 (21.4)	0.989
Underlying liver disease			0.860
Hepatitis B	6 (42.9)	43 (61.4)	
Hepatitis C	2 (14.3)	5 (7.1)	
Alcoholic	2 (14.3)	5 (7.1)	
Autoimmune liver disease	0 (0)	1 (1.4)	
Acute liver failure	1 (7.1)	3 (4.3)	
Biliary atresia	2 (14.3)	9 (12.9)	
Others	1 (7.1)	4 (5.7)	
HCC	9 (64.3)	47 (67.1)	0.876
Pretransplant MELD	10 (7–14)	9 (7–14)	0.909
Donor relation			0.214
Living related	12 (85.7)	44 (62.9)	
Living unrelated	0 (0)	7 (10.0)	
Deceased	2 (14.3)	19 (27.1)	
Donor age (yr)	28 (25–34)	28 (22–34)	0.782
Donor sex, female	5 (35.7)	23 (32.9)	0.974

Values are presented as median (interquartile range) or number (%).

HLA, human leukocyte antigen; BMI, body mass index; HCC, hepatocellular carcinoma; MELD, model for end-stage liver disease.