

SUPPLEMENTARY MATERIALS

Table S1. Change in MAS scores for the wrist, elbow, and finger flexor muscles at 4, 8, and 12 weeks after injection in the per-protocol set

	Wrist flexor MAS			Elbow flexor MAS			Finger flexor MAS		
	n	mean±SD	p-value ^{a)}	n	mean±SD	p-value ^{a)}	n	mean±SD	p-value ^{a)}
Baseline	151	1.95±0.75		151	1.78±0.91		151	2.21±0.86	
Week 4	151	0.98±0.64		151	1.01±0.70		151	1.12±0.71	
Change from baseline	151	-0.97±0.70	<0.001***	151	-0.77±0.68	<0.001***	151	-1.09±0.75	<0.001***
Week 8	151	1.05±0.68		151	1.09±0.69		151	1.23±0.77	
Change from baseline	151	-0.90±0.68	<0.001***	151	-0.69±0.75	<0.001***	151	-0.98±0.76	<0.001***
Week 12	151	1.16±0.72		151	1.20±0.69		151	1.41±0.78	
Change from baseline	151	-0.79±0.68	<0.001***	151	-0.58±0.73	<0.001***	151	-0.79±0.77	<0.001***

MAS, Modified Ashworth Scale; SD, standard deviation.

Friedman test: wrist flexor MAS, p<0.001; elbow flexor MAS, p<0.001; finger flexor MAS, p<0.001.

^{a)}Post-hoc Wilcoxon signed-rank test adjusted by Bonferroni correction, ***p<0.001.