

Supplementary Table 1. Characteristics of patients who underwent cardiac stress testing after referral for deceased donor kidney transplant waitlist placement according to the presence or absence of significant coronary artery disease on coronary angiography

Characteristic	Total (n=63)	CAD (n=27)	No CAD (n=36)	P-value
Age (yr)	53 (44–58)	56 (49–60)	48 (40–58)	0.01
Male	42 (66.7)	22 (81.5)	20 (55.6)	0.04
Ethnicity				0.23
Chinese	42 (66.7)	18 (66.7)	24 (66.7)	
Malay	19 (30.2)	7 (25.9)	12 (33.3)	
Indian	2 (3.2)	2 (7.4)	0	
Others	0	0	0	
Cause of ESKD				0.41
Glomerulonephritis	16 (25.4)	6 (22.2)	10 (27.8)	
Diabetes mellitus	45 (71.4)	19 (70.4)	26 (72.2)	
Hypertension	1 (1.6)	1 (3.7)	0	
ADPKD	0	0	0	
Others	1 (1.6)	1 (3.7)	0	
Dialysis modality				0.75
Hemodialysis	52 (82.5)	23 (85.2)	29 (80.6)	
Peritoneal dialysis	11 (17.5)	4 (14.8)	7 (19.4)	
Dialysis vintage (yr)	1.8 (0.5–5.9)	1.3 (0.3–6.3)	2.9 (0.7–4.8)	0.35
Hypertension	59 (93.7)	26 (96.3)	33 (91.7)	0.63
Diabetes mellitus	49 (77.8)	21 (77.8)	28 (77.8)	1.00
Dyslipidemia	47 (74.6)	17 (63.0)	30 (83.3)	0.08
Current or past smoking	21 (33.3)	11 (40.7)	10 (27.8)	0.30
History of CAD	6 (9.5)	5 (18.5)	1 (2.8)	0.08
History of stroke	0	0	0	NA
Ejection fraction <50%	10 (16.1)	5 (18.5)	5 (14.3)	0.74
Ischemia on the cardiac stress test	34 (54.0)	16 (59.3)	18 (50.0)	0.61

Values are presented as median (interquartile range) or number (%).

CAD, coronary artery disease; ESKD, end-stage kidney disease; ADPKD, autosomal dominant polycystic kidney disease; NA, not applicable.