

Supplementary Table 1. Baseline characteristics of included primary ICH patients and excluded primary ICH patients

Characteristic	Data complete (% of total)	Total group (n=235)	Included primary ICH patients (n=139)	Excluded primary ICH patients* (n=96)	<i>P</i>
Demographic					
Age (yr)	235 (100)	77.0 (68.0–83.0)	76.0 (65.0–82.0)	78.5 (70.3–84.8)	0.024
Male sex	235 (100)	133 (56.6)	88 (63.3)	45 (46.9)	0.012
Clinical					
History of hypertension	233 (99.1)	153 (65.1)	91 (65.5)	62 (64.6)	0.938
Hypercholesterolemia	231 (98.3)	93 (40.3)	54 (38.8)	39 (40.6)	0.670
Anticoagulant use	235 (100)	62 (26.4)	33 (23.7)	29 (30.2)	0.269
Glucose (mmol/L)	226 (96.2)	7.2 (6.0–8.9)	6.7 (5.9–8.5)	7.9 (6.5–9.7)	0.003
GCS	235 (100)	14 (10–15)	15 (13–15)	10 (5–14)	<0.001
mRS after 3 months	220 (93.6)	6 (3–6)	4 (2–6)	6 (4–6)	<0.001
Imaging					
Infratentorial ICH	235 (100)	20 (8.5)	9 (6.5)	11 (11.5)	0.178
Baseline ICH volume (mL)	234 (99.6)	16.6 (6.5–48.5)	12.0 (5.7–27.1)	36.0 (8.8–127.1)	<0.001
Presence of IVH on first CT-scan	235 (100)	103 (43.8)	47 (33.8)	56 (58.3)	<0.001

Values are presented as number (%) or median (interquartile range).

ICH, intracerebral hemorrhage; GCS, Glasgow Coma Scale; mRS, modified Rankin Scale; IVH; intraventricular hemorrhage; CT, computed tomography.

*Patients lacking dual-energy computed tomography angiography (DE-CTA) and patients in whom DE-CTA was performed >3 hours after baseline CT.